

1. Baroni MG, Alcolado JC, Needham EW, Pozzilli P, Stocks J, Galton DJ. Sib-pair analysis of adenosine deaminase locus in NIDDM. *Diabetes*. 41(12):1640-3, 1992.
2. Baroni MG, Alcolado JC, Pozzilli P, Cavallo MG, Li SR, Galton DJ. Polymorphisms at the GLUT2 (beta-cell/liver) glucose transporter gene and non-insulin-dependent diabetes mellitus (NIDDM): analysis in affected pedigree members. *Clin Genet*. 41(5):229-34, 1992.
3. Baroni MG, Oelbaum RS, Pozzilli P, Stocks J, Li SR, Fiore V, Galton DJ. Polymorphisms at the GLUT1 (HepG2) and GLUT4 (muscle/adipocyte) glucose transporter genes and non-insulin-dependent diabetes mellitus (NIDDM). *HumGenet*. 88(5):557-61, 1992.
4. Pozzilli P, Buzzetti R, Bottazzo GF, Tosi R. The selection of control subjects for case/control analysis of susceptibility to type 1 (insulin-dependent) diabetes mellitus. *Diabetologia*.;36(11):1208-9, 1993.
5. Imai Y, Fusco A, Suzuki Y, Lesniak M, D'Alfonso R, Sesti G, Bertoli A, Lauro R, Accili D, Taylor SI. Variant sequences of insulin receptor substrate-1 in patients with noninsulin-dependent diabetes mellitus. *J Clin Endocrinol Metab* 79: 1655-1658, 1994.
6. Pujia A, Gnasso A, Irace C, Dominijanni A, Zingone A, Perrotti N, Colonna A, Mattioli PL. Association between ACE-D/D polymorphism and hypertension in type II diabetic subjects. *J Hum Hypertens* 8:687-91, 1994.
7. Baroni MG, Alcolado JC, Gragnoli C, Franciosi AM, Cavallo MG, Fiore V, Pozzilli P, Galton DJ. Affected sib-pair analysis of the GLUT1 glucose transporter gene locus in non-insulin-dependent diabetes mellitus (NIDDM): evidence for no linkage. *HumGenet*. 93(6):675-80, 1994.
8. Pozzilli P, Buzzetti R, Nisticó L, Romiti A, Giovannini C. Non-HLA genes and the susceptibility to insulin dependent diabetes: the role of the CTLA-4 gene. *Acta Diabetol*. ;33(4):250-2. 1996.
9. Nisticò L, Buzzetti R, Pritchard LE, Van der Auwera B, Giovannini C, Bosi E, Larrad MT, Rios MS, Chow CC, Cockram CS, Jacobs K, Mijovic C, Bain SC, Barnett AH, Vandewalle CL, Schuit F, Gorus FK, Tosi R, Pozzilli P, Todd JA. The CTLA-4 gene region of chromosome 2q33 is linked to, and associated with, type 1 diabetes. *Belgian Diabetes Registry. Hum Mol Genet*. 5(7):1075-80, 1996.
10. Luo DF, Buzzetti R, Rotter JI, Maclaren NK, Raffel LJ, Nisticò L, Giovannini C, Pozzilli P, Thomson G, She JX. Confirmation of three susceptibility genes to

insulin-dependent diabetes mellitus: IDDM4, IDDM5 and IDDM8. *Hum Mol Genet.* 1996.

11. Esposito DL., Mammarella S., Ranieri A., Della Loggia F., Capani F., Consoli A., Mariani-Costantini R., Caramia FG., Cama A., Battista: Deletion of gly723 in the insulin receptor substrate-1 of a patient with non insulin dependent diabetes mellitus. *Hum Mutat*,7:364-366, 1996.
12. Merriman T, Twells R, Merriman M, Eaves I, Cox R, Cucca F, McKinney P, Shield J, Baum D, Bosi E, Pozzilli P, Nisticò L, Buzzetti R, Joner G, Rønningen KS, Thorsby E, Undlien D, Pociot F, Nerup J, Bain S, Barnett A, Todd J. Evidence by allelic association-dependent methods for a type 1 diabetes polygene (IDDM6) on chromosome 18q21. *Hum Mol Genet.* Jul;6(7):1003-10, 1997 *Erratum in: Hum Mol Genet.* 2007 Dec 15;16(24):3197.
13. Imai Y, Neubert P, Sesti G, Accili D, Taylor SI. Expression of variant forms of insulin receptor substrate-1 identified in patients with noninsulin-dependent diabetes mellitus. *J Clin Endocrinol Metab* 82: 4201-4207, 1997.
14. Beguinot F, Haring H, Kiess H, Sesti G, Smith U, Van Obberghen E. The search for NIDDM gene project: A European Community programme aimed to facilitate the identification of NIDDM genes. *Diabetologia* 41 (1 Suppl): 15-16, 1998.
15. Baroni MG, D'Andrea MP, Capici F, Buzzetti R, Cavallo MG, Fallucca F, Giovannini C, Pozzilli P. High frequency of polymorphism but no mutations found in the GLUT1 glucose transporter gene in NIDDM and familial obesity by SSCP analysis. *HumGenet.* Apr;102(4):479-82, 1998.
16. Esposito L, Hill NJ, Pritchard LE, Cucca F, Muxworthy C, Merriman ME, Wilson A, Julier C, Delepine M, Tuomilehto J, Tuomilehto-Wolf E, Ionesco-Tirgoviste C, Nistico' L, Buzzetti R, Pozzilli P, Ferrari M, Bosi E, Pociot F, Nerup J, Bain SC, Todd JA. Genetic analysis of chromosome 2 in type 1 diabetes: analysis of putative loci IDDM7, IDDM12, and IDDM13 and candidate genes NRAMP1 and IA-2 and the interleukin-1 gene cluster. *IMDIAB Group. Diabetes.* 47(11):1797-9, 1998.
17. Nakagawa Y, Kawaguchi Y, Twells RC, Muxworthy C, Hunter KM, Wilson A, Merriman ME, Cox RD, Merriman T, Cucca F, McKinney PA, Shield JP, Tuomilehto J, Tuomilehto-Wolf E, Ionesco-Tirgoviste C, Nisticò L, Buzzetti R, Pozzilli P, Joner G, Thorsby E, Undlien DE, Pociot F, Nerup J, Rønningen KS, Bain SC, Todd JA; Bart's-Oxford Family Study Group. Fine mapping of the diabetes-susceptibility locus, IDDM4, on chromosome 11q13. *Am J Hum Genet.* 1998.

18. Shah VO, Scavini M, Nikolic J, Sun Y, Vai S, Griffith JK, Dorin RI, Stidley C, Jacoub M, Vander Jagt DL, Eaton RP, Zager PG: Z-2 microsatellite allele is linked to increased expression of the aldose reductase gene in diabetic nephropathy. *J Clin Endocrinol Metab* 83: 2886-2891, 1998.
19. Porzio O, Federici M, Hribal ML, Lauro D, Accili D, Lauro R, Borboni P, Sesti G. The Gly(972)-->Arg amino acid polymorphism in IRS-1 impairs insulin secretion in pancreatic beta cells. *J Clin Invest* 104: 357-364, 1999.
20. Mancini FP, Vaccaro O, Sabatino L, Tufano A, Rivellese A A, Riccardi G, Colantuoni V. Pro12Ala substitution in the peroxisome proliferator-activated receptor-gamma2 is not associated with type 2 diabetes. *DIABETES*, vol. 48, p. 1466-1468, 1999.
21. Pizzuti A., Frittitta L., Argiolas A., Baratta R., Goldfine ID., Bozzali M., Ercolino T., Scarlato G., Iacoviello L., Vigneri R., Tassi V., Trischitta V. A polymorphism (K121Q) of the human glycoprotein PC-1 gene coding region is strongly associated with insulin resistance. *Diabetes*, 48: 1881-1884, 1999.
22. Baroni MG, D'Andrea MP, Montali A, Pannitteri G, Barillà F, Campagna F, Mazzei E, Lovari S, Seccareccia F, Campa PP, Ricci G, Pozzilli P, Urbinati G, Arca M. A common mutation of the insulin receptor substrate-1 gene is a risk factor for coronary artery disease. *Arterioscler Thromb Vasc Biol.* 1999.
23. Hribal ML, Federici M, Porzio O, Lauro D, Borboni P, Accili D, Lauro R, Sesti G. The Gly->Arg⁹⁷² amino acid polymorphism in IRS-1 affects glucose metabolism in skeletal muscle cells. *J. Clin. Endocrinol. Metab.* 85: 2004-2013, 2000.
24. Sesti G. Insulin receptor substrates (IRSs) polymorphisms and type 2 diabetes mellitus. *Pharmacogenomics* 1: 343-357, 2000.
25. De Cosmo S., Argiolas A., Miscio G., Thomas S., Piras GP, Trevisan R., Cavallo Perin P., Bacci S., Zucaro L., Margaglione M., Frittitta L., Pizzuti A., Tassi V., Viberti GC, Trischitta V. A PC-1 aminoacid variant (K121Q) is associated with faster progression of renal disease in patients with type 1 diabetes and albuminuria. *Diabetes* 49:521-524, 2000.
26. Rasmussen S.K., Urhammer S.A., Pizzuti A, Echwald S.M., Ekstrom C.T., Hansen L., Hansen T., Borch-Johnsen K., Frittitta L., Trischitta V., Pedersen O. The K121Q variant of human PC-1 gene is not associated with insulin resistance or type 2 diabetes mellitus among Danish Caucasians. *Diabetes*, 49:1601-1603, 2000.

27. . Gu H.F, Almgren P., Lindholm E., Frittitta L., Pizzuti A., Trischitta V, Groop L.C. Association between the human glycoprotein PC-1 gene and elevated glucose and insulin levels in a paired-sibling analysis. *Diabetes*, 49:1608-1611, 2000.
28. Arcaro G., Solini A., Monauni T., Cretti A., Brunato B., Lechi A., Fellin R., Caputo M., Cocco C., Bonora E., Muggeo M., Bonadonna R.C. ACE genotype and endothelium-dependent vasodilation of conduit arteries and forearm microcirculation in humans. *Arterioscl. Thromb. Vasc. Biol.* 21: 1313-1319, 2001.
29. Federici M, Hribal ML, Ranalli M, Marselli L, Porzio O, Lauro D, Borboni P, Lauro R, Marchetti P, Melino G, Sesti G. The common Arg⁹⁷² polymorphism in insulin receptor substrate-1 causes apoptosis of human pancreatic islets. *FASEB J* 15: 22-24, 2001.
30. Accili D, Kido Y, Nakae J, Lauro D, Park BC. Genetics of type 2 diabetes: insight from targeted mouse mutants. *Curr Mol Med.* 1(1):9-23, 2001.
31. Costanzo B., Trischitta V., Di Paola R., Spampinato D., Pizzuti A., Vigneri R., Frittitta L. The Q allele variant (GLN121) of membrane glycoprotein PC-1 interacts with the insulin receptor and inhibits insulin signaling more effectively than the common K allele variant (LYS121). *Diabetes*, 50:831-836, 2001.
32. Frittitta L., Ercolino T., Bozzali M, Argiolas A., Graci S, Santagati M.G, Spampinato D, Di Paola R., Cisternino C., Tassi V., Vigneri R., Pizzuti A., Trischitta V. A cluster of 3 single nucleotide polymorphisms in the 3'-untranslated region of human glycoprotein PC-1 gene stabilizes mRNA and associates with increased protein content and insulin resistance related abnormalities. *Diabetes*, 50:1952-1955, 2001.
33. Paolisso G, Manzella D, Tagliamonte MR, Barbieri M, Marfella R, Zito G, Bonafè M, Giugliano D, Franceschi C, Varricchio M. The BB-paraoxonase genotype is associated with impaired brachial reactivity after acute hypertriglyceridemia in healthy subjects. *J Clin Endocrinol Metab.* 86(3):1078-82, 2001.
34. Frittitta L., Baratta R., Spampinato D., Di Paola R., Pizzuti A., Vigneri R., Trischitta V. The Q121 PC-1 variant and obesity have additive and independent effects in causing insulin resistance. *J Clin Endocrinol Metab*, 86: 5888-5891, 2001.
35. Di Paola R., Frittitta L., Miscio G., Bozali M., Baratta R., Centra M., Spampinato D, Santagati MG, Ercolino T., Cisternino C., Soccio T., Mastroianno S., Tassi V., Almgren P, Pizzuti A., Vigneri R., Trischitta V. A variation in 3'UTR of hPTP1B

increases specific gene expression and associates with insulin resistance. *Am J Hum Genet*, 70: 806-812, 2002.

36. Bugawan TL, Klitz W, Alejandrino M, Ching J, Pabello A, Solfelix CM, Petrone A, Buzzetti R, Pozzilli P, Erlich HA. The association of specific HLA class I and II alleles with type 1 diabetes among Filipinos. *Tissue Antigens*. 59(6):452-69, 2002.
37. Petrone A, Battelino T, Krzisnik C, Bugawan T, Erlich H, Di Mario U, Pozzilli P, Buzzetti R. Similar incidence of type 1 diabetes in two ethnically different populations (Italy and Slovenia) is sustained by similar HLA susceptible/protective haplotype frequencies. *Tissue Antigens*. 60(3):244-53, 2002.
38. Kiechl S., Lorenz E., Reindl M., Wiedermann C.J., Oberhollenzer F., Bonora E., Willeit J., Schwartz D.A. Toll-like receptor 4 polymorphisms and atherogenesis in humans. *N. Engl. J. Med.* 347: 185-192, 2002.
39. Klitz W, Bugawan TL, Pabello A, Solfelix CM, Buzzetti R, Pozzilli P, Steiner L, Alejandrino M, Erlich HA. Association of CTLA-4 variation with type I diabetes in Filipinos. *Immunogenetics*. 54(5):310-3, 2002.
40. De Cosmo S., Tassi V., Thomas S., Piras GP, Trevisan R., Cavallo Perin P., Bacci S., Zucaro L., Cisternino C., Trischitta V, Viberti GC. The Decorin gene 179 allelic variant is associated with a slower progression of renal disease in patients with type 1 diabetes. *Nephron*, 92: 72-76, 2002.
41. De Cosmo S., Miscio G., Zucaro L, Margaglione M., Argiolas A, Thomas S., Piras G., Trevisan R., Cavallo Perin P., Bacci S., Frittitta L., Pizzuti A., Tassi V., Di Minno G., Viberti G., Trischitta V. The role of PC-1 and ACE genes in diabetic nephropathy in type 1 diabetic patients: evidence for a polygenic control of kidney disease progression. *Nephrol Dial Transplant*, 17: 1402-1407, 2002.
42. Menzaghi C., Ercolino T., Di Paola R., Berg AH, Warram JH, Scherer PE, Trischitta V., Doria A. A haplotype at the adiponectin locus is associated with obesity and other features of the insulin resistance syndrome. *Diabetes* 51: 2306-2312, 2002.
43. Ma X, Warram JH., Trischitta V., Doria A. Genetic variants at the resistin locus and risk of type 2 diabetes in Caucasians. *J Clin Endocrinol Metab*, 87: 4407-4410, 2002.
44. Pizzuti A., Argiolas A., Di Paola R., Baratta R., Rauseo A, Bozzali M., Vigneri R., Dallapiccola B, Trischitta V., Frittitta L. An ATG repeat in the 3'UTR of the

- hResistin gene is associated with a decreased risk for insulin resistance. *J Clin Endocrinol Metab*, 87:4403-4406, 2002.
45. Nisticò L, Giorgi G, Giordano M, Galgani A, Petrone A, D'Alfonso S, Federici M, Di Mario U, Pozzilli P, Buzzetti R, Cascino I. IL12B polymorphism and type 1 diabetes in the Italian population: a case-control study. *Diabetes*. 51(5):1649-50, 2002.
 46. Bottini N, Meloni GF, Borgiani P, Giorgini A, Buzzetti R, Pozzilli P, Lucarelli P, Gloria-Bottini F. Genotypes of cytosolic low-molecular-weight protein-tyrosine-phosphatase correlate with age at onset of type 1 diabetes in a sex-specific manner. *Metabolism*. 2002.
 47. Sesti G. Searching for type 2 diabetes genes: prospects in pharmacotherapy. *The Pharmacogenomics J*. 2:25-29, 2002.
 48. Marchetti P, Lupi R, Federici M, Marselli L, Masini M, Boggi U, Del Guerra S, Patane G, Piro S, Anello M, Bergamini E, Purrello F, Lauro R, Mosca F, Sesti G, Del Prato S. Insulin secretory function is impaired in isolated human islets carrying the Gly(972)->Arg IRS-1 polymorphism. *Diabetes* 51:1419-1424, 2002.
 49. Barbieri M, Bonafè M, Marfella R, Ragno E, Giugliano D, Franceschi C, Paolisso G. LL-paraoxonase genotype is associated with a more severe degree of homeostasis model assessment IR in healthy subjects. *J Clin Endocrinol Metab*. 87(1):222-5, 2002.
 50. Bernassola F, Federici M, Corazzari M, Terrinoni A, Hribal ML, De Laurenzi V, Ranalli M, Massa O, Sesti G, McLean WH, Citro G, Barbetti F, Melino G. Role of transglutaminase 2 in glucose tolerance: knockout mice studies and a putative mutation in a MODY patient. *FASEB J* 16:1371-1378, 2002.
 51. D'Alfonso R, Marini MA, Frittitta L, Sorge R, Frontoni S, Porzio O, Mariani LM, Lauro D, Gambardella S, Trischitta V, Federici M, Lauro R, Sesti G. Polymorphisms of the insulin receptor substrate-2 in patients with type 2 diabetes. *J Clin Endocrinol Metab*. 88:317-322, 2003.
 52. De Cosmo S., Trevisan R., Dalla Vestra M., Vedovato M., Argiolas A, Solini A., Saller A., Damone F., Tiengo A, Trischitta V., Fioretto P. The PC-1 aminoacid variant Q121 associates with lower GFR in type 2 diabetic patients with abnormal albumin excretion rate. *Diabetes Care*, 10:2898-902, 2003.
 53. Battelino T, Ursic-Bratina N, Dolzan V, Stopar-Obreza M, Pozzilli P, Krzysnik C, Vidan-Jeras B. The HLA-DRB, -DQB polymorphism and anti-insulin antibody

- response in Slovenian patients with type 1 diabetes. *Eur J Immunogenet.* 30(3):223-7, 2003.
54. Bugawan TL, Mirel DB, Valdes AM, Panelo A, Pozzilli P, Erlich HA. Association and interaction of the IL4R, IL4, and IL13 loci with type 1 diabetes among Filipinos. *Am J Hum Genet.* 72(6):1505-14, 2003.
55. Federici M, Petrone A, Porzio O, Bizzarri C, Lauro D, D'Alfonso R, Cappa M, Patera I, Nisticò L, Baroni M, Sesti G, Di Mario U, Lauro R, Buzzetti R The Gly⁹⁷²→Arg IRS-1 variant is associated with type 1 diabetes in Continental Italy. *Diabetes* 52: 887-890, 2003.
56. Sesti G, Cardellini M, Marini MA, Frontoni S, D'Adamo M, Del Guerra S, Lauro D, De Nicolais P, Sbraccia P, Del Prato S, Gambardella S, Federici M, Marchetti P, Lauro R. A common polymorphism in the promoter of UCP2 contributes to the variation in insulin secretion in glucose-tolerant subjects. *Diabetes* 52: 1280-1283, 2003.
57. Marini MA, Frontoni S, Mineo D, Bracaglia D, Cardellini M, De Nicolais P, Baroni A, D'Alfonso R, Perna M, Lauro D, Federici M, Gambardella S, Lauro R, Sesti G. The Arg⁹⁷²-variant in insulin receptor substrate-1 is associated with an atherogenic profile in offspring of type 2 diabetic patients. *J Clin Endocrinol Metab* 88: 3368-3371, 2003.
58. Ceravolo R, Maio R, Cuda G, Scozzafava A, Sciacqua A, Vatrano M, Bellieni G, D'Angelo G, Schipani F, Sesti G, Perticone F. Relation of fasting insulin related to insertion/deletion polymorphism of angiotensin converting enzyme-gene and cardiac mass in never-treated patients with systemic hypertension. *Am J Cardiol* 92:1234–1237, 2003.
59. Lindi V, Schwab U, Louheranta A, Laakso M, Vessby B, Hermansen K, Storlien L, Riccardi G, A Rivellese A, KANWU Study Group. Impact of the Pro12Ala polymorphism of the PPAR-gamma2 gene on serum triacylglycerol response to n-3 fatty acid supplementation. *MOLECULAR GENETICS AND METABOLISM*, vol. 79, p. 52-60, 2003.
60. Baratta R., Di Paola R., Spampinato D., Fini G., Marucci A.,. Coco A, Vigneri R., Frittitta L., Trischitta V. Evidence for genetic epistasis in human insulin resistance: the combined effect of PC-1 (K121Q) and PPAR γ 2 (P12A) polymorphisms. *J Mol Med*, 81:718-723, 2003.

61. Pucci L, Lucchesi D, Chirulli V, Penno G, Johansson I, Gervasi P, Del Prato S, Longo V. Cytochrome P450 2J2 polymorphism in healthy Caucasians and those with diabetes mellitus. *Am J Pharmacogenomics*. 3: 355-358, 2003.
62. Federici M, Pandolfi A, De Filippis EA, Pellegrini G, Menghini R, Lauro D, Cardellini M, Romano M, Sesti G, Lauro R, Consoli A G972R IRS-1 variant impairs insulin regulation of eNOS in cultured human endothelial cells. *Circulation* 109: 399-405, 2004.
63. Sesti G, Marini MA, Cardellini M, Sciacqua A, Frontoni S, Andreozzi F, Irace C, Lauro D, Gnasso A, Federici M, Perticone F, Lauro R. The Arg972 variant in insulin receptor substrate-1 is associated with an increased risk of secondary failure to sulfonylurea in patients with type 2 diabetes. *Diabetes Care* 27:1394–1398, 2004.
64. Miscio G, Tassi V., Coco A., Soccio T., Di Paola R., Prudente S., Baratta R., Frittitta L., Ludovico O, Padovano L., Dallapiccola B., Di Mario U., De Cosmo S., Trischitta V. The allelic variant of LAR gene promoter –127 bp Ta□A is associated with reduced risk of obesity and other features related to insulin resistance. *J Mol Med*, 82: 459-466, 2004.
65. Bacci S., Menzaghi C., Ercolino T., Ma X., Rauseo A., Salvemini L., Vigna C., Fanelli R., Di Mario U., Doria A, Trischitta V. The +276 G/T single nucleotide polymorphism of the Adiponectin gene is associated with coronary artery disease in type 2 diabetic patients. *Diabetes Care*, 8: 2015-2020, 2004.
66. Ma X., Bacci S., Mlynarski W., Gottardo L., Soccio T., Menzaghi C., Iori E., Lager RA, Shroff AR., Gervino EV., Johnstone MT., Abumrad N., Avogaro A., Trischitta V., Doria A. A Common Haplotype at the CD36 Locus Is Associated With High FFA Levels and Increased Cardiovascular Risk in Caucasians *Hum Mol Genet*. 19:2197-2205, 2004.
67. Menzaghi C., Ercolino T., Salvemini L., Coco A., Kim SH, Fini G., Doria A., Trischitta V. Multigenic control of serum adiponectin levels: evidence for a role of the APM gene and a locus on 14q13. *Physiological Genomics* 2: 170-174, 2004.
68. D'Adamo M, Perego L, Cardellini M, Marini MA, Frontoni S, Andreozzi F, Sciacqua A, Lauro D, Sbraccia P, Federici M, Paganelli M, Pontiroli AE, Lauro R, Perticone F, Folli F, Sesti G. The -866A/A genotype in the promoter of the human uncoupling protein 2 (UCP2) gene is associated with insulin resistance and increased risk of type 2 diabetes mellitus. *Diabetes* 53: 1905–1910, 2004.

69. Perticone F, Sciacqua A, Scozzafava A, Ventura G, Laratta E, Pujia A, Federici M, Lauro R, Sesti G. Impaired endothelial function in never-treated hypertensive subjects carrying the Arg972 polymorphism in the Insulin Receptor Substrate-1 Gene. *J Clin Endocrinol Metab* 89: 3606-3609, 2004.
70. Guo D, Li M, Zhang Y, Yang P, Eckenrode S, Hopkins D, Zheng W, Purohit S, Podolsky RH, Muir A, Wang J, Dong Z, Brusko T, Atkinson M, Pozzilli P, Zeidler A, Raffel LJ, Jacob CO, Park Y, Serrano-Rios M, Larrad MT, Zhang Z, Garchon HJ, Bach JF, Rotter JI, She JX, Wang CY. A functional variant of SUMO4, a new I kappa B alpha modifier, is associated with type 1 diabetes. *Nat Genet.* 36(8):837-41, 2004.
71. Buzzetti R, Galgani A, Petrone A, Del Buono ML, Erlich HA, Bugawan TL, Lorini R, Meschi F, Multari G, Pozzilli P, Locatelli M, Bottazzo G, Di Mario U. Genetic prediction of type 1 diabetes in a population with low frequency of HLA risk genotypes and low incidence of the disease (the DIABFIN study). *Diabetes Metab Res Rev.*, 2004.
72. De Cosmo S, Marucci A, Ciociola E, Di Paola R, Pucci L, Penno G, Del Prato S, Piras GP, Trevisan R, Giunti S, Viberti GC, Trischitta V. Lack of evidence for the 1484insG variant at the 3'-UTR of the protein tyrosine phosphatase 1B (PTP1B) gene as a genetic determinant of diabetic nephropathy development in type 1 diabetic patients. *Nephrol Dial Transplant.* 19: 2419-24120, 2004.
73. Prudente S, Hribal ML, Flex E, Turchi F, Morini E, De Cosmo S, Bacci S, Tassi V, Cardellini M, Lauro R, Sesti G, Dallapiccola B, Trischitta V. The functional Q84R polymorphism of mammalian tribbles homolog TRB3 is associated with insulin resistance and related cardiovascular risk in Caucasians from Italy. *Diabetes* 54:2807-2811, 2005.
74. Bulotta A., Ludovico O., Coco A., Di Paola R., Quattrone A., Carella A., Pellegrini F., Prudente S., Trischitta V. The common -866G/A polymorphism in the promoter region of the UCP-2 gene is associated with reduced risk of type 2 diabetes in Caucasians from Italy. *J Clin Endocrinol Metab.* 90: 1176-80, 2005.
75. Menzaghi C., Salvemini L., Coco A., Fini G., Di Paola R, Doria A, Trischitta V. Lack of evidence for interaction between APM1 and PPAR γ 2 genes in modulating insulin sensitivity in non diabetic Caucasians from Italy. *J. Intern Med* 257: 315-7, 2005.

76. Cardellini M, Perego L, D'Adamo M, Marini MA, Procopio C, Hribal ML, Andreozzi F, Frontoni S, Giacomelli M, Paganelli M, Pontiroli AE, Lauro R, Folli F, Sesti G. C-174G polymorphism in the promoter of the interleukin-6 gene is associated with insulin resistance. *Diabetes Care* 28: 2007–2012, 2005.
77. Lorini R, Minicucci L, Napoli F, Padovani P, Bazzigaluppi E, Tortoioli C, Cherubini V, Bottazzo G, Pozzilli P, Falorni A, Buzzetti R. Screening for type 1 diabetes genetic risk in newborns of continental Italy. Primary prevention (Prevefin Italy)--preliminary data. *Acta Biomed.*;76 Suppl 3:31-5, 2005.
78. Petrone A, Galgani A, Spoletini M, Alemanno I, Di Cola S, Bassotti G, Picardi A, Manfrini S, Osborn J, Pozzilli P, Buzzetti R. Residual insulin secretion at diagnosis of type 1 diabetes is independently associated with both, age of onset and HLA genotype. *Diabetes Metab Res Rev.* 2005.
79. Fanelli M, Filippi E, Sentinelli F, Romeo S, Fallarino M, Buzzetti R, Leonetti F, Baroni MG. The Gly482Ser missense mutation of the peroxisome proliferator-activated receptor gamma coactivator-1 alpha (PGC-1 alpha) gene associates with reduced insulin sensitivity in normal and glucose-intolerant obese subjects. *Dis Markers* 21(4):175-80, 2005.
80. Sesti G, Perego L, Cardellini M, Andreozzi F, Ricasoli C, Vedani P, Guzzi V, Marchi M, Paganelli M, Ferla G, Pontiroli AE, Hribal ML, Folli F. Impact of common polymorphisms in candidate genes for insulin resistance and obesity on weight loss of morbidly obese subjects after laparoscopic adjustable gastric banding and hypocaloric diet. *J Clin Endocrinol Metab* 90: 5064-5069, 2005.
81. Di Paola R, Tassi V., Trischitta V. About the role of Protein Tyrosine Phosphatase 1 B gene on the Metabolic Syndrome and related traits. *J. Intern Med* 258:289-90, 2005.
82. Bacci S., Ludovico O., Prudente S., Zhang Y.Y, Di Paola R., Mangiacotti D., Rauseo A., Nolan D., Duffy J., Fini G., Salvemini L., Amico C., Vigna C., Pellegrini F., Menzaghi C., Doria A., Trischitta V. The K121Q polymorphism of the ENPP1/PC-1 gene is associated with insulin resistance/atherogenic phenotypes including earlier onset of type 2 diabetes and of myocardial infarction. *Diabetes* 54:3021-5, 2005.
83. Rizza S, Tesauro M, Cardellini M, Meneghini R, Bellia A, Marini MA, Lauro D, Sbraccia P, Sesti G, Lauro R, Federici M. Insulin resistance and increased intimal medial thickness in glucose tolerant offspring of type 2 diabetic subjects carrying

the D298D genotype of endothelial nitric oxide synthase. *Arterioscler Thromb Vasc Biol* 26:431-432, 2006.

84. Menzaghi C., Paroni G., De Bonis C., Soccio T, Marucci A. , Bacci S., . Trischitta V. The –318 C>G Single-Nucleotide Polymorphism in the GNAI2 gene Promoter Region Impairs Transcriptional Activity through Specific Binding of Sp1 Transcription Factor and is Associated with High Blood Pressure in Caucasians from Italy. *J Am Soc Nephrol*, 17: S115-S119, 2006.
85. Di Paola R., Ciociola E., Boonyasrisawat W., Nolan D., Duffy J., Miscio G., Cisternino C., Fini G., Tassi V., Doria A., Trischitta V. Association of hGrb10 genetic variations with Type 2 Diabetes. *Diabetes Care*, 29:1181-1183, 2006.
86. Menzaghi C., Coco A., Salvemini L., Thompson R., De Cosmo S., Doria A., Trischitta V. Heritability of serum resistin and its genetic correlation with insulin resistance-related features in non-diabetic Caucasians. *J Clin Endocrinol Metab* 7: 2792-5, 2006.
87. Soccio T., Zhang Y-Y, Bacci S, Mlynarski W., Placha G., Raggio G., Di Paola R., Marucci A., Johnstone M.T., Gervino E.,V, Abumrad N., Klein S., Trischitta V., Doria A. Common Haplotypes At The Adiponectin Receptor 1 (ADIPOR1) Locus Are Associated With Increased Risk of Coronary Artery Disease in Type 2 Diabetes. *Diabetes*, 55:2763-70, 2006.
88. Prudente S., Trischitta V. The pleiotropic effect of the ENPP1 (PC-1) gene on insulin resistance, obesity and type 2 diabetes – Editorial *J Clin Endocrinol Metab*, 91: 4767–4768, 2006.
89. Pedrinelli R, Dell'Omo G, Penno G, Di Bello V, Pucci L, Fotino C, Lucchesi D, Del Prato S, Dal Fiume C, Barlassina C, Cusi D. Alpha-adducin and angiotensin-converting enzyme polymorphisms in hypertension: evidence for a joint influence on albuminuria. *J Hypertens*. 2006; 24 (5): 931-7. *Erratum in: J Hypertens*. 24: 931-937, 2006.
90. Scarpelli D, Cardellini M, Andreozzi F, Laratta E, Hribal ML, Marini MA, Tassi V, Lauro R, Perticone F, Sesti G. Variants of the interleukin-10 promoter gene are associated with obesity and insulin resistance but not type 2 diabetes mellitus in Caucasian Italian subjects. *Diabetes* 55: 1529-1533, 2006.
91. Sesti G, Laratta E, Cardellini M, Andreozzi F, Del Guerra S, Irace C, Gnasso A, Grupillo M, Lauro R, Hribal ML, Perticone F, Marchetti P. The E23K variant of KCNJ11 encoding the pancreatic beta-cell KATP channel subunits Kir6.2 is

- associated with an increased risk of secondary failure to sulfonylurea in patients with type 2 diabetes. *J Clin Endocrinol Metab.* 91: 2334-2339, 2006.
92. Rizza S, Tesauro M, Cardellini M, Menghini R, Bellia A, Marini MA, Lauro D, Sbraccia P, Sesti G, Lauro R, Federici M. Insulin resistance and increased intimal medial thickness in glucose tolerant offspring of type 2 diabetic subjects carrying the D298D genotype of endothelial nitric oxide synthase. *Arterioscler Thromb Vasc Biol.* 26(2):431-2, 2006.
 93. Buzzetti R, Petrone A, Zavarella S, Zampetti S, Spoletini M, Potenziani S, Leto G, Osborn J, Leonetti F. The glucose clamp reveals an association between adiponectin gene polymorphisms and insulin sensitivity in obese subjects. *Int J Obes (Lond)*;31(3):424-8, 2007.
 94. Prudente S., Chandalia M., Morini E., Baratta R., Dallapiccola B., Abate N., Frittitta L., Trischitta V. The Q121/Q121 genotype of ENPP1/PC-1 is associated with lower BMI in non-diabetic Caucasians. *Obesity*, 15: 1-4, 2007.
 95. Boonyasrisawat W., Eberle D., Bacci S., Zhang Y.-Y., Nolan D., Gervino E. V., Johnstone M.T., Trischitta V., Shoelson S. E., Doria A. Tag polymorphisms at the A20 (TNFAIP3) locus are associated with lower gene expression and increased risk of coronary artery disease in type 2 diabetes. *Diabetes*, 54: 499-505, 2007.
 96. Menzaghi C., Trischitta V., Doria A. Perspectives in Diabetes. Genetic Influences of Adiponectin on Insulin Resistance, Type 2 Diabetes, and Cardiovascular Disease. *Diabetes*, 56: 1198-1209, 2007.
 97. Fiorito M, Torrente I, De Cosmo S, Guida V, Colosimo A, Prudente S, Flex E, Menghini R, Miccoli R, Penno G, Pellegrini F, Tassi V, Federici M, Trischitta V, Dallapiccola B. Interaction of DIO2 T92A and PPARgamma2 P12A polymorphisms in the modulation of metabolic syndrome. *Obesity (Silver Spring)* 15: 2889-2895, 2007.
 98. Pascoe L, Tura A, Patel SK, Ibrahim IM, Ferrannini E, Zeggini E, Weedon MN, Mari A, Hattersley AT, McCarthy MI, Frayling TM, Walker M; RISC Consortium; U.K. Type 2 Diabetes Genetics Consortium. Common variants of the novel type 2 diabetes genes CDKAL1 and HHEX/IDE are associated with decreased pancreatic beta-cell function. *Diabetes* ;56(12):3101, 2007.
 99. Prudente S., Flex E., Morini E., Turchi F., Capponi D., De Cosmo S., Tassi V., Guida V., Avogaro A, Folli F., Maiani F., Frittitta L., Dallapiccola B., Trischitta V. A functional variant of the adipocyte glycerol channel Aquaporin 7 gene is

- associated with obesity and related metabolic abnormalities. *Diabetes* 56: 1468-1474, 2007.
100. Bacci S., De Cosmo S., Prudente S., Trischitta V. ENPP1 gene, insulin resistance and related clinical outcomes. *Curr Opin Clin Nutr Metab Care* 10:403–409, 2007.
 101. Prudente S., Ludovico O., Tassi V., Trischitta V. Heterogeneous effects of gene polymorphism on type 2 diabetes risk: lesson from the PPAR γ 2 Pro12Ala – Editorial. *Nutr Metab Cardiovasc Dis*, 17: 629.631, 2007.
 102. Terruzzi I, Senesi P, Fermo I, Lattuada G, Luzi L. Are genetic variants of the methyl group metabolism enzymes risk factors predisposing to obesity? *J Endocrinol Invest*. 30(9):747-53, 2007.
 103. Vaccaro O., Lapice E., Monticelli A., Giacchetti M., Castaldo I., Galasso R., Pinelli M., Donnarumma G., A. Rivellesse, Cocozza S., Riccardi G.. Pro12Ala polymorphism of the PPAR γ 2 locus modulates the relationship between energy intake and body weight in type 2 diabetic patients. *DIABETES CARE*, vol. 30, p. 1156-1161, 2007.
 104. Perticone F, Maio R, Di Paola R, Sciacqua A, Marucci A, De Cosmo S, Perticone M, Sesti G, Trischitta V. Role of PC-1 and ACE genes on insulin resistance and cardiac mass in never-treated hypertensive patients. Suggestive evidence for a digenic additive modulation. *Nutr Metab Cardiovasc Dis* 17:181-187, 2007.
 105. Ludovico O, Pellegrini F, Di Paola R, Minenna A, Mastroianno S, Cardellini M, Marini MA, Andreozzi F, Vaccaro O, Sesti G, Trischitta V. Heterogeneous Effect of Peroxisome Proliferator-activated Receptor γ 2 Ala12 Variant on Type 2 Diabetes Risk. *Obesity* 15:1076-1081, 2007.
 106. De Cosmo S, Prudente S, Andreozzi F, Morini E, Rauseo A, Scarpelli D, Zhang YY, Xu R, Perticone F, Dallapiccola B, Sesti G, Doria A, Trischitta V. Glutamine to arginine substitution at amino acid 84 of mammalian tribbles homolog TRIB3 and CKD in whites with type 2 diabetes. *Am J Kidney Dis* 50:688-689, 2007.
 107. Perticone F, Sciacqua A, Barlassina C, Signorello MC, Andreozzi F, Sesti G, Cusi D. Gly460Trp α -adducin gene polymorphism and endothelial function in untreated hypertensive patients. *J Hypertens* 25:2234-2239, 2007.

108. Spoletini M, Petrone A, Zampetti S, Capizzi M, Zavarella S, Osborn J, Foffi C, Tuccinardi D, Pozzilli P, Buzzetti R; IMDIAB Group. Low-risk HLA genotype in Type 1 diabetes is associated with less destruction of pancreatic B-cells 12 months after diagnosis. *Diabet Med.* 2007.
109. Dell'Omo G, Penno G, Pucci L, Fotino C, Lucchesi D, Del Prato S, Pedrinelli R. Lack of association between endothelial nitric oxide synthase gene polymorphisms, microalbuminuria and endothelial dysfunction in hypertensive men. *J Hypertens.*; 25: 1389-1395, 2007.
110. Iyengar SK, Abboud HE, Goddard KA, Saad MF, Adler SG, Arar NH, Bowden DW, Duggirala R, Elston RC, Hanson RL, Ipp E, Kao WH, Kimmel PL, Klag MJ, Knowler WC, Meoni LA, Nelson RG, Nicholas SB, Pahl MV, Parekh RS, Quade SR, Rich SS, Rotter JI, Scavini M, Schelling JR, Sedor JR, Sehgal AR, Shah VO, Smith MW, Taylor KD, Winkler CA, Zager PG, Freedman BI for the Family Investigation of Nephropathy and Diabetes Research Group. Genome-wide scans for diabetic nephropathy and albuminuria in multiethnic populations: the family investigation of nephropathy and diabetes (FIND). *Diabetes* 56:1577-1585, 2007.
111. Romeo S, Sentinelli F, Cavallo MG, Leonetti F, Fallarino M, Mariotti S, Baroni MG. Search for genetic variants of the SYNTAXIN 1A (STX1A) gene: the -352 A>T variant in the STX1A promoter associates with impaired glucose metabolism in an Italian obese population. *Int J Obes (Lond).* 32(3):413-20, 2008.
112. Tassi V., Coco A., Padovano L., De Bonis C., De Cosmo S., Trischitta V. Akt2 gene common allelic variants in insulin resistance and metabolic syndrome. *Nutr Metab Cardiovasc Dis*, 18:263-70, 2008.
113. Pascoe L, Frayling TM, Weedon MN, Mari A, Tura A, Ferrannini E, Walker M; RISC Consortium. Beta cell glucose sensitivity is decreased by 39% in non-diabetic individuals carrying multiple diabetes-risk alleles compared with those with no risk alleles. *Diabetologia.* 51(11):1989-92, 2008.
114. Lindi V., Schwab U., Louheranta A, Vessby B., Hermansen K., Tapsell L., Riccardi G., Rivellese A, Laakso M., Uusitupa MI: Kanwu Study Group. The G-250A polymorphism in the hepatic lipase gene promoter is associated with changes in hepatic lipase activity and LDL cholesterol: The Kanwu study. *NMCD. NUTRITION METABOLISM AND CARDIOVASCULAR DISEASES*, p. 88-95, 2008.

115. Staiger H, Stančáková A, Zilinskaite J, Vääntinen M, Hansen T, Marini MA, Hammarstedt A, Jansson P-A, Sesti G, Smith U, Pedersen O, Laakso M, Stefan N, Fritsche A, Häring H-U. A candidate type 2 diabetes polymorphism near the HHEX locus affects acute glucose-stimulated insulin release in European populations: results from the EUGENE2 study. *Diabetes* 57:514-517, 2008.
116. McAteer JB., Prudente S., Bacci S., Lyon HN., Hirschhorn JN., Trischitta V., Florez JC. The ENPP1 K121Q polymorphism is associated with type 2 diabetes in European populations: evidence from an updated meta-analysis in 42,042 subjects. *Diabetes* 57: 1125–1130, 2008.
117. Laakso M, Zilinskaite J, Hansen T, Welløv Boesgaard T, Vääntinen M, Stančáková A, Jansson P-A, Pellmé F, Holst JJ, Kuulasmaa T, Hribal ML, Sesti G, Stefan N, Fritsche A, Häring H, Pedersen O, Smith U for the EUGENE2 Consortium. Insulin sensitivity, insulin release and GLP-1 levels in subjects with IFG and/or IGT in the EUGENE2 study. *Diabetologia* 51:502-511, 2008.
118. Boesgaard TW, Zilinskaite J, Vääntinen M, Laakso M, Jansson PA, Hammarstedt A, Smith U, Stefan N, Fritsche A, Häring H, Hribal ML, Sesti G, Zobel DP, Pedersen O, Hansen T; for the EUGENE 2 Consortium. The common SLC30A8 Arg325Trp variant is associated with reduced first phase insulin release in 846 non-diabetic offspring of type 2 diabetes patients-the EUGENE2 study. *Diabetologia* 51:816-820, 2008.
119. Di Castelnuovo A, Soccio M, Iacoviello L, Evangelista V, Consoli A, Vanuzzo D, Diviacco S, Carluccio M, Rignanese L, De Caterina R. The C242T polymorphism of the p22phox component of NAD(P)H oxidase and vascular risk. Two case-control studies and a meta-analysis. *Thromb Haemost.*99(3):594-601, 2008.
120. Stančáková A, Pihlajamäki J, Kuusisto J, Stefan N, Fritsche A, Häring H, Andreozzi F, Succurro E, Sesti G, Boesgaard TW, Hansen T, Pedersen O, Jansson PA, Hammarstedt A, Smith U, Laakso M; EUGENE2 Consortium. SNP rs7754840 of CDKAL1 is associated with impaired insulin secretion in non-diabetic offspring of Type 2 diabetic subjects (the EUGENE2 study) and in a large sample of men with normal glucose tolerance. *J Clin Endocrinol Metab* 93:1924–1930, 2008.
121. Gottardo L., De Cosmo S., Zhang Y-Y., Powers C., Prudente S., Marescotti M. C., Trischitta V., Avogaro A., Doria A. A Polymorphism at the IL6ST (gp130)

Locus is Associated with Traits of the Metabolic Syndrome. *Obesity* 1:205-10, 2008.

122. Rittig K, Peter A, Baltz KM, Tschritter O, Weigert C, Andreozzi F, Perticone F, Siegel-Axel DI, Stefan N, Fritsche A, Salih HR, Schleicher E, Machicao F, Sesti G, Häring HU, Balletshofer BM. The CCR2 promoter polymorphism T-960A, but not the serum MCP-1 level, is associated with endothelial function in prediabetic individuals. *Atherosclerosis* 198:338-346, 2008.
123. Andreozzi F, Formoso G, Prudente S, Hribal ML, Pandolfi A, Bellacchio E, Di Silvestre S, Trischitta V, Consoli A, Sesti G. TRIB3 R84 variant is associated with impaired insulin mediated nitric oxide production in human endothelial cells. *Arterioscler Thromb Vasc Biol* 28:1355-1360, 2008.
124. Morini E., Tassi V., Capponi D., Ludovico O., Dallapiccola B., Trischitta V., Prudente S. Interaction between PPAR γ 2 variants and gender on the modulation of body weight. *Obesity* 16: 1467-1470, 2008.
125. Menzaghi C., Paroni G., De Bonis C., Coco A., Vigna C., Miscio G., Lanna P., Tassi V., Bacci S., Trischitta V. The protein tyrosine phosphatase receptor type f (PTPRF) locus is associated with coronary artery disease in type 2 diabetes. *J. Intern Med* 263: 653-654, 2008.
126. Hribal ML, Tornei F, Pujol A, Menghini R, Barcaroli D, Lauro D, Amoroso R, Lauro R, Bosch F, Sesti G, Federici M. Transgenic mice overexpressing human G972R IRS-1 show impaired insulin action and insulin secretion. *J Cell Mol Med.* 12:2096-106, 2008.
127. Friedrich B, Weyrich P, Stancáková A, Wang J, Kuusisto J, Laakso M, Sesti G, Succurro E, Smith U, Hansen T, Pedersen O, Machicao F, Schäfer S, Lang F, Rislér T, Ullrich S, Stefan N, Fritsche A, Häring HU. Variance of the SGK1 gene is associated with insulin secretion in different European populations: results from the TUEF, EUGENE2, and METSIM studies. *PLoS ONE* 3(11):e3506, 2008.
128. Baratta R., Rossetti P., Prudente S., Barbetti F., Sudano D., Nigro A., Farina M. G, Pellegrini F., Trischitta V., Frittitta L. Role of the ENPP1 K121Q Polymorphism on Glucose Homeostasis. *Diabetes*, 57: 3360–3364, 2008.
129. Noble JA, Martin A, Valdes AM, Lane JA, Galgani A, Petrone A, Lorini R, Pozzilli P, Buzzetti R, Erlich HA. Type 1 diabetes risk for human leukocyte antigen (HLA)-DR3 haplotypes depends on genotypic context: association of DPB1 and

HLA class I loci among DR3- and DR4-matched Italian patients and controls. *Hum Immunol.*; 69(4-5):291-300, 2008.

130. Petrone A, Spoletini M, Zampetti S, Capizzi M, Zavarella S, Osborn J, Pozzilli P, Buzzetti R. Immunotherapy Diabetes (IMDIAB) Group. The PTPN22 1858T gene variant in type 1 diabetes is associated with reduced residual beta-cell function and worse metabolic control. *Diabetes Care.* 31(6):1214-8, 2008.
131. Irace C, Cortese C, Migale M, Liberatoscioli L, Mannucci L, Federici G, Gnasso A. Stromelysin gene promoter polymorphism and common carotid geometry in diabetic subjects. *Int Angiol* 27:413-418, 2008.
132. Barbieri M, Marfella R, Rizzo MR, Boccardi V, Siniscalchi M, Schiattarella C, Siciliano S, Lemme P, Paolisso G. The -8 UTR C/G polymorphism of PSMA6 gene is associated with susceptibility to myocardial infarction in type 2 diabetic patients. *Atherosclerosis.* 201(1):117-23, 2008.
133. Irace C, Cortese C, Fiaschi E, Scavelli F, Liberatoscioli L, Federici G, Gnasso A. The influence of PON1 192 polymorphism on endothelial function in diabetic subjects with or without hypertension. *Hypertens Res.*;31:507-513, 2008.
134. Schelling JR, Abboud HE, Nicholas SB, Pahl MV, Sedor JR, Adler SG, Arar NH, Bowden DW, Elston RC, Freedman BI, Goddard KA, Guo X, Hanson RL, Ipp E, Iyengar SK, Jun G, Kao WH, Kasinath BS, Kimmel PL, Klag MJ, Knowler WC, Nelson RG, Parekh RS, Quade SR, Rich SS, Saad MF, Scavini M, Smith MW, Taylor K, Winkler CA, Zager PG, Shah VO for the Family Investigation of Nephropathy and Diabetes Research Group. Genome-wide scan for estimated glomerular filtration rate in multi-ethnic diabetic populations: the Family Investigation of Nephropathy and Diabetes (FIND). *Diabetes* 57:235-243, 2008.
135. Arar NH, Freedman BI, Adler SG, Iyengar SK, Chew EY, Davis MD, Satko SG, Bowden DW, Duggirala R, Elston RC, Guo X, Hanson RL, Igo RP Jr, Ipp E, Kimmel PL, Knowler WC, Molineros J, Nelson RG, Pahl MV, Quade SR, Rasooly RS, Rotter JI, Saad MF, Scavini M, Schelling JR, Sedor JR, Shah VO, Zager PG, Abboud HE; Family Investigation of Nephropathy and Diabetes Research Group. Heritability of the severity of diabetic retinopathy: the FIND-Eye study. *Invest Ophthalmol Vis Sci* 49:3839-3845, 2008.
136. Malhotra A, Igo RP Jr, Thameem F, Kao WH, Abboud HE, Adler SG, Arar NH, Bowden DW, Duggirala R, Freedman BI, Goddard KA, Ipp E, Iyengar SK, Kimmel PL, Knowler WC, Kohn O, Leehey D, Meoni LA, Nelson RG, Nicholas SB, Parekh

- RS, Rich SS, Chen YD, Saad MF, Scavini M, Schelling JR, Sedor JR, Shah VO, Taylor KD, Thornley-Brown D, Zager PG, Horvath A, Hanson RL; Family Investigation of Nephropathy and Diabetes Research Group. Genome-wide linkage scans for type 2 diabetes mellitus in four ethnically diverse populations-significant evidence for linkage on chromosome 4q in African Americans: the Family Investigation of Nephropathy and Diabetes Research Group. *Diabetes Metab Res Rev* 25:740-7, 2009.
137. Cappelli A, Tumini S, Consoli A, Carinci S, Piersanti C, Ruggiero G, Simonella G, Soletti F, Staffolani P, Pianese L. Novel mutations in GCK and HNF1A genes in Italian families with MODY phenotype. *Diabetes Res Clin Pract.* 83(3):e72-4, 2009.
 138. Dell'Omo G, Penno G, Pucci L, Lucchesi D, Del Prato S, Pedrinelli R. Lack of association between TGF-beta-1 genotypes and microalbuminuria in essential hypertensive men. *Nephrol Dial Transplant.*; 24: 1864-1869, 2009.
 139. De Cosmo S., Minenna A., Zhang Y-Y, Thompson R., Miscio G., Vedovato M., Rauseo A., Saller A, Mastroianno S., Pellegrini F., Trevisan R., Fioretto P., Doria A., Trischitta V. The Q121 Variant of ENPP1 Gene is Associated with Decreased Kidney Function Among Patients With Type 2 Diabetes. *Am J Kidney Dis* 53: 273-80, 2009.
 140. Prudente S., Morini E., Trischitta V. The emerging role of TRIB3 as a gene affecting human insulin resistance and related clinical outcomes. *Acta Diabetol.* 46: 79-84, 2009.
 141. Prudente S, Scarpelli D, Chandalia M, Zhang YY, Morini E, Del Guerra S, Perticone F, Li R, Powers C, Andreozzi F, Marchetti P, Dallapiccola B, Abate N, Doria A, Sesti G, Trischitta V. The TRIB3 Q84R polymorphism and risk of early-onset type 2 diabetes. *J Clin Endocrinol Metab* 94: 190-196, 2009.
 142. Presta I, Andreozzi F, Succurro E, Marini MA, Laratta E, Lauro R, Hribal ML, Perticone F, Sesti G. IL-18 gene polymorphism and metabolic syndrome. *Nutr Metab Cardiovasc Dis* 19:e5-e6, 2009.
 143. Prudente S, Morini E, Trischitta V. Insulin signaling regulating genes: effect on T2DM and cardiovascular risk. *Nat Rev Endocrinol*, 12: 682-693, 2009.
 144. Morini E, Prudente S, Succurro E, Chandalia M, Zhang YY, Mammarella S, Pellegrini F, Powers C, Proto V, Dallapiccola B, Cama A, Sesti G, Abate N, Doria A, Trischitta V. IRS1 G972R polymorphism and type 2 diabetes: a paradigm for the

- difficult ascertainment of the contribution to disease susceptibility of 'low-frequency-low-risk' variants. *Diabetologia* 52:1852–1857, 2009.
145. De Cosmo S., Motterlini N., Prudente S., Pellegrini F., Trevisan R., Bossi A, Remuzzi G., Trischitta V., Ruggenenti P. Impact of the PPAR γ 2 Pro12Ala polymorphism and ACE inhibitor therapy on new-onset microalbuminuria in type 2 diabetes: evidence from Benedict. *Diabetes*, 58: 2920-2929, 2009.
 146. Boesgaard TW, Gjesing AP, Grarup N, Rutanen J, Jansson PA, Hribal ML, Sesti G, Fritsche A, Stefan N, Staiger H, Häring H, Smith U, Laakso M, Pedersen O, Hansen T; EUGENE2 Consortium. Variant near ADAMTS9 known to associate with type 2 diabetes is related to insulin resistance in offspring of type 2 diabetes patients--EUGENE2 study. *PLoS One*. 4:e7236, 2009.
 147. Langenberg C, Pascoe L, Mari A, Tura A, Laakso M, Frayling TM, Barroso I, Loos RJ, Wareham NJ, Walker M; RISC Consortium. Common genetic variation in the melatonin receptor 1B gene (MTNR1B) is associated with decreased early-phase insulin response. *Diabetologia*. 52(8):1537-42, 2009.
 148. Lindroos MM, Majamaa K, Tura A, Mari A, Kalliokoski KK, Taittonen MT, Iozzo P, Nuutila P. m.3243A>G mutation in mitochondrial DNA leads to decreased insulin sensitivity in skeletal muscle and to progressive beta-cell dysfunction. *Diabetes*. 58(3):543-9, 2009.
 149. Barrett JC, Clayton DG, Concannon P, Akolkar B, Cooper JD, Erlich HA, Julier C, Morahan G, Nerup J, Nierras C, Plagnol V, Pociot F, Schuilenburg H, Smyth DJ, Stevens H, Todd JA, Walker NM, Rich SS; Type 1 Diabetes Genetics Consortium. Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. *Nat Genet* 41(6):703-7, 2009.
 150. Beyan H, Buckley LR, Bustin SA, Yousaf N, Pozzilli P, Leslie RD Altered monocyte cyclo-oxygenase response in non-obese diabetic mice. *Clin Exp Immunol* 155(2):304-10, 2009.
 151. Cefalù A. B., Noto D., Arpi M.L., Yin F., Spina R., Hilden H., Barbagallo C.M., Carroccio A., Tarugi P., Squatrito S., Vigneri R., Taskinen M.R., Péterfy M., and Aversa M.R. Novel *LMFI* Nonsense Mutation in a Patient with Severe Hypertriglyceridemia. *J. Clin. Endocrinol. Metab.*: 94(11): 4584-4590, 2009.
 152. Romeo S, Sentinelli F, Dash S, Yeo GS, Savage DB, Leonetti F, Capoccia D, Incani M, Maglio C, Iacovino M, O'Rahilly S, Baroni MG. Morbid obesity exposes

the association between PNPLA3 I148M (rs738409) and indices of hepatic injury in individuals of European descent. *Int J Obes (Lond)*. 34(1):190-4, 2010.

153. Zampetti S, Spoletini M, Petrone A, Capizzi M, Arpi ML, Tiberti C, Di Pietro S, Bosi E, Pozzilli P, Giorgino F, Buzzetti R; Nirad Study Group. Association of F7L2 gene variants with low GAD autoantibody titre in LADA subjects (NIRAD Study 5). *Diabet Med*, 2010.
154. Hilner JE, Perdue LH, Sides EG, Pierce JJ, Wagner AM, Aldrich A, Loth A, Albret L, Wagenknecht LE, Nierras C, Akolkar B; T1DGC. Designing and implementing sample and data collection for an international genetics study: the Type 1 Diabetes Genetics Consortium (T1DGC). *Clin Trials*.;7(1 Suppl):S5-S32, 2010.
155. Zampetaki A., Kiechl S., Drozdov I., Willeit P., Mayr U., Prokopi M., Mayr A., Weger S., Oberhollenzer F., Bonora E., Shah A., Willeit J., Mayr M.. Plasma MicroRNA Profiling Reveals Loss of Endothelial MiR-126 and Other MicroRNAs in Type 2 Diabetes. *Circ. Res.* 107: 810-817, 2010.
156. Marucci A., Miscio G., Padovano L., Boonyasrisawat W., Doria A., Trischitta V., Di Paola R. A functional variant in the gene 3' untranslated region regulates HSP70 expression and is a potential candidate for insulin resistance-related abnormalities. *J. Intern Med* 267: 237-240, 2010.
157. Menzaghi C., Trischitta V. Genetic of serum resistin: a paradigm of population-specific regulation? *Diabetologia* 53: 226-228, 2010.
158. Incani M., Cambuli V.M., Cavalot F, Congiu T., Paderi M., Sentinelli F., Romeo S., Poy P, Soro M., Pilia S., Loche S., Cossu E., Trovati M., Mariotti S., Baroni M.G.. Clinical Application of best practice guidelines for the genetic diagnosis of MODY2 and MODY3. *Diabetic Medicine* 27 (11):1331-3, 2010.
159. Di Paola R, Wojcik J, Succurro E, Marucci A, Chandalia M, Padovano L, Powers C, Merla M, Abate N, Sesti G, Doria A, Trischitta V. GRB10 Gene and type 2 diabetes in whites. *J Intern Med* 267:132-133, 2010.
160. Prudente S, Baratta R, Andreozzi F, Morini E, Farina MG, Nigro A, Copetti M, Pellegrini F, Succurro E, Di Pietrantonio L, Brufani C, Barbetti F, Dallapiccola B, Sesti G, Trischitta V, Frittitta L. TRIB3 R84 variant affects glucose homeostasis by altering the interplay between insulin sensitivity and secretion. *Diabetologia* 53:1354–1361, 2010.

161. Sathananthan A., Man C.D., Micheletto F., Zinsmeister A.R., Camilleri M., Giesler P.D., Laugen J.M., Toffolo G.M., Rizza R.A., Cobelli C., Vella A.: Common genetic variation in GLP1R and insulin secretion in response to exogenous GLP-1 in nondiabetic subjects: a pilot study. *Diabetes Care* 33:2074-2076, 2010.
162. Menzaghi C., Salvemini L., Paroni G., De Bonis C., Mangiacotti D., Fini G, Doria A., Di Paola R, Trischitta V. Circulating HMW adiponectin isoform is heritable and shares a common genetic background with insulin resistance in non diabetic White Caucasians from Italy: evidences from a family based study. *J. Intern Med* 267: 287-294, 2010.
163. Sun Y, Scavini M, Orlando RA, Murata GH, Servilla KS, Tzamaloukas AH, Schrader R, Bedrick EJ, Burge MR, Abumrad NA, Zager PG. Increased CD36 expression signals monocyte activation among patients with type 2 diabetes. *Diabetes Care* 33:2065-7, 2010.
164. MacCluer JW, Scavini M, Shah VO, Cole SA, Laston SL, Voruganti VS, Paine SS, Eaton AJ, Comuzzie AG, Tentori F, Pathak DR, Bobelu A, Bobelu J, Ghahate D, Waikaniwa M, Zager PG. Heritability of measures of kidney disease among Zuni Indians: the Zuni Kidney Project. *Am J Kidney Dis* 56:289-302, 2010.
165. Kiechl S., Parè G., Barbalic M., Dupuis J., Dehghan A., Bis J.C., Laxton R.C., Xiao Q., Bonora E., Willeit J., Xu Q., Witteman J.C., Chasman D., Tracy R.P., Ballantyne C.M., Ridker P.M., Benjamin E.J., Ye S. Association of variation at the ABO locus with circulating levels of sICAM-1, sP-selectin and sE-selectin: a Meta-Analysis. *Circ. Cardiovasc. Genet.* 4: 681-686, 2011.
166. Chiefari E, Tanyolaç S, Paonessa F, Pullinger CR, Capula C, Iiritano S, Mazza T, Forlin M, Fusco A, Durlach V, Durlach A, Malloy MJ, Kane JP, Heiner SW, Filocamo M, Foti DP, Goldfine ID, Brunetti A. Functional variants of the HMGAI gene and type 2 diabetes mellitus. *JAMA*; 305: 903-912, 2011.
167. Brunetti A, Chiefari E, Foti D. Perspectives on the contribution of genetics to the pathogenesis of type 2 diabetes mellitus. *Recenti Prog Med*; 102: 468-75, 2011.
168. Mosca L, Penco S, Patrosso MC, Marocchi A, Lapolla A, Sartore G, Chillelli NC, Paleari R, Mosca A. Genetic variability of the fructosamine 3-kinase gene in diabetic patients. *Clin Chem Lab Med.*;49(5):803-8, 2011.
169. De Cosmo S, Prudente S, Lamacchia O, Lapice E, Morini E, Di Paola R, Copetti M, Ruggenenti P, Remuzzi G, Vaccaro O, Cignarelli M, Trischitta V. PPAR γ 2

- P12A polymorphism and albuminuria in patients with type 2 diabetes: a meta-analysis of case-control studies. *Nephrol Dial Transplant*. 26(12):4011-6, 2011.
170. Bacci S., Rizza S., Prudente S, Spoto B., Powers C., Facciorusso A., Pacilli A., Lauro D., Testa A., Zhang Y-Y, Di Stolfo G., Mallamaci F., Tripepi G., Xu R., Mangiacotti D., Aucella F., Lauro R., Gervino EV., Hauser TH., Copetti M., De Cosmo S., Pellegrini F., Zoccali C., Federici M., Doria A., Trischitta V. The *ENPP1* Q121 variant predicts major cardiovascular events in high-risk individuals. Evidence for interaction with obesity in diabetic patients. *Diabetes* 60:1000-7, 2011.
 171. Formoso G, Di Tomo P, Andreozzi F, Succurro E, Di Silvestre S, Prudente S, Perticone F, Trischitta V, Sesti G, Pandolfi A, Consoli A. The *TRIB3* R84 variant is associated with increased carotid intima-media thickness in vivo and with enhanced MAPK signalling in human endothelial cells. *Cardiovasc Res* 89:184-92, 2011.
 172. Hribal ML, Presta I, Procopio T, Marini MA, Stančáková A, Kuusisto J, Andreozzi F, Hammarstedt A, Jansson PA, Grarup N, Hansen T, Walker M, Stefan N, Fritsche A, Häring HU, Pedersen O, Smith U, Laakso M, Sesti G; on behalf of the EUGENE2 Consortium. Glucose tolerance, insulin sensitivity and insulin release in European non-diabetic carriers of a polymorphism upstream of *CDKN2A* and *CDKN2B*. *Diabetologia* 54:795–802, 2011.
 173. Sharma R, Prudente S, Andreozzi F, Powers C, Mannino G, Bacci S, Gervino EV, Hauser TH, Succurro E, Mercuri L, Goheen EH, Shah H, Trischitta V, Sesti G, Doria A. The type 2 diabetes and insulin-resistance locus near *IRS1* is a determinant of HDL cholesterol and triglycerides levels among diabetic subjects. *Atherosclerosis* 216:157-160, 2011.
 174. Prudente S, Morini E, Larmon J, Andreozzi F, Di Pietro N, Nigro A, Gervino EV, Mannino GC, Bacci S, Hauser TH, Bellacchio E, Formoso G, Pellegrini F, Proto V, Menzaghi C, Frittitta L, Pandolfi A, Sesti G, Doria A, Trischitta V. The *SH2B1* obesity locus is associated with myocardial infarction in diabetic patients and with NO synthase activity in endothelial cells. *Atherosclerosis* 219:667-672, 2011.
 175. Hunt KA, Smyth DJ, Balschun T, Ban M, Mistry V, Ahmad T, Anand V, Barrett JC, Bhaw-Rosun L, Bockett NA, Brand OJ, Brouwer E, Concannon P, Cooper JD, Dias KR, van Diemen CC, Dubois PC, Edkins S, Fölster-Holst R, Fransen K, Glass DN, Heap GA, Hofmann S, Huizinga TW, Hunt S, Langford C, Lee J, Mansfield J, Marrosu MG, Mathew CG, Mein CA, Müller-Quernheim J, Nutland S, Onengut-

- Gumuscu S, Ouwehand W, Pearce K, Prescott NJ, Posthumus MD, Potter S, Rosati G, Sambrook J, Satsangi J, Schreiber S, Shtir C, Simmonds MJ, Sudman M, Thompson SD, Toes R, Trynka G, Vyse TJ, Walker NM, Weidinger S, Zhernakova A, Zoledziwska M; Type 1 Diabetes Genetics Consortium; UK Inflammatory Bowel Disease (IBD) Genetics Consortium; Wellcome Trust Case Control Consortium, Weersma RK, Gough SC, Sawcer S, Wijmenga C, Parkes M, Cucca F, Franke A, Deloukas P, Rich SS, Todd JA, van Heel DA. Rare and functional SIAE variants are not associated with autoimmune disease risk in up to 66,924 individuals of European ancestry. *Nat Genet.* 27;44(1):3-5, 2011.
176. Boccardi V, Rizzo MR, Marfella R, Papa M, Esposito A, Portoghese M, Paolisso G, Barbieri M. -94 ins/del ATTG NFKB1 gene variant is associated with lower susceptibility to myocardial infarction. *Nutr Metab Cardiovasc Dis.* 21(9):679-84, 2011.
 177. Buzzetti R, Cernea S, Petrone A, Capizzi M, Spoletini M, Zampetti S, Guglielmi C, Venditti C, Pozzilli P; DiaPep Trialists Group. C-peptide response and HLA genotypes in subjects with recent-onset type 1 diabetes after immunotherapy with DiaPep277: an exploratory study. *Diabetes.* 60(11):3067-72, 2011.
 178. Bonetti S., Trombetta M., Malerba G., Boselli L., Trabetti E., Muggeo M., Stoico V., Negri C., Pignatti P.F., Bonora E., Bonadonna R.C. Variants and haplotypes of TCF7L2 are associated with {beta}-cell function in patients with newly diagnosed type 2 diabetes: the Verona Newly Diagnosed Type 2 Diabetes Study (VNDS). 1. *J. Clin. Endocrinol Metab.* 96: E389-393, 2011.
 179. Bonetti S., Trombetta M., Boselli M.L., Turrini F., Malerba G., Trabetti E., Pignatti P.F., Bonora E., Bonadonna R.C. Variants of GCKR affect both beta-cell and kidney function in patients with newly diagnosed type 2 diabetes: The Verona Newly Diagnosed Type 2 Diabetes Study. 2. *Diabetes Care* 34: 1205-1210, 2011.
 180. Ragona F, Lattuada G, Ruotolo G, Luzi L, Perseghin G. Lack of association of apoE ε4 allele with insulin resistance. *Acta Diabetol* 49:25-32, 2012.
 181. Spoto B, Testa A, Parlongo RM, Tripepi G, Trischitta V, Mallamaci F, Zoccali C. Insulin resistance and left ventricular hypertrophy in end-stage renal disease: association between the ENPP1 gene and left ventricular concentric remodelling. *Nephrol Dial Transplant.* 27:661-666, 2012.
 182. Mannino GC, Sesti G. Individualized therapy for type 2 diabetes: clinical implications of pharmacogenetic data. *Mol Diagn Ther* 16:285-302, 2012.

- 183.** Andreozzi F, Presta I, Mannino GC, Scarpelli D, Di Silvestre S, Di Pietro N, Succurro E, Sciacqua A, Pandolfi A, Consoli A, Hribal ML, Perticone F, Sesti G. A functional variant of the dimethylarginine dimethylaminohydrolase-2 gene is associated with insulin sensitivity. *PLoS One* 7: e36224, 2012.
- 184.** Trombetta M., Bonetti S., Boselli M., Turrini F., Malerba G., Trabetti E., Pignatti P., Bonora E., Bonadonna R.C. CACNA1E variants affect beta cell function in patients with newly diagnosed type 2 diabetes. the Verona newly diagnosed type 2 diabetes study (VNDS) 3. *PLoS One* 7: e32755, 2012.
- 185.** Brunetti A, Chiefari E, Pullinger CR, Tanyolac S, Foti D, Durlach V, Goldfine ID. Comment on: Marquez et al. Low-frequency variants in HMGA1 are not associated with type 2 diabetes risk. *Diabetes*; 61:524-530, 2012.
- 186.** Qi Q, Menzaghi C, Smith S, Liang L, de Rekeneire N, Garcia ME, Lohman KK, Miljkovic I, Strotmeyer ES, Cummings SR, Kanaya AM, Tylavsky FA, Satterfield S, Ding J, Rimm EB, Trischitta V, Hu FB, Liu Y, Qi L. Genome-wide association analysis identifies TYW3/CRYZ and NDST4 loci associated with circulating resistin levels. *Hum Mol Genet.* 21: 4774-80, 2012.
- 187.** Prudente S, Dallapiccola B, Pellegrini F, Doria A, Trischitta V. Genetic prediction of common diseases. Still no help for the clinical diabetologist! *Nutr Metab Cardiovasc Dis.* 22(11): 929-36, 2012.
- 188.** Maranghi M, Prudente S, D'Erasmus L, Morini E, Ciociola E, Coletta P, Verrienti A, Arciello S, Copetti M, Pellegrini F, Santini SA, Morano S, Filetti S, Trischitta V. The ectonucleotide pyrophosphatase phosphodiesterase 1 (*ENPP1*) K121Q polymorphism modulates the beneficial effect of weight loss on fasting glucose in non diabetic individuals. *Nutr Metab Cardiovasc Dis.* 11: 926-36, 2012.
- 189.** De Cosmo S, Prudente S, Lamacchia O, Lucchesi D, Shah H, Mendonca C, Pucci L, Mercuri L, Gervino EV, Hauser TH, Bailetti D, Penno G, Cignarelli M, Doria A, Trischitta V. The 9p21 coronary artery disease locus and kidney dysfunction in patients with Type 2 diabetes mellitus. *Nephrol Dial Transplant.* 27(12):4411-3, 2012.
- 190.** Chiefari E, Tanyolac S, Iiritano S, Sciacqua A, Capula C, Arcidiacono B, Nocera A, Possidente K, Baudi F, Ventura V, Brunetti G, Brunetti FS, Vero R, Maio R, Greco M, Pavia M, Hodoglulig U, Durlach V, Pullinger CR, Goldfine ID, Perticone F, Foti D, Brunetti A. A polymorphism of HMGA1 is associated with

increased risk of metabolic syndrome and related components. *Sci Rep*; 3:1491, 2013.

191. Barbieri M, Esposito A, Angellotti E, Rizzo MR, Marfella R, Paolisso G. Association of genetic variation in adaptor protein APPL1/APPL2 loci with non-alcoholic fatty liver disease. *PLoS One*.8(8):e71391, 2013.
192. De Cosmo S, Prudente S, Lamacchia O, Pucci L, Lucchesi D, Mendonca C, Bailetti D, Copetti M, Pellegrini F, Cignarelli M, Penno G, Doria A, Trischitta V. The IRS1 G972R polymorphism and glomerular filtration rate in patients with type 2 diabetes of European ancestry. *Nephrol Dial Transplant*. 28(12):3031-4, 2013.
193. Delvecchio M, Ludovico O, Bellacchio E, Stallone R, Palladino T, Mastroianno S, Zelante L, Sacco M, Trischitta V, Carella M. MODY Type 2 P59S GCK mutant: founder effect in South of Italy. *Clin Genet*. 1: 83-7, 2013.
194. Trombetta M., Bonetti S., Boselli M.L., Miccoli R., Trabetti E., Malerba G., Pignatti P.F., Bonora E., Del Prato S., Bonadonna R.C. PPARG2 Pro12Ala and ADAMTS9 rs4607103 as "insulin resistance loci" and "insulin secretion loci" in Italian individuals. The GENFIEV study and the Verona Newly Diagnosed Type 2 Diabetes Study (VNDS) 4. *Acta Diabetol*. 50: 401-408, 2013.
195. Dang MN, Buzzetti R, Pozzilli P. Epigenetics in autoimmune diseases with focus on type 1 diabetes. *Diabetes Metab Res Rev*. 29(1):8-18, 2013.
196. Strollo R, Rizzo P, Spoletini M, Landy R, Hughes C, Ponchel F, Napoli N, Palermo A, Buzzetti R, Pozzilli P, Nissim A. HLA-dependent autoantibodies against post-translationally modified collagen type II in type 1 diabetes mellitus. *Diabetologia*. 56(3):563-72, 2013.
197. Lapice E, Monticelli A, Cocozza S, Pinelli M, Giacco A, Rivellese AA, Cocozza S, Riccardi G, Vaccaro O. The energy intake modulates the association of the -55CT polymorphism of UCP3 with body weight in type 2 diabetic patients. *INT J OBES (Lond)*. 2013.
198. Prudente S, Copetti M, Morini E, Mendonca C, Andreozzi F, Chandalia M, Baratta R; the DIAGRAM consortium, Pellegrini F, Mercuri L, Bailetti D, Abate N, Frittitta L, Sesti G, Florez JC, Doria A, Trischitta V. The SH2B1 obesity locus and abnormal glucose homeostasis: Lack of evidence for association from a meta-analysis in individuals of European ancestry. *Nutr Metab Cardiovasc Dis*. 23:1043-9, 2013.

199. Mannino GC, Greco A, De Lorenzo C, Andreozzi F, Marini MA, Perticone F, Sesti G. A fasting insulin-raising allele at IGF1 locus is associated with circulating levels of IGF-1 and insulin sensitivity. *PLoS One*. 8: e85483, 2013.
200. Qi L, Qi Q, Prudente S, Mendonca C, Andreozzi F, di Pietro N, Sturma M, Novelli V, Mannino GC, Formoso G, Gervino EV, Hauser TH, Muehlschlegel JD, Niewczas MA, Krolewski AS, Biolo G, Pandolfi A, Rimm E, Sesti G, Trischitta V, Hu F, Doria A. Association between a genetic variant related to glutamic acid metabolism and coronary heart disease in individuals with type 2 diabetes. *JAMA* 310: 821-828, 2013.
201. Sesti G, Mannino GC, De Lorenzo C, Greco A, Sciacqua A, Marini MA, Andreozzi F, Perticone F. A functional variant of the dimethylarginine dimethylaminohydrolase-2 gene is associated with chronic kidney disease. *Atherosclerosis* 231: 141–144, 2013.
202. Prudente S, Morini E, Marselli L, Baratta R, Copetti M, Mendonca C, Andreozzi F, Chandalia M, Pellegrini F, Bailetti D, Alberico F, Shah H, Abate N, Sesti G, Frittitta L, Marchetti P, Doria A, Trischitta V. Joint Effect of insulin signaling genes on insulin secretion and glucose homeostasis. *J Clin Endocrinol Metab* 98: E1143–E1147, 2013.
203. Sesti G, Mannino GC, Andreozzi F, Greco A, Perticone M, Sciacqua A, Marini MA, Perticone F. A polymorphism at IGF1 locus is associated with carotid intima media thickness and endothelium-dependent vasodilatation. *Atherosclerosis* 232: 25-30, 2014.
204. Delvecchio M, Di Paola R, Mangiacotti D, Sacco M, Menzaghi C, Trischitta V. Clinical heterogeneity of abnormal glucose homeostasis associated with the HNF4A R311H mutation. *Ital J Pediatr* 40: 58, 2014.
205. Della-Morte D, Palmirotta R, Rehni AK, Pastore D, Capuani B, Pacifici F, De Marchis ML, Dave KR, Bellia A, Fogliame G, Ferroni P, Donadel G, Cacciatore F, Abete P, Dong C, Pileggi A, Roselli M, Ricordi C, Sbraccia P, Guadagni F, Rundek T, Lauro D. Pharmacogenomics and pharmacogenetics of thiazolidinediones: role in diabetes and cardiovascular risk factors. *Pharmacogenomics*. 15(16):2063-82, 2014.
206. Sciacqua A, Presta I, Perticone M, Tassone EJ, Andreozzi F, Quitadamo MC, Sangiuolo FC, Sesti G, Perticone F. 3'-UTR OLR1/LOX-1 gene polymorphism and endothelial dysfunction: molecular and vascular data in never-treated hypertensive patients. *Intern Emerg Med* 9: 273-281, 2014.

207. Pechlaner R., Kiechl S., Willeit P., Demetz E., Haun M., Weger S., Oberhollenzer F., Kronenberg F., Bonora E., Weiss G., Willeit J. Haptoglobin 2-2 genotype is not associated with cardiovascular risk in subjects with elevated glycohemoglobin-results from the Bruneck Study J Am Heart Assoc. 16;3(3):e000732, 2014.
208. Brunetti A, Chiefari E, Foti D. Recent advances in the molecular genetics of type 2 diabetes mellitus. World J Diabetes 5: 128-40, 2014.
209. Pullinger CR, Goldfine ID, Tanyolaç S, Movsesyan I, Faynboym M, Durlach V, Chiefari E, Foti DP, Frost PH, Malloy MJ, Brunetti A, Kane JP. Evidence that an HMGA1 gene variant associates with type 2 diabetes, body mass index, and high-density lipoprotein cholesterol in a Hispanic-American population. Metab Syndr Relat Disord;12: 25-30, 2014.
210. Sambo F., Malovini A., Sandholm N., Stavarachi M., Forsblom C., Mäkinen V.P., Harjutsalo V., Lithovius R., Gordin D., Parkkonen M., Saraheimo M., Thorn L.M., Tolonen N., Wadén J., He B., Osterholm A.M., Tuomilehto J., Lajer M., Salem R.M., McKnight A.J.; GENIE Consortium, Tarnow L, Panduru N.M., Barbarini N., Di Camillo B., Toffolo G., Tryggvason K., Bellazzi R., Cobelli C., FinnDiane Study Group, Groop PH: Novel genetic susceptibility loci for diabetic end-stage renal disease identified through robust naive Bayes classification. Diabetologia 57(8):1611-22, 2014.
211. Giannini C., Dalla Man C., Groop L., Cobelli C., Zhao H., Shaw M.M., Duran E., Pierpont B. Bale A.E Caprio S. Santoro N.:Co-occurrence of risk alleles in or near genes modulating insulin secretion predisposes obese youth to prediabetes. Diabetes Care 37(2):475-82, 2014.
212. Prudente S, Morini E, Lucchesi D, Lamacchia O, Bailetti D, Mercuri L, Alberico F, Copetti M, Pucci L, Fariello S, Giusti L, Cignarelli M, Penno G, De Cosmo S, Trischitta V. IRS1 G972R Missense Polymorphism Is Associated With Failure to Oral Antidiabetes Drugs in White Patients With Type 2 Diabetes From Italy. Diabetes 63(9):3135-40, 2014.
213. Walford GA, Porneala BC, Dauriz M, Vassy JL, Cheng S, Rhee EP, Wang TJ, Meigs JB, Gerszten RE, Florez JC. Metabolite traits and genetic risk provide complementary information for the prediction of future type 2 diabetes Diabetes Care 37(9):2508-14, 2014. *Erratum in: Diabetes. 2018 Jun;67(6):1206..*

- 214.** Cornes BK, Brody JA, Naghmeh N, Morrison AC, Dang HCP, Ahn BS, Wang S, Dauriz M, Barzilay JI, Dupuis J, Florez JC, Coresh J, Gibbs AG, Kao WHL, Liu CT, McKnight B, Muzny D, Pankow JS, Reid JG, White CC, Johnson AD, Wong TY, Psaty BM, Boerwinkle E, Rotter JI, Siscovick DS, Sladek R, and Meigs JB, on behalf of the CHARGE Targeted Sequencing Study Diabetes working group. Association of Levels of Fasting Glucose and Insulin with Rare Variants at the Chromosome 11p11.2-MADD Locus: the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Targeted Sequencing Study. *Circulation: Cardiovascular Genetics*. 7(3):374-82, 2014.
- 215.** Delvecchio M, Ludovico O, Menzaghi C, Di Paola R, Zelante L, Marucci A, Grasso V, Trischitta V, Carella M, Barbetti F, Gallo F, Coccioli MS, Zecchino C, Faienza MF, Cardinale G, Franzese A, Mozzillo E, Iafusco D, Zanfardino A. Low prevalence of HNF1A mutations after molecular screening of multiple MODY genes in 58 Italian families recruited in the pediatric or adult diabetes clinic from a single Italian hospital. *Diabetes Care*. 37: e258-60, 2014.
- 216.** Menzaghi C, Fontana A, Copetti M, Rizza S, Spoto B, Buranasupkajorn P, Tripepi G, Marucci A, Bailetti D, Hastings T, Testa A, Mendonca C, Mallamaci F, De Cosmo S, Bacci S, Federici M, Doria A, Zoccali C, Trischitta V. Joint effect of insulin signaling genes on all-cause mortality. *Atherosclerosis*. 237: 639-44, 2014.
- 217.** Vassy JL, Hivert MF, Dauriz M, Meigs J Response to comment on Vassy et al. Polygenic type 2 diabetes prediction at the limit of common variant detection. *Diabetes* 2014;63:2172-2182”*Diabetes* 63(7): e13, 2014.
- 218.** Dauriz M, Meigs JB. Current Insights into the Joint Genetic Basis of Type 2 Diabetes and Coronary Heart Disease. *Curr Cardiovasc Risk Rep* 8(1):368, 2014.
- 219.** Fares R, Petta S, Lombardi R, Grimaudo S, Dongiovanni P, Pipitone R, Rametta R, Fracanzani AL, Mozzi E, Craxì A, Fargion S, Sesti G, Valenti L. The UCP2 -866 G>A promoter region polymorphism is associated with nonalcoholic steatohepatitis. *Liver Int*. 35:1574-80, 2015.
- 220.** Wessel J, Chu AY, Willems SM, Wang S, Yaghootkar H, Brody JA, Dauriz M, Hivert MF, Raghavan S, Lipovich L, Hidalgo B, Fox K, Huffman JE, An P, Lu Y, Rasmussen-Torvik LJ, Grarup N, Ehm MG, Li L, Baldridge AS, Stančáková A, Abrol R, Besse C, Boland A, Bork-Jensen J, Fornage M, Freitag DF, Garcia ME, Guo X, Hara K, Isaacs A, Jakobsdottir J, Lange LA, Layton JC, Man Li, Hua Zhao J, Meidtner K, Morrison AC, Nalls MA, Peters MJ, Sabater-Lleal M, Schurmann C,

Silveira A, Smith AV, Southam L, Stoiber MH, Strawbridge RJ, Taylor KD, Varga TV, Allin KH, Amin N, Aponte JL, Aung T, Barbieri C, Bihlmeyer NA, Boehnke M, Bombieri C, Bowden DW, Burns SM, Chen Y, Chen Y-D I, Cheng CY, Correa A, Czajkowski J, Dehghan A, Georg B, Ehret, Eiriksdottir G, Escher SA, Farmaki AE, Frånberg M, Gambaro G, Giulianini F, Goddard III WA, Goel A, Gottesman O, Grove ML, Gustafsson S, Hai Y, Hallmans G, Heo J, Hoffmann P, Ikram MK, Jensen RA, Jørgensen ME, Jørgensen T, Karaleftheri M, Khor CC, Kirkpatrick A, Kraja AT, Kuusisto J, Lange EM, Lee IT, Lee WJ, Leong A, Liao J, Liu C, Liu Y, Lindgren CM, Linneberg A, Malerba G, Mamakou V, Marouli E, Maruthur NM, Matchan A, McKean R, McLeod O, Metcalf GA, Mohlke KL, Muzny DM, Ntalla I, Palmer ND, Pasko D, Peter A, Rayner NW, Renström F, Rice K, Sala CF, Sennblad B, Serafetinidis I, Smith JA, Soranzo N, Speliotes EK, Stahl EA, Stirrups K, Nikos Tentolouris, Thanopoulou A, Torres M, Traglia M, Tsafantakis E, Javad S, Yanek LR, Zengini E, Becker DM, Bis JC, Brown JB, L. Cupples A, Hansen T, Ingelsson E, Andrew J. Karter, Carlos Lorenzo, Rasika A. Mathias, Jill M. Norris, Gina M. Peloso, Wayne H.-H. Sheu, Toniolo D, Vaidya D, Varma R, Wagenknecht LE, Boeing H, Erwin P. Bottinger, George Dedoussis, Deloukas P, Ferrannini E, Franco OH, Franks PW, Gibbs RA, Gudnason V, Hamsten A, Harris TB, Andrew T. Hattersley, Hayward C, Hofman A, Jansson JH, Langenberg C, Launer LJ, Levy D, Ben A. Oostra, O'Donnell CJ, O'Rahilly S, Padmanabhan S, Pankow JS, Polasek O, Province MA, Rich SS, Ridker PM, Rudan I, Schulze MB, Smith BH, Uitterlinden AG, Walker M, Watkins H, Wong TY, Zeggini E, Generation Scotland, The EPIC-InterAct Consortium, Laakso M, Borecki IB, Daniel I. Chasman, Oluf Pedersen, Bruce M. Psaty, E. Shyong Tai, van Duijn CM, Wareham NJ, Waterworth DM, Boerwinkle E, Kao WHL, Florez JC, Loos RJJ, Wilson JG, Frayling TM, Siscovick DS, Dupuis J, Rotter JI, Meigs JB, Scott RA, Goodarzi MO. *These authors contributed equally to this work. Rare, low frequency, and common exome chip variants are associated with fasting glucose and influence type 2 diabetes susceptibility - The CHARGE T2D-Glycemia Exome Consortium. *Nature Communications* 6:5897, 2015.

221. Bianco A, Chiefari E, Nobile CG, Foti D, Pavia M, Brunetti A. The Association between HMGA1 rs146052672 Variant and Type 2 Diabetes: A Transethnic Meta-Analysis. *PLoS One*. 10(8):e0136077, 2015.

222. Sentinelli F, Bertoccini L, Barchetta I, Capoccia D, Incani M, Pani MG, Loche S, Angelico F, Arca M, Morini S, Manconi E, Lenzi A, Cossu E, Leonetti F, Baroni MG, Cavallo MG. The vitamin D receptor (VDR) gene rs11568820 variant is associated with type 2 diabetes and impaired insulin secretion in Italian adult subjects, and associates with increased cardio-metabolic risk in children. *Nutr Metab Cardiovasc Dis.* 26(5):407-13, 2016.
223. Sentinelli F, Capoccia D, Bertoccini L, Barchetta I, Incani M, Coccia F, Manconi E, Lenzi A, Cossu E, Leonetti F, Cavallo MG, Baroni MG. Search for Genetic Variant in the Apelin Gene by Resequencing and Association Study in European Subjects. *Genet Test Mol Biomarkers.* 20(2):98-102, 2016.
224. Sentinelli F, Capoccia D, Incani M, Bertoccini L, Severino A, Pani MG, Manconi E, Cossu E, Leonetti F, Baroni MG. The perilipin 2 (PLIN2) gene Ser251Pro missense mutation is associated with reduced insulin secretion and increased insulin sensitivity in Italian obese subjects. *Diabetes Metab Res Rev.*;32(6):550-6, 2016.
225. Trombetta M, Dauriz M, Bonetti S, Travia D, Boselli L, Santi L, Bonora E, Bonadonna RC. Is common genetic variation at IRS1, ENPP1 and TRIB3 loci associated with cardiometabolic phenotypes in type 2 diabetes? An exploratory analysis of the Verona Newly Diagnosed Type 2 Diabetes Study (VNDS) 5. *Nutr Metab Cardiovasc Dis.* 26(3):232-8, 2016.
226. Chiefari E, Ventura V, Capula C, Randazzo G, Scorcio V, Fedele M, Arcidiacono B, Nevolo MT, Bilotta FL, Vitiello M, Palmieri C, Gulletta E, Fusco A, Foti D, Vero R, Brunetti A. A polymorphism of HMGA1 protects against proliferative diabetic retinopathy by impairing HMGA1-induced VEGFA expression. *Sci Rep.* 19;6:39429, 2016.
227. Pacilli A, Prudente S, Copetti M, Fontana A, Mercuri L, Bacci S, Marucci A, Alberico F, Viti R, Palena A, Lamacchia O, Cignarelli M, De Cosmo S, Trischitta V. The PPAR γ 2 P12A polymorphism is not associated with all-cause mortality in patients with type 2 diabetes mellitus. *Endocrine.* 54(1):38-46, 2016
228. Liu CT, Raghavan S, Maruthur N, Kabagambe EK, Hong J, Ng MC, Hivert MF, Lu Y, An P, Bentley AR, Drolet AM, Gaulton KJ, Guo X, Armstrong LL, Irvin MR, Li M, Lipovich L, Rybin DV, Taylor KD, Agyemang C, Palmer ND, Cade BE, Chen WM, Dauriz M, Delaney JA, Edwards TL, Evans DS, Evans MK, Lange LA, Leong A, Liu J, Liu Y, Nayak U, Patel SR, Porneala BC, Rasmussen-Torvik LJ,

Snijder MB, Stallings SC, Tanaka T, Yanek LR, Zhao W, Becker DM, Bielak LF, Biggs ML, Bottinger EP, Bowden DW, Chen G, Correa A, Couper DJ, Crawford DC, Cushman M, Eicher JD, Fornage M, Franceschini N, Fu YP, Goodarzi MO, Gottesman O, Hara K, Harris TB, Jensen RA, Johnson AD, Jhun MA, Karter AJ, Keller MF, Kho AN, Kizer JR, Krauss RM, Langefeld CD, Li X, Liang J, Liu S, Lowe WL Jr, Mosley TH, North KE, Pacheco JA, Peyser PA, Patrick AL, Rice KM, Selvin E, Sims M, Smith JA, Tajuddin SM, Vaidya D, Wren MP, Yao J, Zhu X, Ziegler JT, Zmuda JM, Zonderman AB, Zwinderman AH; AAAG Consortium.; CARE Consortium.; COGENT-BP Consortium.; eMERGE Consortium.; MEDIA Consortium., Adeyemo A, Boerwinkle E, Ferrucci L, Hayes MG, Kardia SL, Miljkovic I, Pankow JS, Rotimi CN, Sale MM, Wagenknecht LE, Arnett DK, Chen YD, Nalls MA; MAGIC Consortium., Province MA, Kao WH, Siscovick DS, Psaty BM, Wilson JG, Loos RJ, Dupuis J, Rich SS, Florez JC, Rotter JI, Morris AP, Meigs JB. Trans-ethnic Meta-analysis and Functional Annotation Illuminates the Genetic Architecture of Fasting Glucose and Insulin. *Am J Hum Genet.* 7;99(1):56-75, 2016.

229. Willems SM, Cornes BK, Brody JA, Morrison AC, Lipovich L, Dauriz M, Chen Y, Liu CT, Rybin DV, Gibbs RA, Muzny D, Pankow JS, Psaty BM, Boerwinkle E, Rotter JI, Siscovick DS, Vasan RS, Kaplan RC, Isaacs A, Dupuis J, van Duijn CM, Meigs JB Association of the IGF1 gene with fasting insulin levels. *Eur J Hum Genet.* 24(9):1337-43, 2016.
230. Lapice E, Monticelli A, Cocozza S, Pinelli M, Massimino E, Giacco A, Rivellese AA, Cocozza S, Riccardi G, Vaccaro O. The combination of UCP3-55CT and PPAR γ 2Pro12Ala polymorphisms affects BMI and substrate oxidation in two diabetic populations. *Nutr Metab Cardiovasc Dis.* ;26(5):400-6, 2016.
231. Politi C, Ciccacci C, D'Amato C, Novelli G, Borgiani P, Spallone V. Recent advances in exploring the genetic susceptibility to diabetic neuropathy. *Diabetes Res Clin Pract.* 120:198-208, 2016.
232. Sandholm N, Van Zuydam N, Ahlqvist E, Juliusdottir T, Deshmukh HA, Rayner NW, Di Camillo B, Forsblom C, Fadista J, Ziemek D, Salem RM, Hiraki LT, Pezzolesi M, Trégouët D, Dahlström E, Valo E, Oskolkov N, Ladenvall C, Marcovecchio ML, Cooper J, Sambo F, Malovini A, Manfrini M, McKnight AJ, Lajer M, Harjutsalo V, Gordin D, Parkkonen M; FinnDiane Study Group., Tuomilehto J, Lyssenko V, McKeigue PM, Rich SS, Brosnan MJ, Fauman E,

- Bellazzi R, Rossing P, Hadjadj S, Krolewski A, Paterson AD; DCCT/EDIC Study Group., Florez JC, Hirschhorn JN, Maxwell AP; GENIE Consortium., Dunger D, Cobelli C, Colhoun HM, Groop L, McCarthy MI, Groop PH; SUMMIT Consortium. The Genetic Landscape of Renal Complications in Type 1 Diabetes. *J Am Soc Nephrol.* 28:557-574, 2017.
- 233.** De Rosa S, Chiefari E, Salerno N, Ventura V, D'Ascoli GL, Arcidiacono B, Ambrosio G, Bilotta FL, Torella D, Foti D, Indolfi C, Brunetti A. HMGA1 is a novel candidate gene for myocardial infarction susceptibility. *Int J Cardiol.* 15;227:331-334, 2017.
- 234.** Bertoccini L, Sentinelli F, Leonetti F, Bailetti D, Capoccia D, Cimini FA, Barchetta I, Incani M, Lenzi A, Cossu E, Cavallo MG, Baroni MG. The vitamin D receptor functional variant rs2228570 (C>T) does not associate with type 2 diabetes mellitus. *Endocr Res.* 42(4):331-335, 2017.
- 235.** Buzzetti R, Prudente S, Copetti M, Dauriz M, Zampetti S, Garofolo M, Penno G, Trischitta V. Clinical worthlessness of genetic prediction of common forms of diabetes mellitus and related chronic complications: A position statement of the Italian Society of Diabetology. *Nutr Metab Cardiovasc Dis.* 27(2):99-114, 2017.
- 236.** Marini MA, Mannino GC, Fiorentino TV, Andreozzi F, Perticone F, Sesti G. A polymorphism at IGF1 locus is associated with anemia. *Oncotarget.* 8(20):32398-32406, 2017.
- 237.** Prudente S, Di Paola R, Copetti M, Lucchesi D, Lamacchia O, Pezzilli S, Mercuri L, Alberico F, Giusti L, Garofolo M, Penno G, Cignarelli M, De Cosmo S, Trischitta V. The rs12917707 polymorphism at the UMOD locus and glomerular filtration rate in individuals with type 2 diabetes: evidence of heterogeneity across two different European populations. *Nephrol Dial Transplant.* 32(10):1718-1722, 2017.
- 238.** Buzzetti R, Prudente S, Copetti M, Dauriz M, Zampetti S, Garofolo M, Penno G, Trischitta V. Clinical worthlessness of genetic prediction of common forms of diabetes mellitus and related chronic complications: A position statement of the Italian Society of Diabetology. *Nutr Metab Cardiovasc Dis.*;27(2):99-114, 2017.
- 239.** Prudente S, Di Paola R, Pezzilli S, Garofolo M, Lamacchia O, Filardi T, Mannino GC, Mercuri L, Alberico F, Scarale MG, Sesti G, Morano S, Penno G, Cignarelli M, Copetti M, Trischitta V. Pharmacogenetics of oral antidiabetes drugs:

- evidence for diverse signals at the IRS1 locus. *Pharmacogenomics J.* 8(3):431-435., 2018.
240. Zusi C, Trombetta M, Bonetti S, Dauriz M, Boselli ML, Trabetti E, Malerba G, Penno G, Zoppini G, Bonora E, Solini A, Bonadonna RC. A renal genetic risk score (GRS) is associated with kidney dysfunction in people with type 2 diabetes. *Diabetes Res Clin Pract.* 144:137-143, 2018.
 241. Iorio A, Ylli D, Polimanti R, Picconi F, Maggio P, Francomano D, Aversa A, Manfellotto D, Fuciarelli M, Frontoni S. Effect of the GSTM1 gene deletion on glycemic variability, sympatho-vagal balance and arterial stiffness in patients with metabolic syndrome, but without diabetes. *Diabetes Res Clin Pract.* 138:158-168, 2018.
 242. Bertoccini L, Bailetti D; SUMMER Study in Diabetes Group, Buzzetti R, Cavallo MG, Copetti M, Cossu E, D'Angelo P, De Cosmo S, Di Mauro L, Leonetti F, Morano S, Morviducci L, Napoli N, Prudente S, Pugliese G, Trischitta V, Baroni MG. Variability in genes regulating vitamin D metabolism is associated with vitamin D levels in type 2 diabetes. *Oncotarget.* 9(79):34911-34918, 2018.
 243. Mannino GC, Fuoco A, Marini MA, Spiga R, Di Fatta C, Mancuso E, Perticone F, Andreozzi F, Sesti G. The polymorphism rs35767 at IGF1 locus is associated with serum urate levels. *Sci Rep.* 8(1):12255, 2018.
 244. Bailetti D, Bertoccini L, Mancina RM, Barchetta I, Capoccia D, Cossu E, Pujia A, Lenzi A, Leonetti F, Cavallo MG, Romeo S, Baroni MG. ANGPTL4 gene E40K variation protects against obesity-associated dyslipidemia in participants with obesity. *Obes Sci Pract.* 5(1):83-90, 2018.
 245. Zusi C, Mantovani A, Olivieri F, Morandi A, Corradi M, Miraglia Del Giudice E, Dauriz M, Valenti L, Byrne CD, Targher G, Maffei C. Contribution of a genetic risk score to clinical prediction of hepatic steatosis in obese children and adolescents. *Dig Liver Dis.* 51(11):1586-1592, 2019.
 246. Nocera D, Menniti M, Belviso S, Bond HM, Lanzillotta D, Spoleti CB, Guagliardi MR, Malatesta P, Trapasso F, Irace C, Perrotti N, Iuliano R. Functional characterization of p.Pro409His variant in HNF1A, a hypomorphic mutation involved in pancreatic β -cell dysfunction. *Acta Diabetol.* 56(8):883-888, 2019.
 247. Sommesse L, Benincasa G, Schiano C, Marfella R, Grimaldi V, Sorriento A, Lucchese R, Fiorito C, Sardu C, Nicoletti GF, Napoli C. Genetic and

- epigenetic-sensitive regulatory network in immune response: a putative link between HLA-G and diabetes. *Expert Rev Endocrinol Metab*, 14:233-241, 2019.
- 248.** Miguel-Escalada I, Bonàs-Guarch S, Cebola I, Ponsa-Cobas J, Mendieta-Esteban J, Atla G, Javierre BM, Rolando DMY, Farabella I, Morgan CC, García-Hurtado J, Beucher A, Morán I, Pasquali L, Ramos-Rodríguez M, Appel EVR, Linneberg A, Gjesing AP, Witte DR, Pedersen O, Grarup N, Ravassard P, Torrents D, Mercader JM, Piemonti L, Berney T, de Koning EJP, Kerr-Conte J, Pattou F, Fedko IO, Groop L, Prokopenko I, Hansen T, Marti-Renom MA, Fraser P, Ferrer J. Human pancreatic islet three-dimensional chromatin architecture provides insights into the genetics of type 2 diabetes. *Nat Genet*, 51:1137-1148, 2019.
- 249.** Mannino GC, Pezzilli S, Averta C, Fuoco A, Spiga R, Mancuso E, Di Fatta C, Perticone F, Prudente S, Trischitta V, Andreozzi F, Sesti G. A functional variant of the dimethylarginine dimethylaminohydrolase-2 gene is associated with myocardial infarction in type 2 diabetic patients. *Cardiovasc Diabetol*, 18:102, 2019.
- 250.** Haydar S, Grigorescu F, Vintilă M, Cogne Y, Lautier C, Tutuncu Y, Brun JF, Robine JM, Pugeat M, Normand C, Poucheret P, Gheorghiu ML, Georgescu C, Badiu C, Băculescu N, Renard E, Ylli D, Badiou S, Sutra T, Cristol JP, Mercier J, Gomis R, Macias JM, Litvinov S, Khusnutdinova E, Poiana C, Pasquali R, Lauro D, Sesti G, Prudente S, Trischitta V, Tsatsoulis A, Abdelhak S, Barakat A, Zenati A, Ylli A, Satman I, Kanninen T, Renato Y, Missoni S. Fine-scale haplotype mapping of MUT, AACS, SLC6A15 and PRKCA genes indicates association with insulin resistance of metabolic syndrome and relationship with branched chain amino acid metabolism or regulation. *PLoS One*, 14:e0214122, 2019.
- 251.** Mannino GC, Andreozzi F, Sesti G. Pharmacogenetics of type 2 diabetes mellitus, the route toward tailored medicine. *Diabetes Metab Res Rev*, 35:e3109, 2019.
- 252.** Latini A, Spallone V, D'Amato C, Novelli G, Borgiani P, Ciccacci C. A common polymorphism in MIR155 gene promoter region is associated with a lower risk to develop type 2 diabetes. *Acta Diabetol*, 56:717-718, 2019.
- 253.** Dwivedi OP, Lehtovirta M, Hastoy B, Chandra V, Krentz NAJ, Kleiner S, Jain D, Richard AM, Abaitua F, Beer NL, Grotz A, Prasad RB, Hansson O,

- Ahlqvist E, Krus U, Artner I, Suoranta A, Gomez D, Baras A, Champon B, Payne AJ, Moralli D, Thomsen SK, Kramer P, Spiliotis I, Ramracheya R, Chabosseau P, Theodoulou A, Cheung R, van de Bunt M, Flannick J, Trombetta M, Bonora E, Wolheim CB, Sarelin L, Bonadonna RC, Rorsman P, Davies B, Brosnan J, McCarthy MI, Otonkoski T, Lagerstedt JO, Rutter GA, Gromada J, Gloyn AL, Tuomi T, Groop L. Loss of ZnT8 function protects against diabetes by enhanced insulin secretion. *Nat Genet*, 51:1596-1606, 2019.
- 254.** Ramos-Rodríguez M, Raurell-Vila H, Colli ML, Alvelos MI, Subirana-Granés M, Juan-Mateu J, Norris R, Turatsinze JV, Nakayasu ES, Webb-Robertson BM, Inshaw JRJ, Marchetti P, Piemonti L, Esteller M, Todd JA, Metz TO, Eizirik DL, Pasquali L. The impact of proinflammatory cytokines on the β -cell regulatory landscape provides insights into the genetics of type 1 diabetes. *Nat Genet*, 51 (11):1588-1595, 2019.
- 255.** Laurenti MC, Dalla Man C, Varghese RT, Andrews JC, Rizza RA, Matveyenko A, De Nicolao G, Cobelli C, Vella A. Diabetes-associated genetic variation in TCF7L2 alters pulsatile insulin secretion in humans. *JCI Insight*. 5(7):e136136, 2020.
- 256.** Galderisi A, Tricò D, Dalla Man C, Santoro N, Pierpont B, Groop L, Cobelli C, Caprio S. Metabolic and Genetic Determinants of Glucose Shape After Oral Challenge in Obese Youths: A Longitudinal Study. *J Clin Endocrinol Metab*. 105(2):534-42, 2020.
- 257.** Moffa S, Mazzuccato G, De Bonis M, De Paolis E, Onori ME, Pontecorvi A, Urbani A, Giaccari A, Capoluongo E, Minucci A. Identification of two novel LDLR variants by Next Generation Sequencing. *Ann Ist Super Sanita*. 56(1):122-127, 2020.
- 258.** Pipino C, Shah H, Prudente S, Di Pietro N, Zeng L, Park K, Trischitta V, Pennathur S, Pandolfi A, Doria A. Association of the 1q25 Diabetes-Specific Coronary Heart Disease Locus With Alterations of the γ -Glutamyl Cycle and Increased Methylglyoxal Levels in Endothelial Cells. *Diabetes*. 69(10):2206-2216, 2020.
- 259.** Morieri ML, Shah HS, Sjaarda J, Lenzini PA, Campbell H, Motsinger-Reif AA, Gao H, Lovato L, Prudente S, Pandolfi A, Pezzolesi MG, Sigal RJ, Paré G, Marcovina SM, Rotroff DM, Patorno E, Mercuri L, Trischitta V, Chew EY, Kraft P, Buse JB, Wagner MJ, Cresci S, Gerstein HC, Ginsberg HN,

- Mychaleckyj JC, Doria A. PPARA Polymorphism Influences the Cardiovascular Benefit of Fenofibrate in Type 2 Diabetes: Findings From ACCORD-Lipid. *Diabetes*. 69(4):771-783, 2020.
- 260.** Masulli M, Della Pepa G, Cocozza S, Capasso M, Pignataro P, Vitale M, Gastaldelli A, Russo M, Dolce P, Riccardi G, Rivellese AA, Vaccaro O. The Pro12Ala polymorphism of PPAR γ 2 modulates beta cell function and failure to oral glucose-lowering drugs in patients with type 2 diabetes. *Diabetes Metab Res Rev*. 11:e3392, 2020.
- 261.** van Zuydam NR, Ladenvall C, Voight BF, Strawbridge RJ, Fernandez-Tajes J, Rayner NW, Robertson NR, Mahajan A, Vlachopoulou E, Goel A, Kleber ME, Nelson CP, Kwee LC, Esko T, Mihailov E, Mägi R, Milani L, Fischer K, Kanoni S, Kumar J, Song C, Hartiala JA, Pedersen NL, Perola M, Gieger C, Peters A, Qu L, Willems SM, Doney ASF, Morris AD, Zheng Y, Sesti G, Hu FB, Qi L, Laakso M, Thorsteinsdottir U, Grallert H, van Duijn C, Reilly MP, Ingelsson E, Deloukas P, Kathiresan S, Metspalu A, Shah SH, Sinisalo J, Salomaa V, Hamsten A, Samani NJ, März W, Hazen SL, Watkins H, Saleheen D, Morris AP, Colhoun HM, Groop L, McCarthy MI, Palmer CNA; SUMMIT Steering Committee; CARDIOGRAMplusC4D Steering Committee*. Genetic Predisposition to Coronary Artery Disease in Type 2 Diabetes Mellitus. *Circ Genom Precis Med*. 13(6):e002769, 2020.
- 262.** Tavaglione F, Targher G, Valenti L, Romeo S. Human and molecular genetics shed lights on fatty liver disease and diabetes conundrum. *Endocrinol Diabetes Metab*. 3(4):e00179, 2020.
- 263.** Mantovani A, Taliento A, Zusi C, Baselli G, Prati D, Granata S, Zaza G, Colecchia A, Maffei C, Byrne CD, Valenti L, Targher G. PNPLA3 I148M gene variant and chronic kidney disease in type 2 diabetic patients with NAFLD: Clinical and experimental findings. *Liver Int*. 40(5):1130-1141, 2020.