

1. Annuzzi G, Vaccaro O, Caprio S, Di Bonito P, Caso P, Riccardi G, Rivellese A. Association between low habitual physical activity and impaired glucose tolerance. *Clinical Physiology*, 5(1):63-70, 1985.
2. Annuzzi G, Jansson E, Kaijser L, Holmquist L, Carlson LA. Increased removal rate of exogenous triglycerides after prolonged exercise: time course and effect of exercise duration. *Metabolism*, 36(5):438-443, 1987.
3. Annuzzi G, Riccardi G, Capaldo B, Kaijser L. Increased insulin-stimulated glucose uptake by exercised human muscles one day after prolonged physical exercise. *European Journal of Clinical Investigation* 21:6-12, 1991
4. Francescato M.P., Geat M., Fusi S., Stupar G., Noacco C., Cattin L.: Carbohydrate requirement and insulin concentration during moderate exercise in type 1 diabetic patients. *Metabolism*; 53:1126-1130, 2004.
5. Balducci S, Leonetti F, Di Mario U, Fallucca F. Is a long-term aerobic plus resistance training program feasible for and effective on metabolic profiles in type 2 diabetic patients? *Diabetes Care*. 27(3):841-2, 2004.
6. Francescato M.P., Cattin L., Geat M., Tosoratti E., Lazzer S., Noacco C., di Prampero P.E.: Glucose pulse. A simple method to estimate the amount of carbohydrates oxidized during exercise in type 1 diabetic patients. *Diabetes Care*; 28(8): 2028-2030, 2005.
7. Finelli C, Gallipoli P, Celentano E, Cacace G, Saldalamacchia G, De Caprio C, Contaldo F, Pasanisi F. Assessment of physical activity in an outpatient obesity clinic in southern Italy: results from a standardized questionnaire. *Nutr Metab Cardiovasc Dis*. 16(3):168-73, 2006.
8. Cohen O., Basu R., Bock G., Dalla Man C., Campioni M., Basu A., Toffolo G., Cobelli C., Rizza R.A.: Prediction of postprandial glycemic exposure: utility of fasting and two hour glucose measurements alone and in combination with assessment of body composition, fitness and strength. *Diabetes Care* 29:2708-13, 2006.
9. Balducci S, Alessi E, Cardelli P, Cavallo S, Fallucca F, Pugliese G. Effects of different modes of exercise training on glucose control and risk factors for complications in type 2 diabetic patients: a meta-analysis: response to Snowling and Hopkins. *Diabetes Care*.; 30:e25, 2007.
10. Benso A, Broglio F, Aimaretti G, Lucatello B, Lanfranco F, Ghigo E, Grottoli S. Endocrine and metabolic responses to extreme altitude and physical exercise in climbers. *Eur J Endocrinol*. 157(6):733-40, 2007.
11. Balducci S, Zanuso S, Massarini M, Corigliano G, Nicolucci A, Missori S, Cavallo S,

- Cardelli P, Alessi E, Pugliese G, Fallucca F; The Italian Diabetes and Exercise Study (IDES): design and methods for a prospective Italian multicentre trial of intensive lifestyle intervention in people with type 2 diabetes and the metabolic syndrome. Italian Diabetes Exercise Study (IDES) Group. *Nutr Metab Cardiovasc Dis.* 18(9):585-95, 2008.
12. Zanuso S, Jimenez A, Pugliese G, Corigliano G, Balducci S. Exercise for the management of type 2 diabetes: a review of the evidence. *Acta Diabetol.*; 25 Suppl 1:S24-28, 2009.
 13. Dalla Man C., Breton M.D., Cobelli C.: Physical activity into the meal glucose-insulin model of type 1 diabetes: in silico studies. *J Diabetes Sci. Technol.* 3:56-67, 2009.
 14. Pugliese G, Zanuso S, Alessi E, Simonelli P, Fallucca S, Fallucca F, Balducci S. Self glucose monitoring and physical exercise in diabetes. *Diabetes Metab Res Rev* 25 Suppl 1:S11-S17, 2009.
 15. Balducci S, Zanuso S, Fernando F, Fallucca S, Fallucca F, Pugliese G; Italian Diabetes Exercise Study (IDES) Group. Physical activity/exercise training in type 2 diabetes. The role of the Italian Diabetes and Exercise Study. *Diabetes Metab Res Rev* 25 Suppl 1:S29-S33, 2009.
 16. Zanuso S, Balducci S, Gutierrez AJ, Bergamin M, Pugliese G. Similar energy expenditure from resistance training at moderate and vigorous intensity in subjects with type 2 diabetes. *Diabetes Res Clin Pract.*; 85:e40-e41, 2009.
 17. Perseghin G, Lattuada G, Ragona F, Alberti G, La Torre A, Luzi L. Free leptin index and thyroid function in male highly trained athletes. *Eur J Endocrinol.* 161(6):871-6, 2009.
 18. Perseghin G, De Cobelli F, Esposito A, Belloni E, Lattuada G, Canu T, Invernizzi PL, Ragona F, La Torre A, Scifo P, Alberti G, Del Maschio A, Luzi L. Left ventricular function and energy metabolism in middle-age men undergoing long lasting sustained aerobic oxidative training. *Heart* 95(8):630-5, 2009.
 19. Fallucca F, Pozzilli P. Physical exercise, public health and quality of life in diabetes. *Diabetes Metab Res Rev.* 25: S1-3, 2009.
 20. IDES Study Investigators. Effect of an intensive exercise intervention strategy on modifiable cardiovascular risk factors in subjects with type 2 diabetes mellitus: a randomized controlled trial: the Italian Diabetes and Exercise Study (IDES). *Arch Intern Med* 170:1794-1803, 2010.
 21. Lauriola V., Codella R., Lattuada G., Caumo A., Capitelli E., Lazzarin A., Luzi L Effects of endurance exercise training on metabolic and inflammatory parameters in HIV-1-infected patients with secondary lipodystrophy and diabetes. *Sport Sci Health* 1:23–26, 2010.
 22. Benedini S., Caimi A., Alberti G, Terruzzi I., Dellerma N., La Torre A., Luzi L. Increase in

homocysteine levels after a half-marathon running: a detrimental metabolic effect of sport? *Sport Sci Health* 1:35–42, 2010.

23. Benevento D, Bizzarri C, Pitocco D, Crinò A, Moretti C, Spera S, Tubili C, Costanza F, Maurizi A, Cipolloni L, Cappa M, Pozzilli P; IMDIAB Group. Computer use, free time activities and metabolic control in patients with type 1 diabetes. *Diabetes Res Clin Pract.* 88(3):e32-4, 2010.
24. Balducci S, Zanuso S, Nicolucci A, Fernando F, Cavallo S, Cardelli P, Fallucca S, Alessi E, Letizia C, Jimenez A, Fallucca F, Pugliese G. Anti-inflammatory effect of exercise training in subjects with type 2 diabetes and the metabolic syndrome is dependent on exercise modalities and independent of weight loss. *Nutr Metab Cardiovasc Dis.* 20:608-617, 2010.
25. Pedone C, Napoli N, Pozzilli P, Lauretani F, Bandinelli S, Ferrucci L, Antonelli-Incalzi R. Quality of diet and potential renal acid load as risk factors for reduced bone density in elderly women. *Bone.* 46(4):1063-7, 2010.
26. Nicolucci A, Balducci S, Cardelli P, Zanuso S, Pugliese G; Italian Diabetes Exercise Study (IDES) Investigators. Improvement of quality of life with supervised exercise training improves quality of life in subjects with type 2 diabetes mellitus. *Arch Intern Med.*; 171:1951-1953, 2011.
27. Balducci S, Zanuso S, Pugliese G, Church TS, Sigal RJ. Diet or diet plus physical activity in patients with early type 2 diabetes. *Lancet.* 378:2066, 2011.
28. Francescato M.P., Geat M., Accardo A., Blokar M., Cattin L., Noacco C.: Exercise and Glycemic Imbalances: a Situation-Specific Estimate of Glucose Supplement. *Med Sci Sports Exerc*; 43(1):2-11, 2011.
29. Terruzzi I., Senesi P., Montesano A., La Torre A., Alberti G., Benedini S., Caumo A., Fermo I., Luzi L. Genetic polymorphisms of the enzymes involved in DNA methylation and synthesis in elite athletes. *Physiological Genomics* 43 (16): 965-73, 2011.
30. Balducci S, Zanuso S, Cardelli P, Salerno G, Fallucca S, Nicolucci A, Pugliese G, for the Italian Diabetes Exercise Study (IDES) Investigators. Supervised Exercise training counterbalances the adverse effects of insulin therapy in overweight/obese subjects with type 2 diabetes. *Diabetes Care.*; 35:39-41, 2012.
31. IDES Study Investigators. Relationship of exercise volume to improvements of quality of life with supervised exercise training in patients with type 2 diabetes in a randomized controlled trial: the Italian Diabetes and Exercise Study (IDES). *Diabetologia*;55:579-588, 2012.
32. Manohar C., Levine J. A, Nandy D. K., Saad A., Dalla Man C., . McCrady-Spitzer S. K,

- Basu R., Cobelli C., Carter R. E., Basu A., Kudva Y. C.: The effect of walking on postprandial glycemic excursion in patients with type 1 diabetes and healthy people. *Diabetes Care* 38:2493-2499, 2012.
33. Balducci S, Zanuso S, Cardelli P, Salvi L, Bazuro A, Pugliese L, Maccora C, Iacobini C, Conti F, Nicolucci A, Pugliese G. Effect of high- versus low-intensity supervised aerobic and resistance training on modifiable cardiovascular risk factors in type 2 diabetes. The Italian Diabetes and Exercise Study (IDES). *PLoSOne*. 7:e49297, 2012.
 34. Balducci S, Zanuso S, Cardelli P, Salvi L, Mazzitelli G, Bazuro A, Iacobini C, Nicolucci A, Pugliese G; for the Italian Diabetes Exercise Study (IDES) Investigators. Changes in physical fitness predict improvements in modifiable cardiovascular risk factors independently of body weight loss in subjects with type 2 diabetes participating in the Italian Diabetes and Exercise Study (IDES). *Diabetes Care*.; 35:1347-1354, 2012.
 35. Nicolucci A, Balducci S, Cardelli P, Cavallo S, Fallucca S, Bazuro A, Simonelli P, Iacobini C, Zanuso S, Pugliese G; Italian Diabetes Exercise Study (IDES) Investigators. Relationship of exercise volume to improvements of quality of life with supervised exercise training in patients with type 2 diabetes in a randomised control trial: the Italian Diabetes and Exercise Study (IDES). *Diabetologia*.; 55:579-588, 2012.
 36. Benedini S., Capitelli E., Casiraghi E., Codella R., Luzi L. A “single-physician” model for diabetes care: best practice for life-style changes? *Sport Sciences For Health*, vol. 7, p. 81-82, 2012.
 37. Bacchi E., Negri C., Trombetta M., Zanolin M.E., Lanza M., Bonora E., Moghetti P. Differences in the acute effects of aerobic and resistance exercise in subjects with type 2 diabetes: results from the RAED2 randomized trial. *PLoS One* 7:e49937, 2012.
 38. Geat M., Stel G., Poser S., Driussi C., Stenner E., Francescato M.P.: Whole-body glucose oxidation rate during prolonged exercise in type 1 diabetic patients under usual life conditions. *Metabolism*; 62(6):836–844, 2013.
 39. Short K.R., Pratt L.V., Teague A.M., Man C.D., Cobelli C.: Postprandial improvement in insulin sensitivity after a single exercise session in adolescents with low aerobic fitness and physical activity. *Pediatric Diabetes* 14:129-37, 2013.
 40. Schiavon M., Hinshaw L., Mallad A., Dalla Man C., Sparacino G., Johnson M.L., Carter R.E., Basu R., Kudva Y.C., Cobelli C., Basu A.: Postprandial glucose fluxes and insulin sensitivity during exercise: a study in healthy individuals. *Am. J. Physiol. Endocrinol. Metab.* 305:E557- E566, 2013.
 41. Sacchetti M, Balducci S, Bazzucchi I, Carlucci F, di Palumbo AS, Haxhi J, Conti F, Di

- Biase N, Calandriello E, Pugliese G. Neuromuscular dysfunction in diabetes: Role of nerve impairment and training status. *Med Sci Sports Exerc.*; 45:52-59, 2013.
42. Francescato M.P., Stel G., Geat M., Cauci S.. Oxidative stress in patients with type 1 diabetes mellitus: is it affected by a single bout of prolonged exercise? *PlosOne*; 9(6):e99062, 2014.
 43. Bozzetto L, Annuzzi G, Costabile G, Costagliola L, Giorgini M, Alderisio A, Strazzullo A, Patti L, Cipriano P, Mangione A, Vitelli A, Vigorito C, Riccardi G, Rivellese AA. A CHO/fibre diet reduces and a MUFA diet increases postprandial lipaemia in type 2 diabetes: no supplementary effects of low-volume physical training. *Acta Diabetol.* 51(3):385-93, 2014.
 44. Annuzzi G, Rivellese AA, Bozzetto L, Riccardi G. The results of Look AHEAD do not row against the implementation of lifestyle changes in patients with type 2 diabetes. *Nutr Metab Cardiovasc Dis.* 24(1):4-9, 2014.
 45. Centis E, Trento M, Dei Cas A, Pontiroli AE, De Feo P, Bruno A, Sasdelli AS, Arturi F, Strollo F, Vigili De' Kreutzenberg S, Invitti C, Di Bonito P, Di Mauro M, Pugliese G, Molteni A, Marchesini G. Stage of change and motivation to healthy diet and habitual physical activity in type 2 diabetes. *Acta Diabetol.*,51:559-66, 2014.
 46. Senesi P, Luzi L, Montesano A, Terruzzi I. DNA demethylation enhances myoblasts hypertrophy during the late phase of myogenesis activating the IGF-I pathway. *Endocrine.* 47(1):244-54, 2014.
 47. Balducci S, Vulpiani MC, Pugliese L, D'Errico V, Menini S, Salerno G, Gargiulo L, Ferretti A, Pugliese G. Effect of supervised exercise training on musculoskeletal symptoms and function in patients with type 2 diabetes: the Italian Diabetes Exercise Study (IDES). *Acta Diabetol.*; 51:647-54, 2014.
 48. Balducci S, Sacchetti M, Haxhi J, Orlando G, D'Errico V, Fallucca S, Menini S, Pugliese G. Physical exercise as therapy for type 2 diabetes mellitus. *Diabetes Metab Res Rev.* ; 30 Suppl 1:13-23, 2014.
 49. Soare A, Khazrai YM, Del Toro R, Roncella E, Fontana L, Fallucca S, Angeletti S, Formisano V, Capata F, Ruiz V, Porrata C, Skrami E, Gesuita R, Manfrini S, Fallucca F, Pianesi M, Pozzilli P. The effect of the macrobiotic Ma-Pi 2 diet vs. the recommended diet in the management of type 2 diabetes: the randomized controlled MADIAB trial. *NutrMetab (Lond).*;11:39, 2014.
 50. Palermo A, Maggi D, Maurizi AR, Pozzilli P, Buzzetti R. Prevention of type 2 diabetes mellitus: is it feasible? *Diabetes Metab Res Rev. Suppl* 1:4-12, 2014.

51. Khazrai YM, Maddaloni E, Altomare M, Cacciapaglia F, Pozzilli P. New technologies aiding dietary programmes for weight control: the oral glucose spray. *Endocrine*. 45(2):288-92, 2014.
52. Pugliese G, Balducci S. NAVIGATOR: physical activity for cardiovascular health? *Lancet*.; 383:1022-1023, 2014.
53. Lombardi C, Carubelli V, Lazzarini V, Vizzardi E, Bordonali T, Ciccarese C, Castrini AI, Dei Cas A, Nodari S, Metra M. Effects of oral administration of orodispersible levo-carnosine on quality of life and exercise performance in patients with chronic heart failure. *Nutrition*.;31(1):72-8, 2015.
54. Balducci S, Sacchetti M, Haxhi J, Orlando G, Zanuso S, Cardelli P, Cavallo S, D'Errico V, Ribaud MC, Di Biase N, Salvi L, Vitale M, Bollanti L, Conti FG, Nicolucci A, Pugliese G; Italian Diabetes and Exercise Study 2 (IDES_2) Investigators. The Italian Diabetes and Exercise Study 2 (IDES-2): a long-term behavioral intervention for adoption and maintenance of a physically active lifestyle. *Trials*. 16:569, 2015.
55. Palermo A, Strollo R, Maddaloni E, Tuccinardi D, D'Onofrio L, Briganti SI, Defeudis G, De Pascalis M, Lazzaro MC, Colleluori G, Manfrini S, Pozzilli P, Napoli N. Irisin is associated with osteoporotic fractures independently of bone mineral density, body composition or daily physical activity. *Clin Endocrinol (Oxf)*. 82:615-9, 2015.
56. Francescato MP, Stel G, Stenner E, Geat M. Prolonged exercise in type 1 diabetes: performance of a customizable algorithm to estimate the carbohydrate supplements to minimize glycemic imbalances. *PLoS One*. 10(4):e0125220, 2015.
57. Codella R, Luzi L, Inverardi L, Ricordi C. The anti-inflammatory effects of exercise in the syndromic thread of diabetes and autoimmunity. *Eur Rev Med Pharmacol Sci*. 19(19):3709-22, 2015.
58. Balducci S, Cardelli P, Pugliese L, D'Errico V, Haxhi J, Alessi E, Iacobini C, Menini S, Bollanti L, Conti FG, Nicolucci A, Pugliese G; Italian Diabetes Exercise Study (IDES) Investigators. Volume-dependent effect of supervised exercise training on fatty liver and visceral adiposity index in subjects with type 2 diabetes The Italian Diabetes Exercise Study (IDES). *Diabetes Res Clin Pract*. 109(2):355-63, 2015.
59. Bacchi E, Bonin C, Zanolini ME, Zambotti F, Livornese D, Donà S, Tosi F, Baldisser G, Ihnatava T, Di Sarra D, Bonora E, Moghetti P. Physical Activity Patterns in Normal-Weight and Overweight/Obese Pregnant Women. *PLoS One*. 9;11(11):e0166254, 2016.
60. Orlando G, Balducci S, Bazzucchi I, Pugliese G, Sacchetti M. Neuromuscular dysfunction in type 2 diabetes: underlying mechanisms and effect of resistance training. *Diabetes Metab*

Res Rev. 32(1):40-50, 2016.

61. Buoite Stella AB, Assaloni R, Tonutti L, Manca E, Tortul C, Candido R, Francescato MP. Strategies Used by Patients with Type 1 Diabetes to Avoid Hypoglycemia in a 24×1-Hour Marathon: Comparison with the Amounts of Carbohydrates Estimated by a Customizable Algorithm. *Can J Diabetes*. 41(2):184-189. 2017
62. Adamo M, Codella R, Casiraghi F, Ferrulli A, Macrì C, Bazzigaluppi E, Terruzzi I, Inverardi L, Ricordi C, Luzi L. Active Subjects with Autoimmune Type 1 Diabetes Have Better Metabolic Profiles than Sedentary Controls. *Cell Transplant*. 24;26(1):23-32, 2017.
63. Accattato F, Greco M, Pullano SA, Carè I, Fiorillo AS, Pujia A, Montalcini T, Foti DP, Brunetti A, Gulletta E Effects of acute physical exercise on oxidative stress and inflammatory status in young, sedentary obese subjects. *PLoS One*. 5;12(6):e0178900, 2017.
64. Codella R, Terruzzi I, Luzi L. Why should people with type 1 diabetes exercise regularly? *Acta Diabetol*. 54(7):615-630, 2017.
65. Adamo M, Codella R, Casiraghi F, Ferrulli A, Macrì C, Bazzigaluppi E, Terruzzi I, Inverardi L, Ricordi C, Luzi L. Active Subjects With Autoimmune Type 1 Diabetes Have Better Metabolic Profiles Than Sedentary Controls. *Cell Transplant*. 26(1):23-32, 2017.
66. Codella R, Adamo M, Maffi P, Piemonti L, Secchi A, Luzi L. Ultra-marathon 100 km in an islet-transplanted runner. *Acta Diabetol*. 54(7):703-706, 2017.
67. Benedini S, Dozio E, Invernizzi PL, Vianello E, Banfi G, Terruzzi I, Luzi L, Corsi Romanelli MM Irisin: A Potential Link between Physical Exercise and Metabolism-An Observational Study in Differently Trained Subjects, from Elite Athletes to Sedentary People. *J Diabetes Res*. 2017:1039161, 2017.
68. Balducci S, D'Errico V, Haxhi J, Sacchetti M, Orlando G, Cardelli P, Vitale M, Bollanti L, Conti F, Zanuso S, Nicolucci A, Pugliese G; Italian Diabetes and Exercise Study 2 (IDES_2) Investigators. Effect of a Behavioral Intervention Strategy for Adoption and Maintenance of a Physically Active Lifestyle: The Italian Diabetes and Exercise Study 2 (IDES_2): A Randomized Controlled Trial. *Diabetes Care*. 40(11):1444-1452, 2017.
69. Codella R, Terruzzi I, Luzi L. Sugars, exercise and health. *J Affect Disord* 15;224:76-86, 2017.
70. Dauriz M, Bacchi E, Boselli L, Santi L, Negri C, Trombetta M, Bonadonna RC, Bonora E, Moghetti P. Association of free-living physical activity measures with metabolic phenotypes in type 2 diabetes at the time of diagnosis. The Verona Newly Diagnosed Type 2 Diabetes Study (VNDS). *Nutr Metab Cardiovasc Dis*. 28(4):343-351, 2018.
71. Indelicato L, Dauriz M, Bacchi E, Donà S, Santi L, Negri C, Cacciatori V, Bonora E,

- Nouwen A, Moghetti P. Sex differences in the association of psychological status with measures of physical activity and sedentary behaviour in adults with type 2 diabetes. *Acta Diabetol.* 55(6):627-635, 2018.
72. Bellavere F, Cacciatori V, Bacchi E, Gemma ML, Raimondo D, Negri C, Thomaseth K, Muggeo M, Bonora E, Moghetti P. Effects of aerobic or resistance exercise training on cardiovascular autonomic function of subjects with type 2 diabetes: A pilot study. *Nutr Metab Cardiovasc Dis.* 28(3):226-233, 2018.
 73. Codella R, Ialacqua M, Terruzzi I, Luzi L. May the force be with you: why resistance training is essential for subjects with type 2 diabetes mellitus without complications. *Endocrine.* 62(1):14-25, 2018.
 74. Codella R, Luzi L, Terruzzi I. Exercise has the guts: How physical activity may positively modulate gut microbiota in chronic and immune-based diseases. *Dig Liver Dis.* 50(4):331-341, 2018.
 75. Buoite Stella A, Yardley J, Francescato MP, Morrison SA Fluid Intake Habits in Type 1 Diabetes Individuals during Typical Training Bouts. *Ann Nutr Metab.* 73(1):10-18, 2018.
 76. Zurlo F, Trevisan C, Vitturi N, Ravussin E, Salvò C, Carraro S, Siffi M, Iob I, Saller A, Previato L, Sergi G, de Kreutzenberg S, Maran A, Avogaro A. One-year caloric restriction and 12-week exercise training intervention in obese adults with type 2 diabetes: emphasis on metabolic control and resting metabolic rate. *J Endocrinol Invest,* 42(12):1497-1507, 2019.
 77. Nasi M, Patrizi G, Pizzi C, Landolfo M, Boriani G, Dei Cas A, Cicero AFG, Fogacci F, Rapezzi C, Sisca G, Capucci A, Vitolo M, Galiè N, Borghi C, Berrettini U, Piepoli M, Mattioli AV. The role of physical activity in individuals with cardiovascular risk factors: an opinion paper from Italian Society of Cardiology-Emilia Romagna-Marche and SIC-Sport. *J Cardiovasc Med (Hagerstown),* 20(10):631-639, 2019.
 78. Francescato MP, Cettolo V. The "independent breath" algorithm: assessment of oxygen uptake during exercise. *Eur J Appl Physiol,* 119(2):495-508, 2019.
 79. Koschate J, Cettolo V, Hoffmann U, Francescato MP. Breath-by-breath oxygen uptake during running: Effects of different calculation algorithms. *Exp Physiol,* 104(12):1829-1840, 2019.
 80. Benedini S, Codella R, Caumo A, Terruzzi I, Luzi L. Single bout of low-intensity exercise produces modestly favorable changes in glycemic and lipidemic profiles after ingestion of non-isoglucidic breakfasts. *Nutrition,* 58:57-64, 2019.
 81. Balducci S, D'Errico V, Haxhi J, Sacchetti M, Orlando G, Cardelli P, Vitale M, Bollanti L, Conti F, Zanuso S, Lucisano G, Nicolucci A, Pugliese G. Italian Diabetes and Exercise

- Study 2 (IDES_2) Investigators. Effect of a Behavioral Intervention Strategy on Sustained Change in Physical Activity and Sedentary Behavior in Patients With Type 2 Diabetes: The IDES_2 Randomized Clinical Trial. *JAMA*, 321:880-890, 2019.
- 82.** Haxhi J, Balducci S, Pugliese G. Invest in METs, Not in Meds. *Am. J. Med*, 132:e756, 2019.
- 83.** Balducci S, Conti F, Sacchetti M, Russo CR, Argento G, Haxhi J, Orlando G, Rapisarda G, D'Errico V, Cardelli P, Pugliese L, Laghi A, Vitale M, Bollanti L, Zanuso S, Nicolucci A, Pugliese G; SWEET BONE Investigators. Study to Weigh the Effect of Exercise Training on BONE quality and strength (SWEET BONE) in type 2 diabetes: study protocol for a randomised clinical trial. *BMJ Open*, 9:e027429, 2019.
- 84.** Francescato MP, Ajčević M, Accardo A. Carbohydrate Requirement for Exercise in Type 1 Diabetes: Effects of Insulin Concentration. *J Diabetes Sci Technol*, 14(6):1116-1121, 2020.
- 85.** Ajčević M, Candido R, Assaloni R, Accardo A, Francescato MP. Personalized Approach for the Management of Exercise-Related Glycemic Imbalances in Type 1 Diabetes: Comparison with Reference Method. *J Diabetes Sci Technol*. [Epub ahead of print], 2020.
- 86.** Romeres D, Olson K, Carter R, Cobelli C, Dalla Man C, Basu A, Basu R. Hyperglycemia But Not Hyperinsulinemia Is Favorable for Exercise in Type 1 Diabetes: A Pilot Study. *Diabetes Care*. 43(9):2176-2182, 2020.
- 87.** Hannemann J, Laing A, Glismann K, Skene DJ, Middleton B, Staels B, Marx N, Grant PJ, Federici M, Niebauer J, Böger R. Timed physical exercise does not influence circadian rhythms and glucose tolerance in rotating night shift workers: The EuRhythDia study. *Diab Vasc Dis Res*. 17(5):1479164120950616, 2020.
- 88.** Garruti G, De Fazio M, Capuano P, Martinez G, Rotelli MT, Puglisi F, Palasciano N, Giorgino F. Exercise and apulian hypocaloric diet affect adipokine changes and gastric banding-induced weight loss: A prospective study on severe obese subjects. *Ann Med Surg (Lond)*. 52:10-15, 2020.
- 89.** Cannata F, Vadalà G, Russo F, Papalia R, Napoli N, Pozzilli P. Beneficial Effects of Physical Activity in Diabetic Patients. *J Funct Morphol Kinesiol*. 5(3):70, 2020.