

1. Wiseman M.J., Mangili R., Alberetto M., Keen H., Viberti G.C. Glomerular response mechanisms to glycaemic changes in insulin-dependent diabetics. *Kidney Int* 31:1012, 1987.
2. Pozzilli P, Visalli N, Corcos A, Andreani D. Why do so few diabetic patients attend dialysis centers in central Italy? *J Diabet Complications*. 1(1):4-6, 1987.
3. . Mangili R, Bending J.J., Scott G.S., Li L.K., Gupta A, Viberti G.C. Increased sodium/lithium countertransport activity in red cells of patients with insulin-dependent diabetes and nephropathy. *N Engl J Med* 318:146, 1988.
4. Mangili R., Viberti G.C, Vergani D. Antibodies to human albumin epitopes in Type I (insulin-dependent) diabetes mellitus. *Diabetologia* 31:639, 1988.
5. Morano S, Cancelli A, Bacci S, Frontoni S, Napoli A, Fallucca F, Gambardella S, Di Mario U. The selective elimination of anionic immunoglobulins as a parameter of kidney damage in diabetes and diabetic pregnancy. *J Diabet Complications*.2:2-4, 1988.
6. Giampietro O, Miccoli R, Anichini R, Penno G, Navalesi R. Short-term vs. overnight urine collection for screening of urinary albumin excretion rate. *Diabetes Care*. 11: 509-10, 1988.
7. Giampietro O, Miccoli R, Clerico A, Penno G, Cruschelli L, Palmieri L, Di Palma L, Gregori G, Anichini R, Bertoli S, Navalesi R. Early diagnosis of diabetic nephropathy. Significance and clinical usefulness of albuminuria and enzymuria. *Minerva Endocrinol*. 13: 187-201,1988.
8. Tilton RG, Pugliese G, Chang K, Kilo C, Williamson JR. Effects of nephrectomy and high protein diets on glomerular hemodynamics and urinary protein excretion in diabetic rats. *J Diab Compl*. 2:30-33, 1988.
9. Tomaselli L., Trischitta V., Vinci C., Frittitta L, Squatrito S., Vigneri R. Evaluation of albumin excretion rate in overnight versus 24-h urine. *Diabetes Care* 12:585-587, 1989.
10. Miccoli R, Giampietro O, Penno G, Odello G, Anichini R, Bertolotto A, Cruschelli L, Bertoli S, Navalesi R. 'Microalbuminuria' in type I (insulin-dependent) diabetic patients with and without retinopathy. *Acta Diabetol Lat*. 26: 163-70, 1989.
11. Giampietro O, Clerico A, Cruschelli L, Penno G, Navalesi R. Microalbuminuria in diabetes mellitus: more on urine storage and accuracy of colorimetric assays. *Clin Chem*. 35: 1560-1562, 1989.
12. Di Mario U, Morano S, Cancelli A, Bacci S, Frontoni S, Pietravallo P, Gambardella S, Andreani D. New parameters to monitor the progression of diabetic nephropathy. *Am J Kidney Dis*.13:45-8, 1989.
13. Gambardella S, Morano S, Cancelli A, Pietravallo P, Frontoni S, Bacci S, Di Mario U, Andreani D. Urinary IgG4: an additional parameter in characterizing patients with incipient diabetic nephropathy. *Diabetes Res*.10:153-7, 1989.

14. Gambardella S, Frontoni S, Grazia Felici M, Spallone V, Gargiulo P, Morano S, Menzinger G. Efficacy of antihypertensive treatment with indapamide in patients with noninsulin-dependent diabetes and persistent microalbuminuria. *Am J Cardiol.*65:46H-50H, 1990.
15. Odetti PR, Borgoglio A, De Pascale A, Rolandi R, Adezati L. Prevention of diabetes-increased aging effect on rat collagen-linked fluorescence by aminoguanidine and rutin. *Diabetes.* 39(7):796-801, 1990.
16. Di Mario U, Cancelli A, Pietravallo P, Altamore G, Mariani G, De Rossi MG, Bernardini G, Pasquale A, Borgia MC, Frontoni S, et al. Anionic versus cationic immunoglobulin clearance in normal subjects: a novel approach to the evaluation of charge permselectivity. *Nephron.*55:400-7, 1990.
17. Clerico A, Giammattei C, Cecchini L, Lucchetti A, Cruschelli L, Penno G, Gregori G, Giampietro O. Exercise-induced proteinuria in well-trained athletes. *Clin Chem.*; 36: 562-4, 1990.
18. Cruschelli L, Clerico A, Penno G, Navalesi R, Giampietro O. Colorimetric versus radioimmunological measurement of glycated and non-glycated serum albumin after affinity chromatography. *Acta Diabetol Lat.*; 27: 349-56, 1990.
19. Giampietro O, Cruschelli L, Penno G, Navalesi R, Clerico A. More on effects of storage time and temperature on measurement of small concentrations of albumin in urine. *Clin Chem.* 37: 591-5, 1991.
20. Clerico A, Penno G, Giampietro O. Albuminuria in the diabetic patient: clinical relevance and methodological features. *Minerva Endocrinol.* 16: 157-161, 1991.
21. Monnier V.M, Sell D.R., Miyata S., Nagaraj R.H., Odetti P., Lapolla. A Advanced Maillard reaction products as markers for tissue damage in diabetes and uremia: relevance to diabetic nephropathy. *Acta Diabetol.* 29, 130-135, 1992.
22. Ido Y, Tilton RG, Chang K, Pugliese G, Williamson JR. Rapid measurement of glomerular filtration rate in small animals. *Kidney Int.* 41:435-439, 1992.
23. Di Mario U, Bacci S, Morano S, Pugliese G, Pietravallo P, Andreani D, Morabito S, Simonetti BM, Pierucci A. Selective decrement of anionic immunoglobulin clearance after induced renal hemodynamic changes. *Am J Physiol.* 262:F381-F388, 1992.
24. Pricci F, Pugliese G, Pugliese F, Menè P, Andreani D, Galli G, Casini A, Rotella CM, Di Mario U. Mechanisms of glucose-induced extracellular matrix overproduction by cultured vascular cells. *Acta Diabetol.* 29:264-267, 1992.
25. Mangili R., Sironi S., Rankel G., Makarovic M., Zerbini G., Del Maschio A., Pozza G. Magnetic resonance imaging of the kidney in Type 1 (insulin-dependent) diabetes mellitus.

Diabetologia 35:1002-1008, 1992.

26. Mangili R., Deferrari G, Di Mario U., Giampietro O., Navalesi R., Nosadini R., Rigamonti G., Crepaldi G, on behalf of the Italian Microalbuminuria Study Group. Prevalence of hypertension and microalbuminuria in adult Type 1 (insulin-dependent) diabetic patients without renal failure in Italy. I. Validation of screening techniques to detect microalbuminuria. *Acta Diabetol* 29:156-166, 1992.
27. Giampietro O, Penno G, Clerico A, Cruschelli L, Lucchetti A, Nannipieri M, Cecere M, Rizzo L, Navalesi R. Which method for quantifying "microalbuminuria" in diabetics? Comparison of several immunological methods (immunoturbidimetric assay, immunonephelometric assay, radioimmunoassay and two semiquantitative tests) for measurement of albumin in urine. *Acta Diabetol.* 28: 239-45, 1992.
28. Penno G, Clerico A, Cruschelli L, Rigamonti G, Crepaldi G, Navalesi R, Giampietro O. External quality control for radioimmunoassay of urinary albumin in diabetic subjects: a one-year experience. *Clin Chem.* 38: 2333-5, 1992.
29. Penno G, Rizzo L, Cecere M, Nannipieri M, Lucchetti A, Rapuano A, Giovannitti G, Miccoli R, Navalesi R, Giampietro O. Microalbuminuria is linked with abnormalities favoring atherosclerosis in type 2 diabetics. *Contrib Nephrol.* 101: 127-34, 1993.
30. Giampietro O, Penno G, Clerico A, Cruschelli L, Cecere M. How and how long to store urine samples before albumin radioimmunoassay: a practical response. *Clin Chem.* 39: 533-6, 1993.
31. Mangili R., Zerbini G., Barlassina C., Cusi D, Pozza G. Sodium-lithium countertransport and triglycerides in Type 1 (insulin-dependent) diabetes mellitus. *Kidney Int* 44:127-133, 1993.
32. Menè P, Pugliese G, Pricci F, Di Mario U, Cinotti GA, Pugliese F. High glucose inhibits cytosolic calcium signalling in cultured rat mesangial cells. *Kidney Int.* 43:585-589, 1993.
33. Mangili R., Deferrari G, Di Mario U., Giampietro O, Navalesi R., Nosadini R, Rigamonti G., Spezia R., Crepaldi, G on behalf of the Italian Microalbuminuria Study Group. Arterial hypertension and microalbuminuria in type 1 diabetes. The Italian Microalbuminuria Study. *Diabetologia* 37:1015-1024, 1994.
34. Pugliese G, Pricci F, Pugliese F, Menè P, Lenti L, Andreani D, Galli G, Casini A, Bianchi S, Rotella CM, Di Mario U. Mechanisms of glucose-enhanced extracellular matrix accumulation in rat glomerular mesangial cells. *Diabetes.* 43:478-490, 1994.
35. Bruno G, Pagano G and the Piedmont Study Group for Diabetes Epidemiology. Low prevalence of microalbuminuria in young Italian insulin-dependent diabetic patients with short duration of disease: a population-based study. *Diabetic Med* 13:889-93, 1996.
36. Pricci F, Pugliese G, Menè P, Romeo G, Romano G, Galli G, Casini A, Rotella CM, Di Mario

- U, Pugliese F. Regulatory role of eicosanoids in extracellular matrix overproduction induced by prolonged exposure to high glucose in cultured rat mesangial cells. *Diabetologia*. 39:1055-1062, 1996.
37. Pugliese G, Pricci F, Locuratolo N, Romeo G, Romano G, Giannini S, Cresci B, Galli G, Rotella CM, Di Mario U. Increased activity of the insulin-like growth factor system in mesangial cells cultured in high glucose conditions: relation to glucose-enhanced extracellular matrix production. *Diabetologia*. 39:775-784, 1996.
 38. Pricci F, Pugliese G, Romano G, Romeo G, Locuratolo N, Menè P, Pugliese F, Galli G, Casini A, Rotella CM, Di Mario U. Insulin-like growth factor (IGF)-I and IGF-II stimulate extracellular matrix production in human mesangial cells. Comparison with transforming growth factor- β (TGF- β). *Endocrinology*. 137:879-885, 1996.
 39. Bruno G, Cavallo-Perin P, Bargero G, Borra M, Calvi V, D'Errico N, Deambrogio P, Pagano G. Prevalence and risk factors for micro- and macroalbuminuria in an Italian population-based cohort of non-insulin-dependent diabetic subjects. *Diabetes Care*. 19:43-47, 1996.
 40. Zerbini G, Mangili R., Gabellini D., Pozza G. Modes of operation of an electroneutral Na^+/Li^+ countertransport in human skin fibroblasts. *Am J Phys*, 272: C1373-C1379, 1997.
 41. Zerbini G, Bazzigaluppi E, Mangili R, Bosi E, Bonifacio E. Prevalence of GAD65 and IA-2 antibodies in IDDM patients affected by diabetic nephropathy. *Diabetologia* 40:485, 1997.
 42. Faronato PP, Maioli M, Tonolo G, Brocco E, Noventa F, Piarulli F, Abaterusso C, Modena F, de Bigontina G, Velussi M, Inchiostro S, Santeusano F, Bueti A, Nosadini R. Clustering of albumin excretion rate abnormalities in Caucasian patients with NIDDM. The Italian NIDDM Nephropathy Study Group. *Diabetologia*. 40(7):816-23, 1997.
 43. Morrone, L. F., S. Di Paolo, F. Logoluso, A. Schena, G. Stallone, F. Giorgino, F. P. Schena. Interference of angiotensin-converting enzyme inhibitors on erythropoiesis in kidney transplant recipients. Role of growth factors and cytokines. *Transplantation* 64:913-918, 1997.
 44. Pugliese G, Pricci F, Romeo G, Menè P, Pugliese F, Giannini S, Cresci B, Galli G, Rotella CM, Vlassara H, Di Mario U. Up-regulation of mesangial growth factor and extracellular matrix synthesis by advanced glycation endproducts (AGEs) via a receptor-mediated mechanism. *Diabetes*. 46:1881-1887, 1997.
 45. Pugliese G, Pricci F, Pesce CM, Romeo G, Lenti E, Vetri M, Caltabiano V, Purrello F, Di Mario U. Early, but not advanced, experimental diabetic glomerulopathy is reversed by pancreatic islet transplants. Correlation with glomerular extracellular matrix mRNA levels. *Diabetes*. 46:1198-1206, 1997.
 46. Menè P, Pugliese G, Pricci F, Di Mario U, Cinotti GA, Pugliese F. High glucose inhibits

capacitative Ca^{2+} influx in cultured rat mesangial cells by a protein kinase C-dependent mechanism. *Diabetologia*. 40:521-527, 1997.

47. Pugliese G, Pricci F, Menè P, Romeo G, Nofroni I, Giannini S, Cresci B, Galli G, Rotella CM, Di Mario U, Pugliese F. High glucose level unmasks a genetic predisposition to enhanced extracellular matrix production in mesangial cells from rats of the Milan normotensive strain. *J Am Soc Nephrol*. 8:406-414, 1997.
48. Crepaldi G, Carta Q., Deferrari G, Mangili R, Navalesi R., Santeusano F., Spalluto A., Vanasia A., Villa G. Nosadini R, on behalf of the Italian Microalbuminuria Study Group. Effects of lisinopril and nifedipine on the progression to overt proteinuria in insulin-dependent diabetic patients with incipient nephropathy and normal blood pressure. *Diabetes Care* 21:104-110, 1998.
49. Traverso N, Menini S, Cosso L, Odetti P, Pronzato MA, Marinari UM. Immunological evidence for increased oxidative stress in diabetic rats. *Diabetologia*. 41:265-70, 1998.
50. Zerbini G, Mangili R, Pozza G. Independence of dimethylamiloride-sensitive Li^+ efflux pathways and Na-Li countertransport in human erythrocytes. *Biochim Biophys Acta* 1371:129-133, 1998.
51. Zerbini G, Maestroni A, Mangili R, Pozza G. Amiloride-Insensitive $\text{Na}^+\text{-H}^+$ Exchange: a candidate mediator of erythrocyte $\text{Na}^+\text{-Li}^+$ Countertransport *J Am Soc Nephrol* 9:2203-11, 1998.
52. Pucci L, Bandinelli S, Penno G, Nannipieri M, Rizzo L, Navalesi R. Iohexol plasma clearance in determining glomerular filtration rate in diabetic patients. *Ren Fail.*, 20: 277-284 1998.
53. Penno G, Chaturvedi N, Talmud PJ, Cotroneo P, Manto A, Nannipieri M, Luong LA, Fuller JH. Effect of angiotensin-converting enzyme (ACE) gene polymorphism on progression of renal disease and the influence of ACE inhibition in IDDM patients: findings from the EUCLID Randomized Controlled Trial. EURODIAB Controlled Trial of Lisinopril in IDDM. *Diabetes*. 47: 1507-11, 1998.
54. Nosadini R, Brocco E, Saller A, Cernigoi AM, Abaterusso C, Bortoloso E, Dalla Vestra M, Maioli M, Piarulli F, Fioretto P Renal function in noninsulin-dependent diabetes mellitus patients treated with angiotensin-converting enzyme inhibitors and calcium channel blockers.. *J Hypertens Suppl*. 16(4):S27-32, 1998.
55. Menè P, Pecci G, Cinotti GA, Pugliese G, Pricci F, Pugliese F. Eicosanoid synthesis in peripheral blood monocytes: a marker of disease activity in lupus nephritis. *Am J Kid Dis*. 32:778-784, 1998.
56. De Cosmo S., Margaglione M., Tassi V., Garrubba M., Thomas S, Olivetti C., Piras GP,

- Trevisan R, Vedovato M., Cavallo Perin P., Bacci S., Colaizzo D., Cisternino C., Zucaro L., Di Minno G., Trischitta V., Viberti GC. ACE, PAI-1, Decorin and Werner Helicase genes are not associated with the development of renal disease in European patients with type 1 diabetes. *Diabetes Metab Res Rev* 15: 247-253, 1999.
57. Nannipieri M, Penno G., Pucci L, Colhoun H., Motti C., Bertacca A., Rizzo L., De Giorgio L., Zerbini G., Mangili R., Navalesi. R. Pronatriodilatin gene polymorphisms, microvascular permeability, and diabetic nephropathy in type 1 diabetes mellitus. *J Am Soc Nephrol.* 10(7):1530-41, 1999.
 58. Zerbini G, Podestà F, Mangili R, Ghelardi R, Meregalli G, Gabellini D, Pozza G. Tetraploid cells in diabetic nephropathy. *Diabetologia* 42:118, 1999.
 59. Pugliese G, Pricci F, Romeo G, Leto G, Amadio L, Iacobini C, Di Mario U. Autocrine and paracrine mechanisms in the early stages of diabetic nephropathy. *J Endocrinol Invest.* 22:708-735, 1999.
 60. Bacci S, De Cosmo S, Garrubba M., Placentino G., Liuzzi A., Barbano F., Di Giorgio A, Trischitta V., Viberti GC. Role of insulin-like growth factor (IGF)-1 in the modulation of renal haemodynamics in type 1 diabetic patients. *Diabetologia*, 43: 922-926, 2000.
 61. Pricci F, Leto G, Amadio L, Iacobini C, Romeo G, Cordone S, Gradini R, Barsotti P, Liu F-T, Di Mario U, Pugliese G. Role of galectin-3 as a receptor for advanced glycosylation endproducts. *Kidney Int.* 58:S31-S39, 2000.
 62. Pugliese G, Pricci F, Leto G, Amadio L, Iacobini C, Romeo G, Lenti L, Sale P, Gradini R, Liu F-T, Di Mario U. The diabetic milieu modulates the AGE-receptor complex in the mesangium by inducing or up-regulating galectin-3 expression. *Diabetes*.49:1249-1257, 2000.
 63. Nosadini R, Velussi M, Brocco E, Bruseghin M, Abaterusso C, Saller A, Dalla Vestra M, Carraro A, Bortoloso E, Sambataro M, Barzon I, Frigato F, Muollo B, Chiesura-Corona M, Pacini G, Baggio B, Piarulli F, Sfriso A, Fioretto P. Course of renal function in type 2 diabetic patients with abnormalities of albumin excretion rate. *Diabetes* 49(3):476-84, 2000.
 64. Di Mario U, Pugliese G. 15th Golgi lecture: from hyperglycaemia to the dysregulation of vascular remodelling in diabetes. *Diabetologia.* 44:674-692, 2001.
 65. Chaturvedi N., Bandinelli S., Mangili R., Penno G., Rottiers R.E., Fuller J.H. Microalbuminuria in type 1 diabetes: rates, risk factors and glycemic threshold. *Kidney Int.* 60(1):219-27, 2001.
 66. Brocco E, Velussi M, Cernigoi AM, Abaterusso C, Bruseghin M, Carraro A, Sambataro M, Piarulli F, Sfriso A, Nosadini R. Evidence of a threshold value of glycated hemoglobin to improve the course of renal function in type 2 diabetes with typical diabetic glomerulopathy. *J Nephrol.* 14(6):461-71, 2001.

67. Pucci L, Bandinelli S, Pilo M, Nannipieri M, Navalesi R, Penno G. Iohexol as a marker of glomerular filtration rate in patients with diabetes: comparison of multiple and simplified sampling protocols. *Diabet Med.* 18: 116-20, 2001.
68. Pugliese G, Pricci F, Iacobini C, Leto G, Amadio L, Barsotti P, Frigeri L, Hsu DK, Vlassara H, Liu F-T, Di Mario U. Accelerated diabetic glomerulopathy in galectin-3/AGE-receptor-3 knockout mice. *FASEB J.* 15:2471-2479, 2001.
69. Pesce C, Menini S, Pricci F, Favre A, Leto G, DiMario U, Pugliese G. Glomerular cell replication and cell loss through apoptosis in experimental diabetes mellitus. *Nephron.* 90:484-8, 2002.
70. Soedamah-Muthu S.S., Colhoun H.M., Abrahamian H., Chan N.N., Mangili R., Reboli G.P, Fuller J.H.; EURODIAB Prospective Complications Study Group. Trends in hypertension management in Type I diabetes across Europe, 1989/1990 - 1997/1999. *Diabetologia.* 45:1362-71, 2002.
71. Zerbini G, Maestroni A, Breviario D., Mangili R., Casari G. Alternative splicing of NHE-1 mediates Na-Li countertransport and associates with activity rate. *Diabetes.* 52:1511-8, 2003.
72. Iacobini C, Amadio L, Oddi G, Ricci C, Barsotti P, Missori S, Sorcini M, Di Mario U, Pricci F, Pugliese G. Role of galectin-3 in diabetic nephropathy. *J Am Soc Nephrol.* 14:S264–S270, 2003.
73. Bruno G, Merletti F, Biggeri A, Bargero G, Ferrero S, Pagano G, Cavallo Perin P. Progression to overt nephropathy in type 2 diabetes: the Casale Monferrato Study. *Diabetes Care* 26: 2150-55, 2003.
74. Bruno G, Biggeri A, Merletti F, Bargero G, Ferrero S, Pagano G, Cavallo Perin P. Low incidence of end-stage renal disease and chronic renal failure in Type 2 diabetes: a 10-years prospective study. *Diabetes Care.* 26:2353-58, 2003.
75. Odetti P, Pesce C, Traverso N, Menini S, Maineri EP, Cosso L, Valentini S, Patriarca S, Cottalasso D, Marinari UM, Pronzato MA. Comparative trial of N-acetyl-cysteine, taurine, and oxerutin on skin and kidney damage in long-term experimental diabetes. *Diabetes.* 52:499-505, 2003.
76. Ruggerenti P, Fassi A, Parvanova AI, Bruno S, Ilian PI, Brusegan V, Rubis N, Ghepari G, Arnoldi F, Ganeva M, Ene-Iordache B, Gaspari F, Perna A, Bossi A, Trevisan R, Dodesini AR, Remuzzi G for the Bergamo Nephrologic Diabetes Complication Trial (BENEDICT) Investigators. Preventing Microalbuminuria in Type 2 Diabetes. *New Eng J Med* 351:1941-1951, 2004.
77. Giorgino F, Laviola L, Cavallo Perin P, Solnica B, Fuller J, Chaturvedi N. Factors associated

with progression to macroalbuminuria in microalbuminuric Type 1 diabetic patients: the EURODIAB Prospective Complications Study. *Diabetologia*.47:1020-1028, 2004.

78. Traverso N, Menini S, Pesce Maineri E, Patriarca S, Odetti P, Cottalasso D, Marinari UM, Pronzato MA. Malondialdehyde, a lipoperoxidation-derived aldehyde, can bring about secondary oxidative damage to proteins. *J Gerontol A Biol Sci Med Sci*. 59:B890-5, 2004.
79. Iacobini C, Menini S, Oddi G, Ricci C, Amadio L, Pricci F, Olivieri A, Sorcini M, Di Mario U, Pesce C, Pugliese G. Galectin-3/AGE-receptor 3 knockout mice show accelerated AGE-induced glomerular injury. Evidence for a protective role of galectin-3 as an AGE-receptor. *FASEB J*. 18:1773-1775, 2004.
80. Menini S, Ricci C, Iacobini C, Bianchi G, Pugliese G, Pesce C. Glomerular number and size in Milan hypertensive and normotensive rats: their relationship to susceptibility and resistance to hypertension and renal disease. *J Hypertens*. 22:2185-2192, 2004.
81. Pohl M.A, Blumenthal S., Cordonnier D.J, De Alvaro F., DeFerrari G., Eisner G., Gilbert R.E., Hunsicker L.G., Lopes de Faria J.B, Mangili R., Moore, Jr. J., Reisin E., Ritz E., Schernthaner G., Spitalowitz S, Tindall H., Rodby R.A, Lewis E.J., For the Collaborative Study Group. Independent and additive impact of blood pressure control and angiotensin II receptor blockade on renal outcomes in the irbesartan diabetic nephropathy trial: clinical implications and limitations. *J Am Soc Nephrol*. 16:2170-9, 2005.
82. De Cosmo S., Minenna A., Ludovico O., Mastroianno S, Di Giorgio A, Pirro L, Trischitta V. Increased urinary albumin excretion, insulin-resistance and related cardiovascular risk factors in patients with type 2 diabetes: evidence of a gender-specific association. *Diabetes Care* 25: 910-915, 2005.
83. Iacobini C, Oddi G, Menini S, Amadio L, Ricci C, Barsotti P, C Di Pippo, Sorcini M, Pricci F, Pugliese F, Pugliese G. Development of age-dependent glomerular lesions in galectin-3/AGE-receptor-3 knockout and wild type mice. *Am J Physiol (Renal Physiol)*. 289:F611-F621, 2005.
84. Pugliese G, Pricci F, Barsotti P, Iacobini C, Ricci C, Oddi G, Romeo G, Leto G, Marano G, Sorcini M, Sabbatini M, Fuiano G, Di Mario U, Pugliese F. Development of diabetic nephropathy in the Milan Normotensive Strain, but not in the Milan Hypertensive Strain. Possible permissive role of hemodynamics. *Kidney Int*. 67:1440-1452, 2005.
85. Solini A, Iacobini C, Ricci C, Chiozzi P, Amadio L, Pricci F, Di Mario U, Di Virgilio F, Pugliese G. Purinergic modulation of mesangial extracellular matrix production: role in diabetic and other glomerular diseases. *Kidney Int*; 67:875-885, 2005.
86. Furfaro AL, Menini S, Patriarca S, Pesce C, Odetti P, Cottalasso D, Marinari UM, Pronzato MA, Traverso N. HNE-dependent molecular damage in diabetic nephropathy and its possible

prevention by N-acetyl-cysteine and oxerutin. *Biofactors*. 24:291-8, 2005.

87. Ji H, Menini S, Mok K, Zheng W, Pesce C, Kim J, Mulroney S, Sandberg K. Gonadal steroid regulation of renal injury in renal wrap hypertension. *American Journal of Physiology: Am J Physiol Renal Physiol*. 288:F513-20,2005.
88. Ji H, Pesce C, Mok K, Zheng W, Menini S, Haiwood JR, Mulroney S, Sandberg K. Sex differences in renal injury and nitric oxide production in renal wrap hypertension. *American Journal of Physiology: Am J Physiol Heart Circ Physiol*. 288:H43-7, 2005.
89. Nosadini R, Velussi M, Brocco E, Abaterusso C, Piarulli F, Morgia G, Satta A, Faedda R, Abhyankar A, Luthman H, Tonolo G Altered transcapillary escape of albumin and microalbuminuria reflects two different pathogenetic mechanisms.. *Diabetes*. 54(1):228-33, 2005.
90. De Cosmo S, Lamacchia O, Rauseo A, Viti R, Gesualdo L, Pilotti A, Trischitta V, Cignarelli M. Cigarette smoking is associated with low glomerular filtration rate in male patients with type 2 diabetes. *Diabetes Care* 29(11):2467-70, 2006.
91. Odetti P, Traverso N, Monacelli F, Menini S, Vazzana J, Tasso B, Pronzato MA, Robaudo C, Deferrari G. Vitamin E-coated filter decreases levels of free 4-hydroxyl-2-nonenal during haemodialysis sessions. *Free Radic Res*. 40:207-12, 2006.
92. Menini S, Amadio L, Oddi G, Ricci C, Pesce C, Pugliese F, Giorgio M, Migliaccio E, Pelicci PG, Iacobini C, Pugliese G. Deletion of p66^{Shc} longevity gene protects against experimental diabetic glomerulopathy by preventing diabetes-induced oxidative stress. *Diabetes*. 55:1642-1650, 2006.
93. Ricci C, Iacobini C, Oddi G, Amadio L, Menini S, Rastaldi MP, Frasheri A, Pricci F, Pugliese F, Pugliese G. Role of TGF- β /GLUT1 axis in susceptibility vs. resistance to diabetic glomerulopathy in the Milan rat model. *Nephrol Dial Transplant*. 21:1514-1524, 2006.
94. Nosadini R, Velussi M, Brocco E, Abaterusso C, Carraro A, Piarulli F, Morgia G, Satta A, Faedda R, Abhyankar A, Luthman H, Tonolo G Increased renal arterial resistance predicts the course of renal function in type 2 diabetes with microalbuminuria.. *Diabetes*. 55(1):234-9, 2006.
95. Menini S, Pesce C. Inside Laboratory Investigation: Glomerular cell apoptosis in diabetes. *Lab Invest.*; 86: 535-37, 2006.
96. Biancofiore G, Pucci L, Cerutti E, Penno G, Pardini E, Esposito M, Bindi L, Pelati E, Romanelli A, Triscornia S, Salvadorini MP, Stratta C, Lanfranco G, Pellegrini G, Del Prato S, Salizzoni M, Mosca F, Filipponi F. Cystatin C as a marker of renal function immediately after liver transplantation. *Liver Transpl*. 12: 285-291, 2006.
97. De Cosmo S., Trevisan R., Minenna A, Vedovato M, Viti R, Santini SA, Dodesini AR, Fioretto

- P, Trischitta V. Insulin Resistance and Cluster of Abnormalities Related to the Metabolic Syndrome are Associated with Reduced Glomerular Filtration Rate in Patients with Type 2 Diabetes. *Diabetes Care* 29: 432-434, 2006.
98. Bruno G, Merletti F, Barger G, Novelli G, Melis D, Soddu A, Perotto M, Pagano G, Cavallo-Perin P. Estimated glomerular filtration rate, albuminuria and mortality in type 2 diabetes: the Casale Monferrato Study. *Diabetologia* 50:941-8, 2007.
99. Ji H, Zheng W, Menini S, Pesce C, Kim J, Wu X, Mulroney SE and Sandberg K. Female protection in progressive renal disease is associated with estradiol attenuation of superoxide production. *Gend Med.* 4:56-71, 2007.
100. Pucci L, Triscornia S, Lucchesi D, Fotino C, Pellegrini G, Pardini E, Miccoli R, Del Prato S, Penno G. Cystatin C and estimates of renal function: searching for a better measure of kidney function in diabetic patients. *Clin Chem.* 53: 480-488, 2007.
101. Maione A, Nicolucci A, Craig JC, Tognoni G, Moschetta A, Palasciano G, Pugliese G, Procaccini DA, Gesualdo L, Pellegrini F, GFM, for the LIRICO Study Group. Angiotensin converting enzyme inhibitors, angiotensin receptor blockers and their combination in patients with micro-macroalbuminuria and one or more cardiovascular risk factors. Protocol of the Long-term Impact of RAS Inhibition on Cardiorenal Outcomes randomized trial (LIRICO). *J Nephrol.* 20:646-655, 2007.
102. Solini A, Santini E, Chimenti D, Chiozzi P, Pratesi F, Cuccato S, Falzoni S, Lupi R, Ferrannini E, Pugliese G, Di Virgilio F. Multiple P2X receptors are involved in the modulation of apoptosis in human mesangial cells: evidence for a key role of P2X4. *Am J Physiol (Renal Physiol).* 292:F1537-F1547, 2007.
103. Lapolla A, Ragazzi E, Andretta B, Fedele D, Tubaro M, Seraglia R, Molin L, Traldi P. Multivariate analysis of matrix-assisted laser desorption/ionization mass spectrometric data related to glycoxidation products of human globins in nephropatic patients. *J Am Soc Mass Spectrom.* 18:1018-1023, 2007.
104. De Cosmo S., Lamacchia O., Gesualdo L., Trischitta V., Cignarelli M. Cigarette Smoking is Associated with Low Glomerular Filtration Rate in Male Patients with Type 2 Diabetes: Response to Saito et al. *Diabetes Care* 30(3):e3, 2007.
105. Davis B, Dei Cas A, Long DA, White KE, Hayward A, Ku CH, Woolf AS, Bilous R, Viberti G, Gnudi L. Podocyte-specific expression of angiopoietin-2 causes proteinuria and apoptosis of glomerular endothelia. *J Am Soc Nephrol.* 18(8):2320-9, 2007.
106. Bianchi C, Penno G, Daniele G, Russo E, Giovannitti MG, Del Prato S, Miccoli R. The metabolic syndrome is related to albuminuria in Type 2 diabetes. *Diabet Med.* 25: 1412-1418,

2008.

107. Lapolla A, Brioschi M, Banfi C, Tremoli E, Bonfante L, Cristoni S, Seraglia R, Traldi P. On the search for glycated lipoprotein ApoA-I in plasma of diabetic and nephropatic patients. *J Mass Spectrom* 43:74-81, 2008.
108. Lapolla A, Brioschi M, Banfi C, Tremoli E, Cosma C, Bonfante L, Cristoni S, Seraglia R, Traldi P. Noenzymatically glycated lipoprotein ApoA-I in plasma of diabetic and nephropatic patients. *Ann NY Acad Sci*, 1126:295-299, 2008.
109. Ji H, Menini S, Zheng W, Pesce C, Wu X, Sandberg K. Role of ACE2 and Ang-(1-7) in 17 β -estradiol regulation of renal pathology in renal wrap hypertension. *Exp Physiol*. 93:648-57, 2008.
110. Cignarelli M., Lamacchia O. Cigarette Smoking and Kidney Dysfunction in Diabetes Mellitus. *Journal of Nephrology* 21(2):180-9, 2008.
111. Lamacchia O, Nicastro V., Camarchio D., Stallone G, Gesualdo L., Cignarelli M.. Waist Circumference Is Strongly Associated with Renal Resistive Index in Normoalbuminuric Patients with Type 2 Diabetes *Am. J Nephrol* 28:54-58, 2008.
112. Ku CH, White KE, Dei Cas A, Hayward A, Webster Z, Bilous R, Marshall S, Viberti G, Gnudi L. Inducible overexpression of sFlt-1 in podocytes ameliorates glomerulopathy in diabetic mice. *Diabetes*. 57(10):2824-33, 2008.
113. Iacobini C, Menini S, Ricci C, Scipioni A, Sansoni V, Mazzitelli G, Cordone S, Pesce C, Pugliese F, Pricci F, Pugliese G. Advanced lipoxidation end-products mediate lipid-induced glomerular injury: role of receptor-mediated mechanisms. *J Pathol*. 218:360-369, 2009.
114. Lapolla A, Seraglia R, Molin L, Williams C. Cosma C, Reitano R, Sechi A, Ragazzi E, Traldi. Low molecular weight proteins in urines from healthy subjects as well as nephropatic and diabetic-nephropatic patients: a MALDI study. *J Mass Spectrom*, 44:419-425, 2009.
115. Cignarelli M, Lamacchia O , Cardinale G, Rauseo A, Mastroianno S , Gesualdo L, De Cosmo S. Metabolic syndrome and albuminuria show an additive effect in modulating glomerular filtration rate in patients with Type 2 Diabetes Mellitus. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* 3 (3): 139-142, 2009.
116. Dessapt C, Baradez MO, Hayward A, Dei Cas A, Thomas SM, Viberti G, Gnudi L. Mechanical forces and TGF β 1 reduce podocyte adhesion through α 3 β 1 integrin downregulation. *Nephrol Dial Transplant*.24(9):2645-55, 2009.
117. Iacobini C, Menini S, Ricci C, Scipioni A, Sansoni V, Mazzitelli G, Cordone S, Pesce C, Pugliese F, Pricci F, Pugliese G. Advanced lipoxidation end-products mediate lipid-induced glomerular injury: role of receptor-mediated mechanisms. *J Pathol*. 218:360-369, 2009.

118. Papale M, Di Paolo S, Magistroni R, Lamacchia O, De Mattia A, Teresa Rocchetti M, Furci L, Pasquali S, De Cosmo S, Cignarelli M, Gesualdo L. Urine proteome analysis may allow non-invasive differential diagnosis of diabetic nephropathy. *Diabetes Care*. 33(11):2409-15, 2010.
119. Lamacchia O, Nicastro V, Camarchio D, Valente U, Grisorio R, Gesualdo L, Cignarelli M. Para- and perirenal fat thickness is an independent predictor of chronic kidney disease, increased renal resistance index and hyperuricaemia in type-2 diabetic patients. *Nephrol Dial Transplant* 26(3):892-8, 2011.
120. Maione A, Navaneethan SD, Graziano G, Mitchell R, Johnson D, Mann JF, Gao P, Craig JC, Tognoni G, Perkovic V, Nicolucci A, De Cosmo S, Sasso A, Lamacchia O, Cignarelli M, Manfreda VM, Gentile G, Strippoli GF. Angiotensin-converting enzyme inhibitors, angiotensin receptor blockers and combined therapy in patients with micro- and macroalbuminuria and other cardiovascular risk factors: a systematic review of randomized controlled trials. *Nephrol Dial Transplant*. 26(9):2827-47, 2011.
121. Pugliese G, Solini A, Fondelli C, Trevisan R, Vedovato M, Nicolucci A, Penno G; Renal Insufficiency And Cardiovascular Events (RIACE) Study Group. Reproducibility of albuminuria in type 2 diabetic subjects. Findings from the Renal Insufficiency And Cardiovascular Events (RIACE) study. *Nephrol Dial Transplant*. 26: 3950-3954, 2011.
122. Ruggenenti P, Lauria G, Iliev IP, Fassi A, Ilieva AP, Rota S, Chiurciu C, Barlovic DP, Sghirlanzoni A, Lombardi R, Penza P, Cavaletti G, Piatti ML, Frigeni B, Filipponi M, Rubis N, Noris G, Motterlini N, Ene-Iordache B, Gaspari F, Perna A, Zaletel J, Bossi A, Dodesini AR, Trevisan R, Remuzzi G; DEMAND Study Investigators. Effects of manidipine and delapril in hypertensive patients with type 2 diabetes mellitus: the delapril and manidipine for nephroprotection in diabetes (DEMAND) randomized clinical trial. *Hypertension*. 58:776-83, 2011.
123. Targher G., Chonchol M., Zoppini G., Abaterusso C., Bonora E. Risk of chronic kidney disease in patients with non-alcoholic fatty liver disease: is there a link? *J. Hepatol*. 54: 1020-1029, 2011.
124. Pugliese G., Solini A., Bonora E., Orsi E., Zerbini G., Giorgino F., Cavalot F., Pontiroli A.E., Baroni M.G., Morano S., Nicolucci A., Penno G. The chronic kidney disease epidemiology collaboration (CKD-EPI) equation provides a better definition of cardiovascular burden associated with CKD than the Modification Of Diet In Renal Disease (MDRD) Study formula in subjects with type 2 diabetes. *Atherosclerosis* 218: 194-199, 2011.
125. Penno G., Solini A., Bonora E., Fondelli C., Orsi E., Zerbini G., Trevisan R., Vedovato M., Gruden G., Cavalot F., Cignarelli M., Laviola L., Morano S., Nicolucci A., Pugliese G.; for the

Renal Insufficiency And Cardiovascular Events (RIACE) Study Group. Clinical significance of nonalbuminuric renal impairment in type 2 diabetes. *J. Hypertens.* 29: 1802-1809, 2011.

126. Lapolla A, Molin L, Seraglia R, Sechi A, Cosma C, Bonfante L, Chilelli NC, Ragazzi E, Traldi P. Urinary peptides as a diagnostic tool for renal failure detected by matrix-assisted laser desorption/ionisation mass spectrometry: an evaluation of their clinical significance. *Eur J Mass Spectrom.* 17(3):245-53, 2011.
127. Menzaghi C., De Cosmo S., Copetti M., Salvemini L., De Bonis C., Mangiacotti D., Fini G, Pellegrini F., Trischitta V. Relationship between ADIPOQ gene, circulating HMW adiponectin and albuminuria in individuals with normal kidney function: evidence from a family-based study. *Diabetologia*, 54: 812-8, 2011.
128. Lapolla A, Porcu S, Traldi P. Mass spectrometry for diabetic nephropathy monitoring: new effective tools for physicians. *ISRN Endocrinol.* 2012:768159, 2012.
129. Menzaghi C., Salvemini L., Fini G., Thompson R., Mangiacotti D., Di Paola R., Morini E., Giorelli M., De Bonis C., De Cosmo S., Doria A., Trischitta V. Serum resistin and kidney function: a family based study in non diabetic, untreated individuals. *PLoS ONE* 7:e38414, 2012.
130. Pugliese G, Mazzitelli G, Bazuro A, Pugliese L, Salvi L. Mechanisms and clinical significance of reduced glomerular filtration rate in diabetes. *G Ital Diabetol Metab.* 32:10-19, 2012.
131. Molin L, Seraglia R, Lapolla A, Ragazzi E, Gonzalez J, Vlahou A, Schanstra JP, Albalat A, Dakna M, Siwy J, Jankowski J, Bitsika V, Mischak H, Zürbig P, Traldi P. A comparison between MALDI-MS and CE-MS data for biomarker assessment in chronic kidney diseases. *J Proteomics*, 2012.
132. Sasso FC, Chiodini P, Carbonara O, De Nicola L, Conte G, Salvatore T, Nasti R, Marfella R, Gallo C, Signoriello S, Torella R, Minutolo R; on behalf of the NID-2 (Nephropathy In Type 2 Diabetes) Study Group. High cardiovascular risk in patients with Type 2 diabetic nephropathy: the predictive role of albuminuria and glomerular filtration rate. The NID-2 Prospective Cohort Study. *Nephrol Dial Transplant.* 27(6):2269-2274, 2012.
133. Targher G., Pichiri I., Zoppini G., Trombetta M., Bonora E. Increased prevalence of chronic kidney disease in patients with Type 1 diabetes and non-alcoholic fatty liver. *Diabet. Med.* 29: 220-226, 2012.
134. Solini A., Penno G., Bonora E., Fondelli C., Orsi E., Arosio M., Trevisan R., Vedovato M., Cignarelli M., Andreatozzi F., Nicolucci A., Pugliese G.; for the Renal Insufficiency And Cardiovascular Events (RIACE) Study Group. Diverging association of reduced glomerular

filtration rate and albuminuria with coronary and noncoronary events in patients with type 2 diabetes: The Renal Insufficiency and Cardiovascular Events (RIACE) Italian Multicentre Study. *Diabetes Care* 35: 143-149, 2012.

135. Doria A, Niewczasz MA, Fiorina P. Can existing drugs approved for other indications retard renal function decline in patients with type 1 diabetes and nephropathy? *Semin Nephrol.* 32(5):437-44, 2012.
136. Zoppini G., Targher G., Chonchol M., Ortalda V., Negri C., Stoico V., Bonora E. Predictors of estimated GFR decline in patients with type 2 diabetes and preserved kidney function. *Clin. J. Am. Soc. Nephrol.* 7: 401-408, 2012.
137. Ruggenenti P, Porrini E, Gaspari F, Cannata AN, Carrara F, Cella C, Ferrari S, Stucchi N, Parvanova Ilieva A, Iliev IP, Dodesini AR, Trevisan R, Bossi AC, Zaletel J, G Remuzzi for the GFR Study Investigators. Glomerular Hyperfiltration and Renal Disease Progression in Type 2 Diabetes. *Diabetes Care* 35:2061–2068, 2012.
138. Ruggenenti P, Porrini E, Motterlini N, Perna A, Parvanova Ilieva A, Iliev IP, Dodesini AR, Trevisan R, Bossi AC, Sampietro G, Capitoni E, Gaspari F, Rubis N, Ene-Iordache B, and Remuzzi G, for the BENEDICT Study Investigators. Measurable Urinary Albumin Predicts Cardiovascular Risk among Normoalbuminuric Patients with Type 2 Diabetes. *J Am Soc Nephrol*, 23:1717-24, 2012.
139. De Cosmo S, Menzaghi C, Prudente S, Trischitta V. Role of insulin resistance in kidney dysfunction: insights into the mechanism and epidemiological evidence. *Nephrol Dial Transplant.* 1:29-36, 2013.
140. Ruggenenti P, Porrini E, Bossi AC, Parvanova A, Perna A, Gaspari F, Petrov, Trevisan R, Zaletel J, Remuzzi G. The GFR and GFR decline cannot be accurately estimated in type 2 diabetics. *Kidney Int* 84: 164-173, 2013.
141. Rurali E, Noris M, Chianca A, Donadelli R, Banterla F, Galbusera M, Gherardi G, Gastoldi S, Parvanova Ilieva A, Petrov Iliev I, Bossi AC, Haeffliger C, Trevisan R, Remuzzi G, Ruggenenti P, for the BENEDICT Study Group. ADAMTS13 predicts renal and cardiovascular events in type 2 diabetics and response to therapy. *Diabetes* 62 (10): 3599-609, 2013.
142. Fiorentino L, Cavalera M, Menini S, Marchetti V, Mavilio M, Fabrizi M, Conserva F, Casagrande V, Menghini R, Pontrelli P, Arisi I, D'Onofrio M, Lauro D, Khokha R, Accili D, Pugliese G, Gesualdo L, Lauro R, Federici M. Loss of TIMP3 underlies diabetic nephropathy via FoxO1/STAT1 interplay. *EMBO Mole Med.* 5:441-455, 2013.
143. Fiorentino L, Cavalera M, Menini S, Marchetti V, Mavilio M, Fabrizi M, Conserva F, Casagrande V, Menghini R, Pontrelli P, Arisi I, D'Onofrio M, Lauro D, Khokha R, Accili D,

Pugliese G, Gesualdo L, Lauro R, Federici M. Loss of TIMP3 underlies diabetic nephropathy via FoxO1/STAT1 interplay. *EMBO Mol Med.* 5(3):441-55, 2013.

144. Bonomini M, Di Liberato L., Del Rosso G, Stingone A, Marinangeli G, Consoli A, Bertoli S, De Vecchi A, Bosi E, Russo R, Corciulo R, Gesualdo L, Giorgino F, Cerasoli P, Di Castelnuovo A., Monaco M. P, Shockley T, Rossi C, Arduini A. Effect of an l-Carnitine-Containing Peritoneal Dialysate on Insulin Sensitivity in Patients Treated With CAPD: A 4-Month, Prospective, Multicenter Randomized Trial. *American Journal of Kidney Diseases* 62:929-938, 2013.
145. Solini A, Menini S, Rossi C, Ricci C, Santini E, Blasetti Fantauzzi C, Iacobini C, Pugliese G. The purinergic 2X7 receptor participates in renal inflammation and injury induced by high fat diet: possible role of NLRP3 inflammasome activation. *J Pathol.* 231:342-353, 2013.
146. Penno G, Solini A, Bonora E, Fondelli C, Orsi E, Zerbini G, Morano S, Cavalot F, Lamacchia O, Laviola L, Nicolucci A, Pugliese G; for the Renal Insufficiency And Cardiovascular Events (RIACE) Study Group. HbA1c variability as an independent correlate of nephropathy, but not retinopathy, in patients with type 2 diabetes: The Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicenter Study. *Diabetes Care* 36: 2301-2310, 2013.
147. Barutta F, Tricarico M, Corbelli A, Annaratone L, Pinach S, Grimaldi S, Bruno G, Cimino D, Taverna D, Deregibus MC, Rastaldi MP, Perin PC, Gruden G. Urinary Exosomal MicroRNAs in Incipient Diabetic Nephropathy. *PLoS One.* 8:e73798, 2013.
148. Barutta F, Grimaldi S, Franco I, Bellini S, Gambino R, Pinach S, Corbelli A, Bruno G, Rastaldi MP, Aveta T, Hirsch E, Di Marzo V, Gruden G. Deficiency of cannabinoid receptor of type 2 worsens renal functional and structural abnormalities in streptozotocin-induced diabetic mice. *Kidney Int.* 86 (5): 979-90, 2014.
149. Bruno G, Barutta F, Landi A, Ghezzo G, Baldin C, Cavallo Perin P, Gruden G. The predictive role of low-moderate reduction of renal function on cardiovascular mortality of diabetic patients is due to the strong independent effect of NT-proBNP: the population-based Casale Monferrato Study. *Plos One* 9:e114855, 2014.
150. Pugliese G., Solini A., Bonora E., Fondelli C., Orsi E., Nicolucci A., Penno G., RIACE Study Group. Chronic kidney disease in type 2 diabetes: lessons from the Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicentre Study. *Nutr Metab Cardiovasc Dis.* 24: 815-822, 2014.
151. Targher G., Mantovani A., Pichiri I., Mingolla L., Cavalieri V., Mantovani W. Pancheri S., Trombetta M., Zoppini G., Chonchol M., Byrne C.D., Bonora E. Nonalcoholic fatty liver disease is independently associated with an increased incidence of chronic kidney disease in

patients with type 1 diabetes. *Diabetes Care* 37: 1729-1736, 2014.

152. De Cosmo S, Lamacchia O, Pacilli A, Fariello S, Pinnelli S, Fontana A, Di Mauro L, Cignarelli M, Trischitta V. Normoalbuminuric renal impairment and all-cause mortality in type 2 diabetes mellitus.. *Acta Diabetol.* 51(4):687-9, 2014.
153. Pugliese G, Solini A, Bonora E, Orsi E, Zerbini G, Fondelli C, Gruden G, Cavalot F, Lamacchia O, Trevisan R, Vedovato M, Penno G; RIACE Study Group. Distribution of cardiovascular disease and retinopathy in patients with type 2 diabetes according to different classification systems for chronic kidney disease: a cross-sectional analysis of the renal insufficiency and cardiovascular events (RIACE) Italian multicenter study. *Cardiovasc Diabetol.* 13(1):59, 2014.
154. Fiorina P, Vergani A, Bassi R, Niewczas MA, Altintas MM, Pezzolesi MG, D'Addio F, Chin M, Tezza S, Ben Nasr M, Mattinzoli D, Ikehata M, Corradi D, Schumacher V, Buvall L, Yu CC, Chang JM, La Rosa S, Finzi G, Solini A, Vincenti F, Rastaldi MP, Reiser J, Krolewski AS, Mundel PH, Sayegh MH. Role of Podocyte B7-1 in Diabetic Nephropathy. *J Am Soc Nephrol.* 25 (7):1415-29, 2014.
155. Papa G, Iurato MP, Licciardello C, Maiorana R, Finocchiaro C, Pezzino V. Aging, diabetic nephropathy and multiple macrovascular involvement are associated with atrial fibrillation in type 2 diabetes mellitus. *British Journal of Medicine & Medical Research* 4(32):5154-5166, 2014.
156. Savarese G, Dei Cas A, Rosano G, D'Amore C, Musella F, Mosca S, Reiner MF, Marchioli R, Trimarco B, Perrone-Filardi P. Reduction of albumin urinary excretion is associated with reduced cardiovascular events in hypertensive and/or diabetic patients. A meta-regression analysis of 32 randomized trials. *Int J Cardiol.* 172(2):403-10, 2014.
157. Pugliese G. Updating the natural history of diabetic nephropathy. *Acta Diabetol.* 51:905–915, 2014.
158. Barutta F, Bruno G, Grimaldi S, Gruden G. Inflammation in diabetic nephropathy: moving toward clinical biomarkers and targets for treatment. *Endocrine* 48:730-42, 2015.
159. Ortega Moreno L, Lamacchia O, Copetti M, Salvemini L, De Bonis C, De Cosmo S, Cignarelli M, Trischitta V, Menzaghi C. Serum Adiponectin and Glomerular Filtration Rate in Patients with Type 2 Diabetes. *PLoS One.* 10(10):e0140631, 2015.
160. Penno G, Solini A, Zoppini G, Fondelli C, Trevisan R, Vedovato M, Cavalot F, Gruden G, Lamacchia O, Laviola L, Orsi E, Pugliese G; Renal Insufficiency Cardiovascular Events (RIACE) Study Group Independent correlates of urinary albumin excretion within the normoalbuminuric range in patients with type 2 diabetes: The Renal Insufficiency And

Cardiovascular Events (RIACE) Italian Multicentre Study. *Acta Diabetol.* 52(5):971-81, 2015.

161. Piccoli GB, Grassi G, Cabiddu G, Nazha M, Roggero S, Capizzi I, De Pascale A, Priola AM, Di Vico C, Maxia S, Loi V, Asunis AM, Pani A, Veltri A. Diabetic Kidney Disease: A Syndrome Rather Than a Single Disease. *Rev Diabet Stud* 12(1-2):87-109, 2015.
162. Tesauro M, Nisticò S, Noce A, Tarantino A, Marrone G, Costa A, Rovella V, Di Cola G, Campia U, Lauro D, Cardillo C, Di Daniele N. The possible role of glutathione-S-transferase activity in diabetic nephropathy. *Int J Immunopathol Pharmacol.* 28(1):129-33, 2015.
163. Penno G, Solini A, Zoppini G, Fondelli C, Trevisan R, Vedovato M, Cavalot F, Gruden G, Lamacchia O, Laviola L, Orsi E, Pugliese G; Renal Insufficiency Cardiovascular Events (RIACE) Study Group. Independent correlates of urinary albumin excretion within the normoalbuminuric range in patients with type 2 diabetes: The Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicentre Study. *Acta Diabetol.* 52(5):971-81, 2015.
164. Chilelli NC, Cremasco D, Cosma C, Ragazzi E, Francini Pesenti F, Bonfante L, Lapolla A. Effectiveness of a diet with low advanced glycation end products, in improving glycooxidation and lipid peroxidation: a long-term investigation in patients with chronic renal failure. *Endocrine.* 54(2):552-555, 2016.
165. Bonomini M, Pandolfi A. Chemerin in renal dysfunction and cardiovascular disease. *Vascul Pharmacol.* 77:28-34, 2016.
166. Di Pietro N, Giardinelli A, Sirolli V, Riganti C, Di Tomo P, Gazzano E, Di Silvestre S, Panknin C, Cortese-Krott MM, Csonka C, Kelm M, Ferdinandy P, Bonomini M, Pandolfi A. Nitric oxide synthetic pathway and cGMP levels are altered in red blood cells from end-stage renal disease patients. *Mol Cell Biochem.* 417(1-2):155-67, 2016.
167. Sasso FC, Lascar N, Ascione A, Carbonara O, De Nicola L, Minutolo R, Salvatore T, Rizzo MR, Cirillo P, Paolisso G, Marfella R; NID-2 study group. Moderate-intensity statin therapy seems ineffective in primary cardiovascular prevention in patients with type 2 diabetes complicated by nephropathy. A multicenter prospective 8 years follow up study. *Cardiovasc Diabetol.* 15(1):147, 2016.
168. Bassi R, Fornoni A, Doria A, Fiorina P. CTLA4-Ig in B7-1-positive diabetic and non-diabetic kidney disease. *Diabetologia.* 59(1):21-9, 2016.
169. Solini A, Manca ML, Penno G, Pugliese G, Cobb JE, Ferrannini E. Prediction of Declining Renal Function and Albuminuria in Patients With Type 2 Diabetes by Metabolomics. *J Clin Endocrinol Metab.* 101(2):696-704, 2016.
170. Barutta F, Grimaldi S, Gambino R, Vemuri K, Makriyannis A, Annaratone L, di Marzo V, Bruno G, Gruden G. Dual therapy targeting the endocannabinoid system prevents experimental

diabetic nephropathy. *Nephrol Dial Transplant*. 32(10):1655-1665, 2017.

171. Pontrelli P, Conserva F, Papale M, Oranger A, Barozzino M, Vocino G, Rocchetti MT, Gigante M, Castellano G, Rossini M, Simone S, Laviola L, Giorgino F, Grandaliano G, Di Paolo S, Gesualdo L. Lysine 63 ubiquitination is involved in the progression of tubular damage in diabetic nephropathy. *FASEB J*. 31(1):308-319, 2017.
172. Menini S, Iacobini C, Pugliese G, Pesce C. Dietary interventions to contrast the onset and progression of diabetic nephropathy: A critical survey of new data. *Crit Rev Food Sci Nutr*. 27:1-10, 2017.
173. Penno G, Russo E, Garofolo M, Daniele G, Lucchesi D, Giusti L, Sancho Bornez V, Bianchi C, Dardano A, Miccoli R, Del Prato S. Normoalbuminuric chronic kidney disease in type 1 diabetes: is it real and is it serious? Reply to Rigalleau V, Blanco L, Alexandre L et al *Diabetologia*. 60(10):2123-2125, 2017 (letter)
174. Penno G, Russo E, Garofolo M, Daniele G, Lucchesi D, Giusti L, Sancho Bornez V, Bianchi C, Dardano A, Miccoli R, Del Prato S. Evidence for two distinct phenotypes of chronic kidney disease in individuals with type 1 diabetes mellitus. *Diabetologia*. 60(6):1102-1113, 2017.
175. Fiorentino TV, Marini MA, Succurro E, Sciacqua A, Andreozzi F, Perticone F, Sesti G. Elevated hemoglobin glycation index identify non-diabetic individuals at increased risk of kidney dysfunction. *Oncotarget*. 8(45):79576-79586, 2017.
176. Rossi L, Nicoletti MC, Carmosino M, Mastrofrancesco L, Di Franco A, Indrio F, Lella R, Laviola L, Giorgino F, Svelto M, Gesualdo L, Procino G. Urinary Excretion of Kidney Aquaporins as Possible Diagnostic Biomarker of Diabetic Nephropathy. *J Diabetes Res*. . 2017:4360357, 2017.
177. Ruospo M, Saglimbene VM, Palmer SC, De Cosmo S, Pacilli A, Lamacchia O, Cignarelli M, Fioretto P, Vecchio M, Craig JC, Strippoli GF. Glucose targets for preventing diabetic kidney disease and its progression. *Cochrane Database Syst Rev*. 6:CD010137, 2017.
178. Mantovani A, Zaza G, Byrne CD, Lonardo A, Zoppini G, Bonora E, Targher G. Nonalcoholic Fatty Liver Disease Increases Risk of Incident Chronic Kidney Disease: a Systematic Review and Meta-Analysis. *Metabolism*. 87:1-12 2018.
179. Mantovani A, Rigolon R, Turino T, Pichiri I, Falceri A, Rossi A, Temporelli PL, Bonapace S, Lippi G, Zoppini G, Bonora E, Byrne CD, Targher G. Association between decreasing estimated glomerular filtration rate and risk of cardiac conduction defects in patients with type 2 diabetes. *Diabetes Metab*. 44(6):473-481, 2018.
180. Fadini GP, Solini A, Manca ML, Zatti G, Karamouzis I, Di Benedetto A, Frittitta L,

Avogaro A; DARWIN-T2D Network. Phenotyping normal kidney function in elderly patients with type 2 diabetes: a cross-sectional multicentre study. *Acta Diabetol.* 55(11):1121-1129, 2018.

181. Penno G, Solini A, Orsi E, Bonora E, Fondelli C, Trevisan R, Vedovato M, Cavalot F, Lamacchia O, Scardapane M, Nicolucci A, Pugliese G; Renal Insufficiency And Cardiovascular Events (RIACE) Study Group. Non-albuminuric renal impairment is a strong predictor of mortality in individuals with type 2 diabetes: the Renal Insufficiency And Cardiovascular Events (RIACE) Italian multicentre study. *Diabetologia.* 61(11):2277-2289, 2018.
182. Mantovani A, Zaza G, Byrne CD, Lonardo A, Zoppini G, Bonora E, Targher G. Nonalcoholic fatty liver disease increases risk of incident chronic kidney disease: A systematic review and meta-analysis. *Metabolism.* 79:64-76, 2018.
183. Mirijello A, Viazzi F, Fioretto P, Giorda C, Ceriello A, Russo GT, Guida P, Pontremoli R, De Cosmo S; AMD ANNALS Study Group. Association of kidney disease measures with risk of renal function worsening in patients with type 1 diabetes. *BMC Nephrol.* 19(1):347, 2018.
184. Barutta F, Bellini S, Mastrocola R, Gambino R, Piscitelli F, di Marzo V, Corbetta B, Vemuri VK, Makriyannis A, Annaratone L, Bruno G, Gruden G. Reversal of albuminuria by combined AM6545 and perindopril therapy in experimental diabetic nephropathy. *Br J Pharmacol.* 175(23):4371-4385, 2018.
185. Barutta F, Mastrocola R, Bellini S, Bruno G, Gruden G. Cannabinoid Receptors in Diabetic Kidney Disease. *Curr Diab Rep.* 18(2), 2018.
186. Penno G, Solini A, Bonora E, Orsi E, Fondelli C, Zerbini G, Trevisan R, Vedovato M, Cavalot F, Laviola L, Nicolucci A, Pugliese G; Renal Insufficiency and Cardiovascular Events (RIACE) Study Group. Defining the contribution of chronic kidney disease to all-cause mortality in patients with type 2 diabetes: the Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicenter Study. *Acta Diabetol.* 55(6):603-612, 2018.
187. Iacobini C, Menini S, Blasetti Fantauzzi C, Pesce CM, Giaccari A, Salomone E, Lapolla A, Orioli M, Aldini G, Pugliese G. FL-926-16, a novel bioavailable carnosinase-resistant carnosine derivative, prevents onset and stops progression of diabetic nephropathy in db/db mice. *Br J Pharmacol.* 175(1):53-66, 2018.
188. Fadini GP, Solini A, Manca ML, Zatti G, Karamouzis I, Di Benedetto A, Frittitta L, Avogaro A; DARWIN-T2D Network. Phenotyping normal kidney function in elderly patients with type 2 diabetes: a cross-sectional multicentre study. *Acta Diabetol.* 55(11):1121-1129, 2018.
189. Lamacchia O, Viazzi F, Fioretto P, Mirijello A, Giorda C, Ceriello A, Russo G, Guida P,

Pontremoli R, De Cosmo S. Normoalbuminuric kidney impairment in patients with T1DM: insights from annals initiative. *Diabetol Metab Syndr*. 10:60, 2018.

190. Menini S, Iacobini C, Pugliese G, Pesce C Dietary interventions to contrast the onset and progression of diabetic nephropathy: A critical survey of new data. *Crit Rev Food Sci Nutr*. 58(10):1671-1680, 2018.
191. Pugliese G, Penno G, Natali A, Barutta F, Di Paolo S, Reboldi G, Gesualdo L, De Nicola L; Italian Diabetes Society and the Italian Society of Nephrology. Diabetic kidney disease: New clinical and therapeutic issues. Joint position statement of the Italian Diabetes Society and the Italian Society of Nephrology on "The natural history of diabetic kidney disease and treatment of hyperglycemia in patients with type 2 diabetes and impaired renal function". *Nutr Metab Cardiovasc Dis*, 29(11):1127-1150, 2019.
192. Perrini S. Leptin: a marker of renal injury. *Intern Emerg Med*, 14(4):493-494, 2019.
193. Niewczas MA, Pavkov ME, Skupien J, Smiles A, Md Dom ZI, Wilson JM, Park J, Nair V, Schlafly A, Saulnier PJ, Satake E, Simeone CA, Shah H, Qiu C, Looker HC, Fiorina P, Ware CF, Sun JK, Doria A, Kretzler M, Susztak K, Duffin KL, Nelson RG, Krolewski AS. A signature of circulating inflammatory proteins and development of end-stage renal disease in diabetes. *Nat Med*, 25(5):805-813, 2019.
194. Conserva F, Barozzino M, Pesce F, Divella C, Oranger A, Papale M, Sallustio F, Simone S, Laviola L, Giorgino F, Gallone A, Pontrelli P, Gesualdo L. Urinary miRNA-27b-3p and miRNA-1228-3p correlate with the progression of Kidney Fibrosis in Diabetic Nephropathy. *Sci Rep*, 9(1):11357, 2019.
195. Rossing P, Strand J, Avogaro A, Becka M, Kanefendt F, Otto C. Effects of the chymase inhibitor fulacimstat in diabetic kidney disease-results from the CADA DIA trial. *Nephrol Dial Transplant*. gfaa299, 2020.
196. Penno G, Orsi E, Solini A, Bonora E, Fondelli C, Trevisan R, Vedovato M, Cavalot F, Gruden G, Laviola L, Nicolucci A, Pugliese G; Renal Insufficiency And Cardiovascular Events (RIACE) Study Group. Renal hyperfiltration is independently associated with increased all-cause mortality in individuals with type 2 diabetes: a prospective cohort study. *BMJ Open Diabetes Res Care*. 8(1):e001481, 2020.
197. Lamacchia O, Sorrentino MR, Berti G, Paradiso M, Corrado A, Cantatore FP, De Cosmo S. Glomerular filtration rate is associated with trabecular bone score in patients with type 2 diabetes mellitus. *Diabetes Res Clin Pract*. 164:108164, 2020.
198. Pugliese G, Penno G, Natali A, Barutta F, Di Paolo S, Reboldi G, Gesualdo L, De Nicola L; Italian Diabetes Society and the Italian Society of Nephrology. Diabetic kidney disease: new

clinical and therapeutic issues. Joint position statement of the Italian Diabetes Society and the Italian Society of Nephrology on "The natural history of diabetic kidney disease and treatment of hyperglycemia in patients with type 2 diabetes and impaired renal function". *J Nephrol.* 33(1):9-35, 2020.

- 199.** Di Paolo S, Fiorentino M, De Nicola L, Reboldi G, Gesualdo L, Barutta F, Natali A, Penno G, Fioretto P, Pugliese G; Italian Society of Nephrology and the Italian Diabetes Society. Indications for renal biopsy in patients with diabetes. Joint position statement of the Italian Society of Nephrology and the Italian Diabetes Society. *Nutr Metab Cardiovasc Dis.* 30(12):2123-2132, 2020.
- 200.** Vitale M, Haxhi J, Cirrito T, Pugliese G. Renal protection with glucagon-like peptide-1 receptor agonists. *Curr Opin Pharmacol.* 54:91-101, 2000.
- 201.** Benetti E, Liberto E, Bressanello D, Bordano V, Rosa AC, Miglio G, Haxhi J, Pugliese G, Balducci S, Cordero C. Sedentariness and Urinary Metabolite Profile in Type 2 Diabetic Patients, a Cross-Sectional Study. *Metabolites.* 10(5):205, 2020.
- 202.** Mantovani A, Petracca G, Beatrice G, Csermely A, Lonardo A, Schattenberg JM, Tilg H, Byrne CD, Targher G. Non-alcoholic fatty liver disease and risk of incident chronic kidney disease: an updated meta-analysis. *Gut.* [gutjnl-2020-323082](https://doi.org/10.1136/gutjnl-2020-323082), 2020.