

1. Prando R, Cordera R, De Micheli A, Maiello M, Odetti P, Viviani G, Corsi L, Adezati L. I.V. glucose tolerance test: correlation between FFA, glucose and IRI in normal, obese and diabetic subjects. *Acta Diabetol Lat.* 15(5-6):259-72, 1978.
2. Prando R, Cordera R, Odetti P, De Micheli A, Maiello M, Viviani G, Adezati L. Exploration of the early insulin response by two small successive loads of I.V. glucose in normal and obese subjects. *Acta Diabetol Lat.* 15(1-2):53-67, 1978.
3. Bergman R.N., Ider Y.Z., Bowden C.R., Cobelli C. Quantitative estimation of insulin sensitivity. *Am. J. Physiol.* 236: E667 -677, 1979.
4. Cobelli C., Pacini G., Salvan A. On a simple model of insulin secretion. *Med. Biol. Eng. Comput.* 18: 457-463, 1980.
5. Bergman R.N., Cobelli C.: Minimal modeling, partition analysis and the estimation of insulin sensitivity. *Fed. Proc.* 39: 110-115, 1980.
6. Bergman R.N., Cobelli C., Toffolo G.: Minimal model of glucose/insulin dynamics in the intact organism. A novel approach for evaluation of factors controlling glucose tolerance. *Trans. Inst. Meas. Control* 3: 207-216, 1981.
7. Riggio O, Merli M, Cangiano C, Capocaccia R, Cascino A, Lala A, Leonetti F, Mauceri M, Pepe M, Rossi Fanelli F, Savioli M, Tamburrano G, Capocaccia L. Glucose intolerance in liver cirrhosis. *Metabolism.* 31(6):627-34, 1982.
8. Del Prato S., Nosadini R., Tiengo A., Tessari P., Avogaro A., Trevisan R., Valerio A., Muggeo M., Cobelli C., Toffolo G.: Insulin-mediated glucose disposal in Type-I diabetes. Evidence for insulin resistance, *J. Clin. Endocrin. Metab.* 57: 904-910, 1983.
9. Bonora E., Orioli S., Coscelli C., Buzzelli G., Gentilini P., Butturini U. Possible roles of glucagon, growth hormone and free fatty acids in the pathogenesis of insulin resistance of subjects with chronic liver diseases. *Acta Diabetol. Lat.* 21: 241-250, 1984.
10. Riggio O, Merli M, Cantafora A, Di Biase A, Lalloni L, Leonetti F, Miazzi P, Rinaldi V, Rossi-Fanelli F, Tamburrano G, et al. Total and individual free fatty acid concentrations in liver cirrhosis. *Metabolism.* 33(7):646-51, 1984.
11. Bonora E., Coscelli C., Butturini U. Relationship between insulin resistance, insulin secretion and insulin metabolism in simple obesity. *Int. J. Obes.* 9: 307-312, 1985.
12. Ferrannini E., Smith D.J., Cobelli C., Toffolo G., Pilo A., DeFronzo R.A.: Effect of insulin on the distribution and disposition of glucose in man. *J. Clin. Invest.* 76: 357-364, 1985.

13. Cordera R, Bertolini S, Andraghetti G, Pistocchi G, de Alessi M, Gherzi R. Insulin receptor binding on red cells of hypertriglyceridemic patients. Effect of a low fat, low carbohydrate diet. *Diabete Metab.* 11(3):137-40, 1985.
14. Cordera R, Andraghetti G, Bonadonna R, Gherzi R, Maiello M, Odetti P, Viviani G, Berri F, Adezati L. Influence of cell age and ketoaminic linkage on rapidglycosylation of hemoglobin in human red cells in vitro. *Horm Metab Res.* 17(4):201-4, 1985.
15. Marchetti P, Benzi L, Trischitta V, Brunetti A, Cecchetti P, Ciccarone AM, Pezzino V, Vigneri R, Navalesi R. Complementary use of gel permeation and reversed-phase liquid chromatography for the analysis of A14-[125I]insulin and its degradation products in isolated human monocytes. *J Chromatogr*; 377: 339-344, 1986.
16. Cobelli C., Pacini G., Toffolo G., Sacca' L.: Estimation of insulin sensitivity and glucose clearance from minimal model: new insights from labelled IVGTT. *Am. J. Physiol.* 250: E591-598, 1986.
17. Trischitta V, Gullo D, Squatrito S, Pezzino V, Goldfine ID, Vigneri R. Insulin internalization into monocytes is decreased in patients with type II diabetes mellitus. *J Clin Endocrinol Metab*; 62:522-528, 1986.
18. Bertolini S, Elicio N, Cordera R, Gapitanio GL, Montagna G, Croce S, Saturnino M, Balestreri R, De Cecco L. Effects of three low-dose oral contraceptive formulations on lipid metabolism. *Acta Obstet Gynecol Scand.*;66(4):327-32, 1987.
19. Bonora E., Manicardi V., Zavaroni I., Coscelli C., Butturini U. Relationships between insulin secretion, insulin metabolism and insulin resistance in mild glucose intolerance. *Diabete Metab.* 13: 116-121, 1987.
20. Purrello F, Gullo D, Brunetti A, Buscema M, Italia S, Goldfine ID, Vigneri R. Direct effects of biguanides on glucose utilization in vitro. *Metabolism*; 36: 774-776, 1987.
21. Trischitta V, Benzi L, Brunetti A, Cecchetti P, Marchetti P, Vigneri R, Navalesi R. Intracellular insulin processing is altered in monocytes from patients with type II diabetes mellitus. *J Clin Endocrinol Metab*; 64: 914-920, 1987.
22. Benzi L, Marchetti P, Brunetti A, Ciccarone AM, Cecchetti P, Trischitta E, Di Cianni G, Squatrito S, Navalesi R. Insulin degradation into monocytes from normal subjects: a high performance liquid chromatographic analysis *J Endocrinol Invest*; 11: 303-307, 1988.

23. Pezzino V, Papa V, Trischitta V, Brunetti A, Goodman PA, Treutelaar MK, Williams JA, Maddux BA, Vigneri R, Goldfine ID. Human insulin receptor radioimmunoassay: applicability to insulin-resistant states. *Am J Physiol*; 257:E451-E457, 1989.
24. Trischitta V, Wong KY, Brunetti A, Scalisi R, Vigneri R, Goldfine ID. Endocytosis, recycling, and degradation of the insulin receptor. Studies with monoclonal antireceptor antibodies that do not activate receptor kinase. *J Biol Chem*; 264: 5041-5046, 1989.
25. Hawley DM, Maddux BA, Patel RG, Wong KY, Mamula PW, Firestone GL, Brunetti A, Verspohl E, Goldfine ID. Insulin receptor monoclonal antibodies that mimic insulin action without activating tyrosine kinase. *J Biol Chem*; 264: 2438-2444, 1989.
26. Giaccari A, Rossetti L. Isocratic high-performance liquid chromatographic determination of the concentration and specific radioactivity of phosphoenolpyruvate and uridine diphosphate glucose in tissue extracts. *J Chromatogr*. 29;497:69-78, 1989.
27. Bonora E., Moghetti P., Zancanaro C., Cigolini M., Querena M., Cacciatori V., Corngnati A., Muggeo M. Estimates of in vivo insulin action in man: comparison of insulin tolerance tests and euglycemic and hyperglycemic glucose clamp studies. *J. Clin. Endocrinol. Metab.* 68: 374-378, 1989.
28. Merli M, Leonetti F, Riggio O, Giaccari A, Romiti A, Sbraccia P, Tamburrano G. Resistance to insulin suppression of plasma free fatty acids in liver cirrhosis. *J Endocrinol Invest*. 13(10):787-95, 1990.
29. Benzi L, Trischitta V, Ciccarone A, Cecchetti P, Brunetti A, Squatrito S, Marchetti P, Vigneri R, Navalesi R. Improvement with metformin in insulin internalization and processing in monocytes from NIDDM patients. *Diabetes*; 39: 844-849, 1990.
30. Zancanaro C., Cigolini M., Bonora E., Moghetti P., Querena M., Muggeo M. Glucose and insulin suppression of plasma free fatty acids in obese subjects with normal glucose tolerance or mild newly-diagnosed type II (non-insulin dependent) diabetics. *J. Endocrinol. Invest*. 13: 55-59, 1990.
31. Rossetti L, Giaccari A, Klein-Robbenhaar E, Vogel LR. Insulinomimetic properties of trace elements and characterization of their in vivo mode of action. *Diabetes*. Oct;39(10):1243-50. 1990.
32. Rossetti L, Giaccari A, DeFronzo RA. Glucose toxicity. *Diabetes Care*.

Jun;13(6):610-30, 1990.

33. Rossetti L, Giaccari A. Relative contribution of glycogen synthesis and glycolysis to insulin-mediated glucose uptake. A dose-response euglycemic clamp study in normal and diabetic rats. *J Clin Invest.* Jun;85(6):1785-92, 1990.
34. Rossetti L, DeFronzo RA, Gherzi R, Stein P, Andraghetti G, Falzetti G, Shulman GI, Klein-Robbenhaar E, Cordera R. Effect of metformin treatment on insulin action in diabetic rats: in vivo and in vitro correlations. *Metabolism.* 39(4):425-35, 1990.
35. Cordera R, Andraghetti G, DeFronzo RA, Rossetti L. Effect of in vivo vanadate treatment on insulin receptor tyrosine kinase activity in partially pancreatectomized diabetic rats. *Endocrinology.* 126(4):2177-83, 1990.
36. Frontoni S, Choi SB, Banduch D, Rossetti L. In vivo insulin resistance induced by amylin primarily through inhibition of insulin-stimulated glycogen synthesis in skeletal muscle. *Diabetes.*,40:568-73, 1991.
37. Cobelli C., Ruggeri A.: A reduced sampling schedule for estimating the parameters of the glucose minimal model from a labelled IVGTT. *IEEE Trans. Biomed. Eng. BME-38:* 1023-1029, 1991.
38. Rossetti L, Frontoni S, Dimarchi R, DeFronzo RA, Giaccari A. Metabolic effects of IGF-I in diabetic rats. *Diabetes.*,40:444-8, 1991.
39. Caumo A., Giacca A., Morgese M., Pozza G., Micossi P., Cobelli C.: Minimal models of glucose disappearance: lessons from the labelled IVGTT. *Diabetic Medicine* 8: 822-832, 1991.
40. Choi SB, Frontoni S, Rossetti L. Mechanism by which calcitonin gene-related peptide antagonizes insulin action in vivo. *Am J Physiol.*,260(2 Pt 1):E321-5, 1991.
41. Bruno A, Pagano G, Benzi L, Di Ciani G, Spallone V, Calabrese G, Passeri A. Change in glucose metabolism after long-term treatment with deflazacort and betamethasone. *Eur J Clin Pharmacol* 43:47-50, 1992.
42. Barbetti F., Gejman P.V., Taylor S.I., Raben N., Cama A., Bonora E., Pizzo P., Moghetti P., Muggeo M., Roth J. Detection of mutations in insulin receptor gene by denaturing gradient gel electrophoresis. *Diabetes* 41: 408-415, 1992.
43. Giaccari A, Rossetti L. Predominant role of gluconeogenesis in the hepatic glycogen repletion of diabetic rats. *J Clin Invest.* 89(1):36-45, 1992.
44. Librenti MC, Cocchi M, Orsi E, Pozza G, Micossi P. Effect of soya and cellulose fibers on postprandial glycemic response in type II diabetic patients. *Diabetes Care.*

15:111-113, 1992.

45. Bonora E., Bonadonna R.C., Del Prato S., Gulli G., Solini A., Matsuda M., DeFronzo R.A.: In vivo glucose metabolism in obese and type II diabetic subjects with and without hypertension. *Diabetes* 42: 764-772, 1993.
46. Butler P., Caumo A., Zerman A., O'Brien P., Cobelli C., Rizza R.: Methods for assessment of the rate of onset and offset of insulin action during nonsteady state in humans. *Am. J. Physiol.* 264: E548-560, 1993.
47. Katz H., Butler P., Homan M., Zerman A., Caumo A., Cobelli C., Rizza R.: Hepatic and extrahepatic insulin action in humans: measurement in the absence of non-steady-state error. *Am. J. Physiol.* 264: E561-566, 1993.
48. Del Prato S., Bonadonna R.C., Bonora E., Gulli G., Solini A., Shank M., DeFronzo R.A.: Characterization of cellular defects of insulin action in type II (non-insulin-dependent) diabetes mellitus. *J. Clin. Invest.* 91: 484-494, 1993.
49. Bonadonna R.C., Del Prato S., Saccomani M.P., Bonora E., Gulli G., Ferrannini E., Bier D., Cobelli C., DeFronzo R.A. Transmembrane glucose transport in skeletal muscle of patients with non-insulin-dependent diabetes mellitus. *J. Clin. Invest.* 92: 486-494, 1993.
50. Bonadonna R., Del Prato S., Bonora E., Gulli G., Solini A., DeFronzo R.A. The effects of physiologic hyperinsulinemia on the intracellular metabolic partition of plasma glucose. *Am. J. Physiol.* 265: E943-E953, 1993.
51. Carrera P, Cordera R, Ferrari M, Cremonesi L, Taramelli R, Andraghetti G, Carducci C, Dozio N, Pozza G, Taylor SI, et al. Substitution of Leu for Pro-193 in the insulin receptor in a patient with a genetic form of severe insulin resistance. *Hum Mol Genet* 2(9):1437-41, 1993.
52. Cangemi C, Lombardo G, Sinagra D. Insulin resistance: related clinical syndromes. *Recent Prog Med.* 84(7-8):541-50, 1993.
53. Rossetti L, Farrace S, Choi SB, Giaccari A, Sloan L, Frontoni S, Katz MS. Multiple metabolic effects of CGRP in conscious rats: role of glycogen synthase and phosphorylase. *Am J Physiol* 264(1 Pt 1):E1-10, 1993.
54. Rossetti L, Giaccari A, Barzilai N, Howard K, Sebel G, Hu M. Mechanism by which hyperglycemia inhibits hepatic glucose production in conscious rats. Implications for the pathophysiology of fasting hyperglycemia in diabetes. *J Clin Invest.* 92(3):1126-34., 1993.
55. Moore MC, Shulman GI, Giaccari A, Pagliassotti MJ, Cline G, Neal D, Rossetti L,

- Cherrington AD. Effect of hepatic nerves on disposition of an intraduodenal glucose load. *Am J Physiol.* 265(3 Pt 1):E487-96, 1993.
56. Blasi C., Jeanrenaud B.: "Insulin Resistance Syndrome: Defective GABA Neuromodulation as a Possible Hereditary Pathogenetic Factor (The 'GABA Hypothesis'). *Med Hyp.* 40: 197-206, 1993.
 57. Rodriguez G, Nobili F, Celestino MA, Francione S, Gulli G, Hassan K, Marengo S, Rosadini G, Cordera R. Regional cerebral blood flow and cerebrovascular reactivity in IDDM. *Diabetes Care.* Feb;16(2):462-8, 1993.
 58. Katz H., Homan M., Jensen M., Caumo A., Cobelli C., Rizza R.: Assessment of insulin action in NIDDM in the presence of dynamic changes in insulin and glucose concentration. *Diabetes* 43:289-296, 1994.
 59. Alzaid A., Dinneen S., Turk D., Caumo A., Cobelli C., Rizza R.: Assessment of insulin action and glucose effectiveness in diabetic and nondiabetic humans. *J. Clin. Invest.* 94: 2341-2348, 1994.
 60. Maggi D, Laurino C, Andraghetti G, Cordera R. The overexpression of insulin receptor makes CHO cells resistant to the action of IGF-1: role of IRS-1. *Biochem Biophys Res Commun.* 30;205(1):693-9, 1994.
 61. Laurino C, Bertolini S, Cordera R. Linkage analysis does not support a role for glucokinase gene in the aetiology of type 2 diabetes mellitus among north western Italians. *Mol Cell Endocrinol.* 104(2):147-51, 1994.
 62. Spallarossa P, Cordera R, Andraghetti G, Bertero G, Brunelli C, Caponnetto S. Association between plasma insulin and angiographically documented significant coronary artery disease. *Am J Cardiol.* 15;74(2):177-9, 1994.
 63. Del Prato S, Leonetti F, Simonson DC, Sheehan P, Matsuda M, DeFronzo RA. Effect of sustained physiologic hyperinsulinaemia and hyperglycaemia on insulin secretion and insulin sensitivity in man. *Diabetologia.* 37(10):1025-35, 1994.
 64. Maggi D, Andraghetti G, Cordera R. A ser for Cys mutation in the extracellular portion of insulin receptor beta subunit impairs the insulin-insulin receptor complex internalization in CHO cells. *Biochem Biophys Res Com* 25;210(3):931-7, 1995.
 65. Bonora, E., Gulli G., Bonadonna, R.C., Del Prato, S., Solini, A., DeFronzo, R.A. Insulin sensitivity is not impaired in Mexican American women without family history of diabetes mellitus. *Diabetes Care* 18: 825-833, 1995.
 66. Hwang JH, Perseghin G, Rothman DL, Cline GW, Magnusson I, Petersen KF, Shulman GI. Impaired hepatic glycogen synthesis in insulin dependent diabetic

- subjects: a ^{13}C nuclear magnetic resonance spectroscopy study. *J Clin Invest*; 95: 783-787, 1995.
67. Battezzati A, Terruzzi I, Perseghin G, Bianchi E, Di Carlo V, Pozza G, Luzi L. Defective insulin action on protein and glucose metabolism during chronic hyperinsulinemia in subjects with benign insulinoma. *Diabetes*; 44: 837-844, 1995.
 68. Roden M, Perseghin G, Petersen KF, Hwang JH, Cline GW, Gerow K, Rothman DL, Shulman GI. The roles of insulin and glucagon in the regulation of hepatic glycogen synthesis and turnover in humans. *J Clin Invest*; 97: 642-648, 1996.
 69. Price TB, Perseghin G, Duleba A, Chen W, Chase J, Rothman DL, Shulman RG, Shulman GI. NMR studies of muscle glycogen synthesis in insulin resistant offspring of NIDDM parents immediately following glycogen depleting exercise. *Proc Natl Acad Sci USA*; 93: 5329-5334, 1996.
 70. Avogaro A.P., Vicini P., Valerio A., Caumo A., Cobelli C. The hot but not the cold minimal model allows precise assessment of insulin sensitivity in NIDDM subjects. *Am. J. Physiol.* 33:E532-540, 1996.
 71. Caumo A., Vicini P., and Cobelli C. Is the minimal model too minimal? *Diabetologia*, 39:997-1000, 1996.
 72. Saccomani M.P., Bonadonna R., Bier D.M., De Fronzo R.A., and Cobelli C.: A model to measure insulin effects on glucose transport and phosphorylation in muscle: a three-tracer study. *Am. J. Physiol.*, 33:E170-185, 1996.
 73. Zachwieja J.J., Toffolo G., Cobelli C., Bier D.M., and Yarasheski K.E. Resistance exercise and growth hormone administration in older men: effects on insulin sensitivity and secretion during a stable-label intravenous glucose tolerance test. *Metabolism* 45:254-260, 1996.
 74. Avogaro A., Toffolo G., Miola M., Valerio A., Tiengo A., Cobelli C., Del Prato S. Intracellular lactate- and pyruvate-interconversion rates are increased in muscle tissue of non-insulin-dependent diabetic individuals. *J. Clin. Invest.* 98:1-8, 1996.
 75. Basu A., Alzaid A., Dinneen S., Caumo A., Cobelli C., Rizza R. Effects of a change in the pattern of insulin delivery on carbohydrate tolerance in diabetic and nondiabetic humans in the presence of differing degrees of insulin resistance. *J. Clin. Invest.* 97:2351-2361, 1996.
 76. Roden M, Price TB, Perseghin G, Petersen KF, Cline GW, Shulman GI. Mechanism of free fatty acid induced insulin resistance in humans. *J Clin Invest* 97: 2859-2865, 1996.

77. Velho G, Petersen KF, Perseghin G, Hwang JH, Rothman DL, Pueyo ME, Cline GW, Froguel P, Shulman GI. Impaired hepatic glycogen synthesis in glucokinase deficient (Mody-2) subjects. *J Clin Invest*; 98: 1755-1761, 1996.
78. Perseghin G, Price TB, Petersen KF, Roden M, Cline GW, Gerow K, Rothman DL, Shulman GI. Increased glucose transport/phosphorylation and muscle glycogen synthesis after exercise training in insulin resistant subjects. *N Engl J Med*; 335: 1357-1362, 1996.
79. Bonadonna R., Del Prato S., Bonora E., Saccomani M.P., Gulli G., Natali A., Frascerra S., Pecori N., Ferrannini E., Bier D., Cobelli C., DeFronzo R.A. Roles of glucose transport and glucose phosphorylation in muscle insulin resistance of type 2 (non-insulin-dependent) diabetes mellitus. *Diabetes* 45: 915-925, 1996.
80. Barbeti F, Rocchi M, Bossolasco M, Cordera R, Sbraccia P, Finelli P, Consalez GG. The human skeletal muscle glycogenin gene: cDNA, tissue expression and chromosomal localization. *Biochem Biophys Res Commun.* 7;220(1):72-7. 1996.
81. Spallarossa P, Brunelli C, Minuto F, Caruso D, Battistini M, Caponnetto S, Cordera R. Insulin-like growth factor-I and angiographically documented coronary artery disease. *Am J Cardiol.* 15;77(2):200-2, 1996.
82. Targher G., Alberiche M., Zenere MB, Bonadonna RC, Muggeo M., Bonora E. Cigarette smoking and insulin resistance in patients with non-insulin-dependent diabetes mellitus (NIDDM). *J. Clin. Endocrinol. Metab.* 82: 3619-3624, 1997.
83. Basu A., Caumo A., Bettini F., Gelisio A., Alzaid A., Cobelli C., Rizza R.: Impaired basal glucose effectiveness in NIDDM-Contribution of defects in glucose disappearance and production, measured using an optimized minimal model independent protocol. *Diabetes* 46: 421-432, 1997.
84. Cobelli C., Vicini P., Caumo A.: If the minimal model is too minimal, who suffers more: SG or SI? *Diabetologia* 40: 362-363, 1997.
85. Utriainen T., Nuutila P., Takala T., Vicini P., Ruotsalainen U., Rnnemaa T., Tolvanen T., Raitakari M., Haaparanta M., Kirvel O., Cobelli C., Yki-Jarvinen H.: Intact insulin stimulation of skeletal muscle blood flow, its heterogeneity and redistribution but not of glucose uptake in non-insulin dependent diabetes mellitus. *J. Clin. Invest.* 100: 777-785, 1997.
86. Vicini P., Caumo A., Cobelli C.: The hot IVGTT two compartment minimal model: indices of glucose effectiveness and insulin sensitivity. *Am. J. Physiol.* 273: E1024-1031, 1997.

87. Perseghin G, Ghosh S, Gerow K, Shulman GI. Metabolic defects in lean, non-diabetic offspring of NIDDM parents: a cross-sectional study. *Diabetes*; 46:1001-1009, 1997.
88. Cobelli C., Bettini F., Caumo A, Quon M.J.: Overestimation of minimal model glucose effectiveness in presence of insulin response is due to undermodeling. *Am. J. Physiol.* 275: E1031-1036, 1998.
89. Bonadonna R., Saccomani M.P., Del Prato S., Bonora E., DeFronzo R.A., Cobelli C. Role of tissue specific blood flow and tissue recruitment in insulin-mediated glucose uptake of skeletal muscle. *Circulation* 98: 234-241, 1998.
90. Maggi D, Andraghetti G, Carpentier JL, Cordera R. Cys860 in the extracellular domain of insulin receptor beta-subunit is critical for internalization and signal transduction. *Endocrinology*. 139(2):496-504, 1998.
91. Pinducciu C, Borgonovo G, Arezzo A, Torre GC, Giordano G, Cordera R. Toxic thyroid adenoma: absence of DNA mutations of the TSH receptor and Gs alpha. *Eur J Endocrinol.* 138(1):37-40, 1998
92. Guazzini B, Gaffi D, Mainieri D, Multari G, Cordera R, Bertolini S, Pozza G, Meschi F, Barbetti F. Three novel missense mutations in the glucokinase gene (G80S; E221K; G227C) in Italian subjects with maturity-onset diabetes of the young (MODY). *Mutations in brief* no. 162. Online. *Hum Mutat.*;12(2):136, 1998.
93. Perseghin G, Corno A, Santoro F, Biagioli B, Paolini G, Battezzati A, Benedini S, Donatelli F, Pozza G, Grossi A, Luzi L. Myocardial metabolism studied during warm blood antero-retrograde reperfusion in ischemic human hearts. *Acta Diabetologica*; 35: 67-73, 1998.
94. Perseghin G, Scifo P, De Cobelli F, Pagliato E, Battezzati A, Arcelloni C, Vanzulli A, Testolin G, Pozza G, Del Maschio A, Luzi L. Intramyocellular triglyceride content is a determinant of in vivo insulin resistance in humans: a 1H-13C NMR spectroscopy assessment in offspring of type 2 diabetic parents. *Diabetes*; 48: 1600-1606, 1999.
95. Spallarossa P, Rossettin P, Minuto F, Caruso D, Cordera R, Battistini M, Barreca A, Masperone MA, Brunelli C. Evaluation of growth hormone administration in patients with chronic heart failure secondary to coronary artery disease. *Am J Cardiol.* 84(4):430-3, 1999.
96. Maggi D, Barbetti F, Cordera R. Role of proline 193 in the insulin receptor post-translational processing. *Diabetologia.* Apr;42(4):435-42, 1999.

97. Bonora E., Targher G., Alberiche M., Bonadonna R.C., Saggiani F., Zenere M.B., Uleri S., Muggeo M. Effect of chronic treatment with lacidipine or lisinopril on intracellular partitioning of glucose metabolism in type 2 diabetes mellitus. *J. Clin. Endocrinol. Metab.* 84: 1544-1550, 1999.
98. Merli M, Leonetti F, Riggio O, Valeriano V, Ribaud MC, Strati F, Tisone G, Casciani CU, Capocaccia L. Glucose intolerance and insulin resistance in cirrhosis are normalized after liver transplantation. *Hepatology*. Sep;30(3):649-54, 1999. *Erratum in: Hepatology 2000 Aug;32(2):446. Sprati F [corrected to Strati F].*
99. Pedrinelli R, Dell'Omo G, Giampietro O, Giorgi D, Di Bello V, Bandinelli S, Penno G, Mariani M. Dissociation between albuminuria and insulinaemia in hypertensive and atherosclerotic men. *J Hum Hypertens.*; 13: 129-34, 1999.
100. Graci S., Baratta R, Degano C., Iuppa A., Vigneri R., Frittitta L., Trischitta V. The intravenous insulin tolerance test is an accurate method for screening a general population for insulin resistance and related abnormalities. *J Endocrinol Invest* 22: 472-475, 1999.
101. Frittitta L., Camastra S., Baratta R., Costanzo B., D'Adamo M., Graci S, Spampinato D., Maddux BA., Vigneri R, Ferrannini E., Trischitta V. A soluble PC-1 circulates in human plasma: a new marker of insulin resistance and associated abnormalities. *J Clin Endocrinol Metab* 84: 3620-3625, 1999.
102. Bonora E., Targher G., Alberiche M., Bonadonna R.C., Saggiani F., Zenere M.B., Monauni T., Muggeo M. Homeostasis Model Assessment (HOMA) closely mirrors the glucose clamp technique in the assessment of insulin sensitivity. Studies in subjects with various degrees of glucose tolerance and insulin sensitivity. *Diabetes Care* 23: 57-63, 2000.
103. Hoffman R.P., Vicini P., Sivitz W.I., Cobelli C.: Pubertal adolescent male-female differences in insulin sensitivity and glucose effectiveness determined by the one compartment minimal model. *Pediatric Research* 48:384-388, 2000.
104. Bonora E. Relationship between regional fat distribution and insulin resistance. *Int. J. Obes.* 24 (suppl. 2): S32-S35, 2000.
105. Monauni T., Zenti M.G., Cretti A., Daniels M., Targher G., Caruso B., Caputo M., McClain D., Del Prato S., Giaccari A., Muggeo M., Bonora E., Bonadonna R. Effects of glucosamine infusion *Diabetes* 49: 926-935, 2000.
106. Cella F, Giordano G, Cordera R. Serum leptin concentrations during the menstrual cycle in normal-weight women: effects of an oral triphasic estrogen-

progestin medication. *Eur J Endocrinol.* 142(2):174-8, 2000.

107. Nielsen M.F., Nyholm B., Caumo A., Chandramouli V., Schumann W.C., Cobelli C., Landau, B.R., Rizza R.A., Schmitz O.: Prandial glucose effectiveness and fasting gluconeogenesis in insulin resistant first-degree relatives of patients with type 2 diabetes. *Diabetes* 49:2135-2141, 2000.
108. Caumo A., Bergman RN., Cobelli C.: Insulin sensitivity from meal tolerance tests in normal subjects: a minimal model index. *J. Clin. End. Metab.* 85:4396-4402, 2000.
109. Muggeo M, Bolli G, Bompiani G, Brunetti P, Capani F, Cavallo-Perin P, Comaschi M, Cordera R, Crepaldi G, Cucinotta D, Di Mario U, Fedele D, Ferrannini E, Giorgino R, Giugliano D, Gnudi A, Mancini M, Menzinger G, Pagano G, Pontiroli AE, Pozza G, Santeusano F, Tiengo BA, Trovati M, Vigneri R. Glycemic control and cardiovascular diseases in Type 2 diabetes mellitus. Beyond fasting glycemia and glycosylated hemoglobin. *Diabetes Nutr Metab.* 13(4):182-5, 2000.
110. Perseghin G, Mazzaferro V, Piceni Sereni L, Regalia E, Benedini S, Bazzigaluppi E, Pulvirenti A, Antonio Silva Leão A, Calori G, Romito R, Baratti D, Luzi L. Contribution of reduced insulin sensitivity and secretion to the pathogenesis of hepatogenous diabetes: effect of liver transplantation. *Hepatology*; 31: 694-703, 2000.
111. Perseghin G. Pathogenesis of obesity and diabetes mellitus: insights provided by indirect calorimetry in humans. *Acta Diabetol*; 38: 7-21, 2001.
112. Perseghin G, Scifo P, Pagliato E, Battezzati A, Soldini L, Benedini S, Testolin G, Del Maschio A, Luzi L. Gender factors affect fatty acids-induced insulin resistance in nonobese humans: effects of oral steroidal contraception. *J Clin Endocrinol Metab*; 86: 3188-3196, 2001.
113. Sunehag A.L., Treuth M.S., Toffolo G., Butte N.F., Cobelli C., Bier D.M., Haymond M.W.: Glucose production, gluconeogenesis, and insulin sensitivity in children and adolescents: an evaluation of their reproducibility. *Pediatr. Res.* 50: 115-23, 2001.
114. Perseghin G, Caumo A, Caloni M, Testolin G, Luzi L. Incorporation of the fasting plasma FFA concentration into QUICKI improves its association with insulin sensitivity in nonobese individuals. *J Clin Endocrinol Metab*; 86: 4776-4781, 2001.

115. Spallarossa P, Schiavo M, Rossettin P, Cordone S, Olivotti L, Cordera R, Brunelli C. Sulfonylurea treatment of type 2 diabetic patients does not reduce the vasodilator response to ischemia. *Diabetes Care*. 24(4):738-42, 2001.
116. Maggi D, Cordera R. Cys 786 and Cys 776 in the posttranslational processing of the insulin and IGF-I receptors. *Biochem Biophys Res Commun*. 280(3):836-41, 2001.
117. Bonora E., Targher G., Alberiche M., Bonadonna R.C., Zenere M.B., Saggiani F., Muggeo M. Intracellular partition of plasma glucose disposal in hypertensive and normotensive subjects with type 2 diabetes mellitus. *J. Clin. Endocrinol. Metab*. 86: 2073-2079, 2001.
118. Targher G., Bonadonna R.C., Alberiche M.A., Zenere M.B., Muggeo M., Bonora E. Relationship between soluble adhesion molecules and insulin sensitivity in type 2 diabetic individuals. Role of adipose tissue. *Diabetes Care* 24: 1961-1966, 2001.
119. Bonora E., Targher G., Alberiche M., Formentini G., Calcaterra F., Lombardi S., Marini F., Poli M., Zenari L., Raffaelli A., Perbellini S., Zenere M.B., Saggiani F., Bonadonna R.C., Muggeo M. Predictors of insulin sensitivity in type 2 diabetes mellitus. *Diabet. Med*. 19: 535-542, 2002.
120. Leonetti F, Iacobellis G, Zappaterreno A, Di Mario U. Clinical, physiopathological and dietetic aspects of metabolic syndrome. *Dig Liver Dis*;34 Suppl 2:S134-9, 2002.
121. Dalla Man C., Caumo A., Cobelli C.: The oral glucose minimal model: estimation of insulin sensitivity from a meal test. *IEEE Trans. Biomed. Eng*. 49: 419-429, 2002.
122. Hribal ML, Oriente F, Accili D. Mouse models of insulin resistance. *Am J Physiol Endocrinol Metab*. 282:E977-81, 2002.
123. Hoffman R.P., Vicini P., Cobelli C. Comparison of insulin sensitivity and glucose effectiveness determined by the one- and two-compartment-labeled minimal model in late prepubertal children and early adolescents. *Metabolism* 51: 1582-1586, 2002.
124. Maggi D, Biedi C, Segat D, Barbero D, Panetta D, Cordera R. IGF-I induces caveolin 1 tyrosine phosphorylation and translocation in the lipid rafts. *Biochem Biophys Res Commun*. Aug 2;295(5):1085-9, 2002.
125. Perseghin G, Caumo A, Arcelloni C, Benedini S, Lanzi R, Pagliato E, Piceni

- Sereni L, Testolin G, Battezzati A, Comi G, Comola M, Luzi L. Contribution of abnormal insulin secretion and insulin resistance to the pathogenesis of type 2 diabetes in myotonic dystrophy. *Diabetes Care*; 26: 2112-2118, 2003.
126. Perseghin G, Petersen KF, Shulman GI. Cellular mechanisms of insulin resistance; potential links with inflammation. *Int J Obes*; 27: S6-S11, 2003.
 127. Perseghin G, Lattuada G, Danna M, Piceni Sereni L, Maffi P, Battezzati A, De Cobelli F, Secchi A, Del Maschio A, Luzi L. Insulin resistance, intramyocellular lipid content and plasma adiponectin concentrations in patients with type 1 diabetes. *Am J Physiol Endocrinol Metab*; 285: E1174-E1181, 2003.
 128. Briatore L, Andraghetti G, Cordera R. Acute plasma glucose increase, but not early insulin response, regulates plasma ghrelin. *Eur J Endocrinol*. 149(5):403-6, 2003.
 129. Biedi C, Panetta D, Segat D, Cordera R, Maggi D. Specificity of insulin-like growth factor I and insulin on Shc phosphorylation and Grb2 recruitment in caveolae. *Endocrinology*. 144(12):5497-503, 2003.
 130. Benzi L, Cecchetti P, Ciccarone AM, Novelli S, Paoli A, Bertacca A, Maffei M, Maggi D, Andraghetti G, Del Prato S, Cordera R. The extracellular portion of the insulin receptor beta-subunit regulates the cellular trafficking of the insulin-insulin receptor complex. Studies on Chinese hamster ovary cells carrying the Cys 860-->Ser insulin receptor mutation. *Eur J Endocrinol*. 148(3):365-71, 2003.
 131. Ronchi CL, Orsi E, Giavoli C, Cappiello V, Epaminonda P, Beck-Peccoz P, Arosio M. Evaluation of insulin resistance in acromegalic patients before and after treatment with somatostatin analogues. *J Endocrinol Invest*. 26:533-538, 2003.
 132. Giavoli C, Porretti S, Ronchi CL, Cappiello V, Ferrante E, Orsi E, Arosio M, Beck-Peccoz P. Long-term monitoring of insulin sensitivity in growth hormone-deficient adults on substitutive recombinant human growth hormone therapy. *Metabolism*. 53:740-743, 2004.
 133. Balducci S, Leonetti F, Di Mario U, Fallucca F. Is a long-term aerobic plus resistance training program feasible for and effective on metabolic profiles in type 2 diabetic patients? *Diabetes Care*. 27(3):841-2, 2004.
 134. Filippi E, Sentinelli F, Trischitta V, Romeo S, Arca M, Leonetti F, Di Mario U, Baroni MG. Association of the human adiponectin gene and insulin resistance. *Eur J Hum Genet*. 12(3):199-205, 2004.
 135. Dalla Man C., Caumo A., Basu R., Rizza R., Toffolo G., Cobelli C.: Minimal

- model estimation of glucose absorption and insulin sensitivity from oral test: validation with a tracer method. *Am. J. Physiol.* 287:E637-343, 2004.
- 136.** Spallarossa P, Barsotti A, Cordera R, Ghigliotti G, Maggi D, Brunelli C. Reduction of cardiovascular morbidity and mortality in type 2 diabetes. A rational approach to hypoglycemic therapy. *J Endocrinol Invest.* 27(5):485-95, 2004.
- 137.** Panetta D, Biedi C, Repetto S, Cordera R, Maggi D. IGF-I regulates caveolin 1 and IRS1 interaction in caveolae. *Biochem Biophys Res Commun.* , 316(1):240-3, 2004.
- 138.** Hanusch-Enserer U, Cauza E, Spak M, Endler G, Dunky A, Tura A, Wagner O, Rosen HR, Pacini G, Prager R. Improvement of insulin resistance and early atherosclerosis in patients after gastric banding. *Obes Res.* 12(2):284-91, 2004.
- 139.** Dell'Omo G, Penno G, Pucci L, Mariani M, Del Prato S, Pedrinelli R. Abnormal capillary permeability and endothelial dysfunction in hypertension with comorbid Metabolic Syndrome. *Atherosclerosis.*; 172: 383-389, 2004.
- 140.** Pontiroli AE, Pizzocri P, Caumo A, Perseghin G, Luzi L. Evaluation of insulin release and insulin sensitivity through oral glucose tolerance test: differences between NGT, IFG, IGT, and type 2 diabetes mellitus. A cross-sectional and follow-up study. *Acta Diabetol*; 41: 70-76, 2004.
- 141.** Perseghin G, Comola M, Scifo P, Benedini S, De Cobelli F, Lanzi R, Costantino F, Lattuada G, Battezzati A, Del Maschio A, Luzi L. Postabsorptive and insulin-stimulated energy and protein metabolism in patients with Myotonic Dystrophy type 1. *Am J Clin Nutr*; 80: 357-364, 2004.
- 142.** Lattuada G, Piceni Sereni L, Ruggieri D, Scollo A, Benedini S, Ragona F, Costantino F, Battezzati A, Luzi L, Perseghin G. Postabsorptive and insulin-stimulated energy homeostasis and leucine turnover in offspring of T2DM patients *Diabetes Care* 27: 2716-2722, 2004.
- 143.** Lattuada G, Costantino F, Caumo A, Ragona F, Scifo P, De Cobelli F, Del Maschio A, Luzi L, Perseghin G. Reduced whole body lipid oxidation is associated with insulin resistance but not with intramyocellular lipid content in offspring of type 2 diabetic patients. *Diabetologia*; 48: 741-747, 2005.
- 144.** Dalla Man C., Caumo A., Basu R., Rizza R., Toffolo G., Cobelli C.: Measurement of selective effect of insulin on glucose disposal from labeled glucose oral test minimal model. *Am. J. Physiol. Endocrinol. Metab.* 289:E909-914, 2005.
- 145.** Dalla Man C., Yarasheski K.E., Caumo A., Robertson H., Toffolo G., Polonsky

- K.S., Cobelli C.: Insulin sensitivity by oral glucose minimal models: validation against clamp. *Am. J. Physiol. Endocrinol. Metab.* 289:E954-959, 2005.
- 146.** Sesti G, Sciacqua A, Cardellini M, Marini MA, Maio R, Vatrano M, Succurro E, Lauro R, Federici M, Perticone F. Plasma concentration of insulin-like growth factor-I is independently associated with insulin sensitivity in subjects with different degree of glucose tolerance. *Diabetes Care* 28: 120-125, 2005.
- 147.** Irace C, Cortese C, Fiaschi E, Carallo C, Sesti G, Farinaro E, Gnasso A. Components of the metabolic syndrome and carotid atherosclerosis. *Hypertension* 45: 597-601, 2005.
- 148.** Repetto S, Salani B, Maggi D, Cordera R. Insulin and IGF-I phosphorylate eNOS in HUVECs by a caveolin-1 dependent mechanism. *Biochem Biophys Res Commun.* 337(3):849-52, 2005.
- 149.** Salani B, Repetto S, Cordera R, Maggi D. Glimepiride activates eNOS with a mechanism Akt but not caveolin-1 dependent. *Biochem Biophys Res Commun.*;335(3):832-5, 2005.
- 150.** Foti D, Chieffari E, Fedele M, Iuliano R, Brunetti L, Paonessa F, Manfioletti G, Barbetti F, Brunetti A, Croce CM, Fusco A, Brunetti A. Lack of the architectural factor HMGA1 causes insulin resistance and diabetes in humans and mice. *Nat Med*; 11: 765-773, 2005.
- 151.** Pandolfi A, Solini A, Pellegrini G, Mincione G, Di Silvestre S, Chiozzi P, Giardinelli A, Di Marcantonio MC, Piccirelli A, Capani F, Consoli A. Selective insulin resistance affecting nitric oxide release but not plasminogen activator inhibitor-1 synthesis in fibroblasts from insulin-resistant individuals. *Arterioscler Thromb Vasc Biol.* 25(11):2392-7, 2005.
- 152.** Dell'Omo G, Penno G, Pucci L, Pellegrini G, Scotti A, Del Prato S, Pedrinelli R. The vascular effects of doxazosin in hypertension complicated by metabolic syndrome. *Coron Artery Dis.*; 16: 67-73, 2005.
- 153.** Dell'Omo G, Penno G, Del Prato S, Pedrinelli R. Chlorthalidone improves endothelial-mediated vascular responses in hypertension complicated by nondiabetic metabolic syndrome. *J Cardiovasc Pharmacol Ther.*; 10: 265-272, 2005.
- 154.** Perseghin G. Muscle lipid metabolism in the metabolic syndrome. *Curr Opin Lipidol* 16: 416-420, 2005.
- 155.** Perseghin G, Lattuada G, De Cobelli F, Esposito A, Costantino F, Canu T, Ragona F, Scifo P, De Taddeo F, Maffi P, Secchi A, Del Maschio A, Luzi L.

- Reduced intra-hepatic fat content is associated with increased whole body lipid oxidation in patients with type 1 diabetes. *Diabetologia*; 48: 2615-2621, 2005.
- 156.** Fragasso G, Perseghin G, De Cobelli F, Esposito A, Pallosi A, Lattuada G, Scifo P, Calori G, Del Maschio A, Margonato A. Effects of metabolic modulation by trimetazidine on left ventricular function and phosphocreatine/adenosine triphosphate ratio in patients with heart failure. *Eur Heart J*; 27: 942-948, 2006.
- 157.** Pilonetto G., Caumo A., Sparacino G., Cobelli C.: A new dynamic index of insulin sensitivity. *IEEE Trans. Biomed. Eng.* 53:369-79, 2006.
- 158.** De Cobelli F, Pieroni M, Esposito A, Belloni E, Chimenti C, Mellone R, Canu T, Perseghin G, Gaudio C, Maseri A, Frustaci A, Del Maschio A. Delayed gadolinium-enhanced cardiac magnetic resonance in patients with chronic myocarditis presenting with heart failure or recurrent arrhythmias. *J Am Coll Cardiol*; 47: 1649-1654, 2006.
- 159.** Fragasso G, Montano C, Perseghin G, Pallosi A, Calori G, Lattuada G, Oggionni S, Bassanelli G, Locatelli M, Lopaschuk G, Margonato A. The anti-ischemic effect of trimetazidine in patients with postprandial myocardial ischemia is unrelated to meal composition. *Am Heart J*; 151: 1238.e1-1238.e8, 2006.
- 160.** Perseghin G, Burska A, Lattuada G, Alberti G, Costantino F, Ragogna F, Oggionni S, Scollo A, Terruzzi I, Luzi L. Increased serum resistin in elite endurance athletes with high insulin sensitivity. *Diabetologia*; 49:1893-1900, 2006.
- 161.** Perseghin G, Bonfanti R, Magni S, Lattuada G, De Cobelli F, Canu T, Esposito A, Scifo P, Ntali G, Costantino F, Bosio L, Ragogna F, Del Maschio A, Chiumello G, Luzi L. Insulin resistance and whole body energy homeostasis in obese adolescents with fatty liver disease. *Am J Physiol Endocrinol Metab*; 291: E697-E703, 2006.
- 162.** Caumo A, Perseghin G, Brunani A, Luzi L. New insights on the simultaneous assessment of insulin sensitivity and β -cell function with the HOMA2 method. *Diabetes Care* 29: 2733-2734, 2006.
- 163.** Perseghin G, Lattuada G, De Cobelli F, Ntali G, Esposito A, Burska A, Belloni E, Canu T, Ragogna F, Scifo P, Del Maschio A, Luzi L. Serum resistin and intra-hepatic fat content in non-diabetic individuals. *J Clin Endocrinol Metab* 91: 5122-5125, 2006.
- 164.** Salani B, Briatore L, Andraghetti G, Adami GF, Maggi D, Cordera R. High-molecular weight adiponectin isoforms increase after biliopancreatic diversion in

- obese subjects. *Obesity* (Silver Spring). 14(9):1511-4, 2006.
165. Penno G, Miccoli R, Pucci L, Del Prato S. The metabolic syndrome. Beyond the insulin resistance syndrome. *Pharmacol Res.*; 53: 457-68, 2006.
 166. Gilardini L, McTernan PG, Girola A, da Silva NF, Alberti L, Kumar S, Invitti C. Adiponectin is a candidate marker of metabolic syndrome in obese children and adolescents. *Atherosclerosis* 189: 401-407, 2006.
 167. Invitti C., Maffei C., Gilardini L., Pontiggia B., Mazzilli G., Morabito F., Viberti G.C. Metabolic syndrome in obese Caucasian children: prevalence using WHO-derived criteria and association with non traditional cardiovascular risk factors. *Int J Obesity* 30:627-633, 2006.
 168. Amato, M.C., Galluzzo, A., Merlino, S., Mattina, A., Richiusa, P., Criscimanna, A., Giordano, C Lower insulin sensitivity differentiates hirsute from non-hirsute Sicilian women with polycystic ovary syndrome.. *European Journal of Endocrinology* 155 (6):859-865, 2006.
 169. De Filippis E, Cusi K, Ocampo G, Berria R, Buck S, Consoli A, Mandarino LJ. Exercise-induced improvement in vasodilatory function accompanies increased insulin sensitivity in obesity and type 2 diabetes mellitus. *J Clin Endocrinol Metab.* 91:4903-10, 2006.
 170. Formoso G, Chen H, Kim J, Montagnani M, Consoli A, Quon M DHEA mimics acute actions of insulin to stimulate production of both NO and ET-1 via distinct PI 3-kinase- and MAP-kinase-dependent pathways in vascular endothelium. *Mol Endocrinol* 20:1153-63, 2005.
 171. Lo Iacono, O., Venezia, G., Petta, S., Mineo, C., De Lisi, S., Di Marco, V., Rodolico, V., Giordano C., Almasio PL, Craxí, A. The impact of insulin resistance, serum adipocytokines and visceral obesity on steatosis and fibrosis in patients with chronic hepatitis C. *Alimentary Pharmacology and Therapeutics* 25 (10):1181-1191, 2007.
 172. Bock G., Chittilapilly E., Basu R., Toffolo G., Cobelli C., Chandramouli V., Landau B.R., Rizza R.A.: Contribution of hepatic and extrahepatic insulin resistance to the pathogenesis of impaired fasting glucose: role of increased rates of gluconeogenesis. *Diabetes* 56:1703-1711, 2007.
 173. Bianchi C, Penno G, Romero F, Del Prato S, Miccoli R. Treating the metabolic syndrome. *Expert Rev Cardiovasc Ther.*; 5: 491-506, 2007
 174. Sesti G, Capaldo B, Cavallo Perin P, Del Prato S, Frittitta L, Frontoni S, Hribal

ML, Marchesini G, Paolisso G, Piatti PM, Solini A, Bonora E, on behalf of the Group of Italian Scientists of Insulin Resistance (GISIR). Correspondence Between the International Diabetes Federation Criteria for Metabolic Syndrome and Insulin Resistance in a Cohort of Italian Nondiabetic Caucasians. The GISIR database. *Diabetes Care* 30: e33, 2007.

175. Madonna R, Massaro M, Pandolfi A, Consoli A, De Caterina R. The prominent role of p38 mitogen-activated protein kinase in insulin-mediated enhancement of VCAM-1 expression in endothelial cells. *Int J Immunopathol Pharmacol.*20(3):539-55., 2007.
176. Andreozzi F, Succurro E, Mancuso MR, Perticone M, Sciacqua A, Perticone F, Sesti G. Metabolic and cardiovascular risk factors in subjects with impaired fasting glucose: the 100 versus 110 mg/dL threshold. *Diabetes Metab Res Rev* 23: 547-550 2007.
177. Bonora E., Kiechl S., Willeit J., Oberhollenzer F., Egger G., Meigs J., Bonadonna R., Muggeo M. Insulin resistance as estimated by homeostasis model assessment predicts incident symptomatic cardiovascular disease in Caucasian subjects from the general population. The Bruneck Study. *Diabetes Care* 30: 318-324, 2007.
178. Perseghin G, Lattuada G, De Cobelli F, Ragona F, Ntali G, Esposito A, Belloni E, Canu T, Terruzzi I, Scifo P, Del Maschio A, Luzi L. Habitual physical activity is associated with the intra-hepatic fat content in humans *Diabetes Care*; 30: 683-688, 2007.
179. Di Filippo C, Verza M, Coppola L, Rossi F, D'Amico M, Marfella R. Insulin resistance and postprandial hyperglycemia the bad companions in natural history of diabetes: effects on health of vascular tree. *Curr Diabetes Rev.* Nov;3(4):268-73, 2007.
180. Abbatecola AM, Ferrucci L, Marfella R, Paolisso G. Insulin resistance and cognitive decline may be common soil for frailty syndrome. *Arch Intern Med.* 167(19):2145-6, 2007.
181. Caumo A, Perseghin G, Brunani A, Luzi L. Disposition index by the HOMA2 method. Reply to Nagasaka et al. *Diabetes Care* 30: e43, 2007.
182. Fragasso G, Pallosi A, Perseghin G. Letter regarding article by Tuunanen et al, "Free fatty acid depletion acutely decreases cardiac work and efficiency in cardiomyopathic heart failure". *Circulation*; 115: e546, 2007.

183. Morbiducci U, Di Benedetto G, Kautzky-Willer A, Pacini G, Tura A. Improved usability of the minimal model of insulin sensitivity based on an automated approach and genetic algorithms for parameter estimation. *Clin Sci (Lond)*. 112(4):257-63, 2007.
184. Amato, M.C., Galluzzo, A., Finocchiaro, S., Criscimanna, A., Giordano, C. The evaluation of metabolic parameters and insulin sensitivity for a more robust diagnosis of the polycystic ovary syndrome. *Clinical Endocrinology* 69 (1): 52-60, 2008.
185. Galluzzo, A., Amato, M.C., Giordano, C. Insulin resistance and polycystic ovary syndrome. *Nutrition, Metabolism and Cardiovascular Diseases* 18 (7): 511-518, 2008.
186. Perseghin G, Lattuada G, De Cobelli F, Esposito A, Belloni E, Ntali G, Canu T, Ragogna F, Scifo P, Del Maschio A, Luzi L. Increased mediastinal fat and impaired left ventricular energy metabolism in young men with newly found fatty liver *Hepatology*; 47: 51-58, 2008.
187. Bellentani S, Delle Grave R, Suppini A, Marchesini G, Bedogni G, Bugianesi E, Crocè L, Gastaldelli A, Marra F, Perseghin G, Svegliati-Baroni G, Tell G, Tiribelli C. Behavior therapy for nonalcoholic fatty liver disease: the need for a multidisciplinary approach. *Hepatology*; 47: 746-754, 2008.
188. Perseghin G. Is a nutritional therapeutic approach unsuitable for metabolically healthy but obese women? *Diabetologia*; 51: 1567-1569, 2008.
189. Bugianesi E, Bellentani S, Bedogni G, Tiribelli C, and the Fatty Liver Italian Network (FLIN) (Members of FLIN: Svegliati-Baroni G, Crocè LS, Gastaldelli A, Marchesini G, Marra F, Perseghin G, Tell G). Clinical update on non-alcoholic fatty liver disease and steatohepatitis. *Annals of Hepatology* 7: 157-160, 2008.
190. Belloni E, De Cobelli F, Esposito A, Mellone R, Perseghin G, Canu T, Del Maschio A. MRI of cardiomyopathy *AJR Am J Roentgenol* 191: 1702-1710, 2008.
191. Ghiadoni L, Penno G, Giannarelli C, Plantinga Y, Bernardini M, Pucci L, Miccoli R, Taddei S, Salvetti A, Del Prato S. Metabolic syndrome and vascular alterations in normotensive subjects at risk of diabetes mellitus. *Hypertension*.; 51: 440-445, 2008.
192. Plantinga Y, Ghiadoni L, Magagna A, Giannarelli C, Penno G, Pucci L, Taddei S, Del Prato S, Salvetti A. Peripheral wave reflection and endothelial function in untreated essential hypertensive patients with and without the metabolic syndrome.

J Hypertens.; 26: 1216-1222, 2008.

193. Sesti G, Andreozzi F, Bonadonna R, De Mattia G, Leonetti F, Luzi L, Marini MA, Natali A, Vettor R, Bonora E. on behalf of the Group of Italian Scientists of Insulin Resistance (GISIR). Impact of lowering the criterion for impaired fasting glucose on identification of individuals with insulin resistance. The GISIR database. *Diabetes Metab Res Rev* 24:130-136, 2008.
194. Luzi L., Zoppini G., Targher G., Battezzati A., Muggeo M., Bonora E. Insulin effect on serum potassium and auto-inhibition of insulin secretion is intact in a patient with leprechaunism despite severe impairment of substrates metabolism. *Diabetes Metab. Res. Rev.* 24: 205-210, 2008.
195. Bonora E., Capaldo B., Cavallo Perin P., Del Prato S., De Mattia G., Frittitta L., Frontoni S., Legnetti F., Luzi L., Marchesini G., Marini M.A., Natali A., Paolisso G., Piatti P.M., Pujia A., Solini A., Vettor R., Bonadonna R.C. on behalf of the Group of Italian Scientists of Insulin Resistance (GISIR). Hyperinsulinemia and insulin resistance are independently associated with plasma lipids, uric acid and blood pressure in nondiabetic subjects. The GISIR database. *Nutr. Metab. Cardiovasc. Dis.* 18: 624-631, 2008.
196. Puddu A, Salani B, Cordera R, Viviani GL, Maggi D. Caveolin-1 is essential for glimepiride-induced insulin secretion in the pancreatic betaTC-6 cell line. *Biochem Biophys Res Commun.* 375(2):235-7, 2008.
197. Resmini E, Andraghetti G, Rebora A, Cordera R, Vera L, Giusti M, Minuto F, Ferone D. Leptin, ghrelin, and adiponectin evaluation in transsexual subjects during hormonal treatments. *J Androl.* 29(5):580-5, 2008.
198. Briatore L, Salani B, Andraghetti G, Danovaro C, Sferrazzo E, Scopinaro N, Adami GF, Maggi D, Cordera R. Restoration of acute insulin response in T2DM subjects 1 month after biliopancreatic diversion. *Obesity (Silver Spring).* 16(1):77-81, 2008.
199. Benedini S, Cortelli P, Caumo A, Terruzzi I, Beelke M, Perseghin G, Montagna P, Luzi L. Insulin resistance to both glucose and aminoacid metabolism in a patient with Fatal Familial Insomnia. *Nutr Metab Cardiovasc Dis,* 18(9):e47-8, 2008.
200. Salani B, Briatore L, Garibaldi S, Cordera R, Maggi D. Caveolin-1 down-regulation inhibits insulin-like growth factor-I receptor signaltransduction in H9C2 rat cardiomyoblasts. *Endocrinology.* 149(2):461-5, 2008.
201. Consoli C, Martelli E, D'Adamo M, Menghini R, Arcelli D, Porzio O, Pandolfi

- A, Pistolese GR, Consoli A, Lauro R, Ippoliti A, Federici M. Insulin resistance affects gene expression in endothelium. *Arterioscler Thromb Vasc Biol.* 28(2):e7-9, 2008.
- 202.** Dalla Man C., Toffolo G., Basu R., Rizza R.A., Cobelli C. Use of labeled oral minimal model to measure hepatic insulin sensitivity. *Am. J. Physiol. Endocrinol. Metab.* 295:E1152-9, 2008.
- 203.** Basu A., Dalla Man C., Basu R., Toffolo G., Cobelli C., Rizza R.A.: Effects of Type 2 diabetes on insulin secretion, insulin action, glucose effectiveness and postprandial glucose metabolism. *Diabetes Care* 32:866-72, 2009.
- 204.** Dalla Man C., Toffolo G., Basu R., Rizza R.A., Cobelli C.: Use of labeled oral minimal model to measure hepatic insulin sensitivity. *Am. J. of Physiol. Endocrinol. Metab.* 295:1152-1159, 2009.
- 205.** Penno G, Pucci L, Dell'Omo G, Lucchesi D, Miccoli R, Del Prato S, Solini A, Pedrinelli R. Soluble CD40 ligand levels in essential hypertensive men: evidence of a possible role of insulin resistance. *Am J Hypertens.* 22: 1007-13, 2009.
- 206.** Succurro E, Marini MA, Grembiale A, Lugarà M, Andreozzi F, Sciaqua A, Hribal ML, Lauro R, Perticone F, Sesti G. differences in cardiovascular risk profile based on relationship between post-load plasma glucose and fasting plasma levels. *Diabetes Metab Res Rev* 25: 351-356, 2009.
- 207.** Succurro E, Marini MA, Arturi F, Grembiale A, Lugarà M, Andreozzi F, Sciacqua A, Lauro R, Hribal ML, Perticone F, Sesti G. Elevated one-hour post-load plasma glucose levels identifies subjects with normal glucose tolerance but early carotid atherosclerosis. *Atherosclerosis* 207: 245–249, 2009.
- 208.** Succurro E, Andreozzi F, Marini MA, Lauro R, Hribal ML, Perticone F, Sesti G. Low plasma insulin-like growth factor-1 levels are associated with reduced insulin sensitivity and increased insulin secretion in nondiabetic subjects. *Nutr Metab Cardiovasc Dis* 19: 713-719, 2009.
- 209.** Salani B, Briatore L, Contini P, Passalacqua M, Melloni E, Paggi A, Cordera R, Maggi D. IGF-I induced rapid recruitment of integrin beta1 to lipid rafts is Caveolin-1 dependent. *Biochem Biophys Res Commun.* 380(3):489-92, 2009.
- 210.** Chiefari E, Paonessa F, Iiritano S, Le Pera I, Palmieri D, Brunetti G, Lupo A, Colantuoni V, Foti D, Gulletta E, De Sarro G, Fusco A, Brunetti A. The cAMP-HMGA1-RBP4 system: a novel biochemical pathway for modulating glucose homeostasis. *BMC Biol*; 7:24, 2009.

211. Carallo C, Mancuso G, Mauro G, Laghi F, Madafferi B, Irace C, Gnasso A, Scavelli F, Dell'Aquila F, Bartone M, Gullo F, Ferraro M, Spagnuolo V, Belmonte M, Ferrara A, Silvano Rotondaro A, Brandolino N, Parasporo F, Scopelliti F. Hepatic steatosis, carotid atherosclerosis and metabolic syndrome: the STEATO Study.. *J Gastroenterol*;44:1156-1161, 2009.
212. Leonardini A, Laviola L, Perrini S, Natalicchio A, Giorgino F. Cross-Talk between PPARgamma and Insulin Signaling and Modulation of Insulin Sensitivity. *PPAR Res*;2009:818945, 2009.
213. Cordera R, Salani B, Briatore L. Insulin analogues: fears, facts and fantasies. *Diabetes Metab Res Rev*. Jan;25(1):50-1, 2009.
214. Esposito A, De Cobelli F, Perseghin G, Pieroni M, Belloni E, Mellone R, Canu T, Gentinetta F, Scifo P, Chimenti C, Frustaci A, Luzi L, Maseri A, Del Maschio A. Impaired left ventricular energy metabolism in patients with hypertrophic cardiomyopathy is related to the extension of fibrosis at gadolinium delayed enhanced MR imaging. *Heart* 95: 228-233, 2009.
215. De Cobelli F, Esposito A, Belloni E, Pieroni M, Perseghin G, Frustaci A, Del Maschio A. Delayed-Gadolinium enhancement cardiac magnetic resonance for differentiation of Anderson-Fabry disease from symmetric hypertrophic cardiomyopathy. *AJR Am J Roentgenol*; 192: W97-102, 2009.
216. Esposito A, De Cobelli F, Perseghin G, Del Maschio A. The Authors response to: "Impaired left ventricular energy metabolism in hypertrophic cardiomyopathy is not due to fibrosis". *Heart*; 95: 505, 2009.
217. Perseghin G, Caumo A, Lattuada G, De Cobelli F, Ntali G, Esposito A, Belloni E, Canu T, Ragogna F, Scifo P, Del Maschio A, Luzi L. Elevated fasting plasma C-peptide occurs in non-diabetic individuals with fatty liver irrespectively of insulin resistance. *Diabet Med*; 26: 847-854, 2009.
218. Perseghin G. Where does insulin resistance start ? The Liver. *Diabetes Care*; 32 (Suppl 2): S164-S167, 2009.
219. Perseghin G, Lattuada G, Ragogna F, Alberti G, La Torre A, Luzi L. Free leptin index and thyroid function in male highly trained athletes. *Eur J Endocrinol*; 161: 871-876, 2009.
220. Perseghin G. The role of non alcoholic fatty liver disease in cardiovascular disease. *Dig Dis*; 28: 210-213, 2010.
221. Perseghin G. Exploring the in vivo mechanisms of action of glucokinase

- activators in type 2 diabetes. *J Clin Endocrinol Metab* 95: 4871-4873, 2010.
222. Natali A, Muscelli E, Mari A, Balkau B, Walker M, Tura A, Anderwald C, Golay A, Ferrannini E; Relationship between Insulin Sensitivity and Cardiovascular disease investigators. Insulin sensitivity and beta-cell function in the offspring of type 2 diabetic patients: impact of line of inheritance. *J Clin Endocrinol Metab*. Oct;95(10):4703-11, 2010.
 223. Mari A, Tura A, Natali A, Laville M, Laakso M, Gabriel R, Beck-Nielsen H, Ferrannini E; RISC Investigators. Impaired beta cell glucose sensitivity rather than inadequate compensation for insulin resistance is the dominant defect in glucose intolerance. *Diabetologia*. 53(4):749-56, 2010.
 224. Chiefari E, Iiritano S, Paonessa F, Le Pera I, Arcidiacono B, Filocamo M, Foti D, Liebhaber SA, Brunetti A. Pseudogene-mediated posttranscriptional silencing of HMGA1 can result in insulin resistance and type 2 diabetes. *Nat Commun* 1:40, 2010.
 225. Pillonetto G., Caumo A., Cobelli C.: Dynamic insulin sensitivity index: importance in diabetes. *Am. J. Physiol. Endocrinol. Metab*. 298:E440-E448, 2010.
 226. Tura A, Sbrignadello S, Succurro E, Groop L, Sesti G, Pacini G. An empirical index of insulin sensitivity from short IVGTT: validation against the minimal model and glucose clamp indices in subjects with different clinical characteristics. *Diabetologia* 53: 144-152, 2010.
 227. Salani B, Passalacqua M, Maffioli S, Briatore L, Hamoudane M, Contini P, Cordera R, Maggi D. IGF-IR internalizes with Caveolin-1 and PTRF/Cavin in HaCat cells. *PLoS One*. 30;5(11):e14157, 2010.
 228. Cimmino MA, Andraghetti G, Briatore L, Salani B, Parodi M, Cutolo M, Cordera R. Changes in adiponectin and leptin concentrations during glucocorticoid treatment: a pilot study in patients with polymyalgia rheumatica. *Ann N Y Acad Sci*. 1193:160-3, 2010.
 229. Marini C, Bezante G, Gandolfo P, Modonesi E, Morbelli SD, Depascale A, Rollando D, Maggi D, Albertelli M, Armonino R, Balbi M, Brunelli C, Cordera R, Sambuceti G. Optimization of flow reserve measurement using SPECT technology to evaluate the determinants of coronary microvascular dysfunction in diabetes. *Eur J Nucl Med Mol Imaging*. 37(2):357-67, 2010.
 230. Muscogiuri G, Sorice GP, Priolella A, Policola C, Della Casa S, Pontecorvi A, Giaccari A. 25-Hydroxyvitamin D concentration correlates with insulin-sensitivity

and BMI in obesity. *Obesity* (Silver Spring). 18(10):1906-10, 2010.

231. Terruzzi I, Senesi P, Magni C, Montesano A, Scarafoni A, Luzi L, Duranti M. Insulin-mimetic action of conglutin-gamma, a lupin seed protein, in mouse myoblasts. *Nutr Metab Cardiovasc Dis*. 2010.
232. Bianchi C, Penno G, Crisci I, Daniele G, Agostini A, Del Prato S, Miccoli R. Serum gamma-glutamyltransferase levels are related to insulin sensitivity and secretion in subjects with abnormal glucose regulation. *Diabetes Metab Res Rev*.; 26: 181-186, 2010.
233. Mari A, Tura A, Natali A, Anderwald C, Balkau B, Lalic N, Walker M, Ferrannini E; RISC Investigators. Influence of hyperinsulinemia and insulin resistance on in vivo β -cell function: their role in human β -cell dysfunction. *Diabetes*. 60(12):3141-7, 2011.
234. Vigili de Kreutzenberg S., Fadini G.P., Boscari F., Rossi E., Guerra S., Sparacino G., Cobelli C., Ceolotto G., Bottero M., Avogaro A.: Impaired hemodynamic response to meal intake in insulin-resistant subjects: an impedance cardiography approach. *Am J Clin Nutr*. 93:926-33, 2011.
235. Anderwald C, Tura A, Grassi A, Krebs M, Szendroedi J, Roden M, Bischof MG, Luger A, Pacini G. Insulin infusion during normoglycemia modulates insulin secretion according to whole-body insulin sensitivity. *Diabetes Care*.;34(2):437-41, 2011.
236. Succurro E, Marini MA, Arturi F, Grembiale A, Fiorentino TV, Andreozzi F, Sciacqua A, Lauro R, Hribal ML, Perticone F, Sesti G. Usefulness of hemoglobin A1C as a criterion to define the metabolic syndrome in a cohort of italian nondiabetic white subjects. *Am J Cardiol* 107:1650-1655, 2011.
237. Mammi C, Pastore D, Lombardo MF, Ferrelli F, Caprio M, Consoli C, Tesaro M, Gatta L, Fini M, Federici M, Sbraccia P, Donadel G, Bellia A, Rosano GM, Fabbri A, Lauro D. Sildenafil reduces insulin-resistance in human endothelial cells. *PLoS One*. 6(1):e14542, 2011.
238. Bianchi C, Miccoli R, Bonadonna RC, Giorgino F, Frontoni S, Faloia E, Marchesini G, Dolci MA, Alviggi L, Gnasso A, Consoli A, Cavalot F, Cavallo MG, Leonetti F, Giaccari A, Del Prato S; GENFIEV Investigators. Metabolic syndrome in subjects at high risk for type 2 diabetes: the genetic, physiopathology and evolution of type 2 diabetes (GENFIEV) study. *Nutr Metab Cardiovasc Dis*.;21:699-705, 2011.

239. Muscogiuri G, Kamat A, Balas B, Giaccari A, Defronzo RA, Musi N, Katz MS. β -Adrenergic Responsive Induction of Insulin Resistance in Liver of Aging Rats. *Endocr Res.*;36(2):74-82, 2011.
240. Sorice GP, Muscogiuri G, Mezza T, Priolella A, Giaccari A. Metabolic syndrome in transplant patients: an academic or a health burden? *Transplant Proc.* Jan-43(1):313-7, 2011.
241. Muscogiuri G, Sorice GP, Priolella A, Mezza T, Cipolla C, Salomone E, Giaccari A, Pontecorvi A, Della Casa S. The size of adrenal incidentalomas correlates with insulin resistance. Is there a cause-effect relationship? *Clin Endocrinol (Oxf)*. Mar;74(3):300-5, 2011.
242. Fragasso G, Montano C, Lattuada G, Salerno A, Palloshi A, Calori G, Luzi L, Perseghin G, Margonato A. A high carbohydrate meal yields a lower ischemic threshold than a high fat meal in patients with stable coronary disease. *Int J Cardiol*, 147: 209-213, 2011.
243. Perseghin G. Lipids in the wrong place: visceral fat and nonalcoholic steatohepatitis (NASH). *Diabetes Care*; 34: S367-S370, 2011.
244. Lattuada G, Ragogna F, Perseghin G. Why does NAFLD predict type 2 diabetes. *Current Diabetes Report*; 11: 167-172, 2011.
245. Calori G, Lattuada G, Ragogna F, Garancini MP, Crosignani P, Villa M, Bosi E, Ruotolo G, Piemonti L, Perseghin G. Fatty Liver Index (FLI) and mortality: the Cremona Study in the 15th year of follow up *Hepatology*; 54: 145-152, 2011.
246. Fragasso G, De Cobelli F, Spoladore R, Esposito A, Salerno A, Calori G, Montanaro C, Maranta F, Lattuada G, Margonato A, Del Maschio A, Perseghin G. Resting cardiac energy metabolism is inversely associated with heart rate in normal young adult males. *Am Heart J*; 162: 136-141, 2011.
247. Fragasso G, Salerno A, Lattuada G, Cuko A, Calori G, Scollo A, Arioli F, Bassanelli G, Spoladore R, Luzi L, Margonato A, Perseghin G. Effect of partial inhibition of fatty-acid oxidation by trimetazidine on whole body energy metabolism in patients with chronic heart failure. *Heart* 97: 1495-1500, 2011.
248. Perseghin G, Lattuada G, De Cobelli F, Esposito A, Canu T, Ragogna F, Maffi P, Scifo P, Secchi A, Del Maschio A, Luzi L. Left ventricular function and energy homeostasis in patients with type 1 diabetes with and without microvascular complications. *Int J Cardiol*; 154: 111–115, 2012.
249. Esposito A, De Cobelli F, Colantoni C, Perseghin G, Del Vecchio A, Canu T,

- Calandrino R, Del Maschio A. Gender influence on the dose saving allowed by prospective-triggered 64-slice multi detector computed tomography coronary angiography as compared with retrospective-gated mode. *Int J Cardiol* 158: 253-259, 2012.
250. Marini MA, Succurro E, Frontoni S, Mastroianni S, Arturi F, Sciacqua A, Lauro R, Hribal ML, Perticone F, Sesti G. Insulin sensitivity, β -cell function, and incretin effect in individuals with elevated 1-h postload plasma glucose levels. *Diabetes Care* 35: 868-872, 2012.
251. GENFIEV Investigators. Pathogenetic mechanisms and cardiovascular risk: differences between HbA(1c) and oral glucose tolerance test for the diagnosis of glucose tolerance. *Diabetes Care* 35:2607-2612, 2012.
252. Bruzzone S, Ameri P, Briatore L, Mannino E, Basile G, Andraghetti G, Grozio A, Magnone M, Guida L, Scarfi S, Salis A, Damonte G, Sturla L, Nencioni A, Fenoglio D, Fiory F, Miele C, Beguinot F, Ruvolo V, Bormioli M, Colombo G, Maggi D, Murialdo G, Cordera R, De Flora A, Zocchi E. The plant hormone abscisic acid increases in human plasma after hyperglycemia and stimulates glucose consumption by adipocytes and myoblasts. *FASEB J.* 26(3):1251-60, 2012.
253. Salani B, Maffioli S, Hamoudane M, Parodi A, Ravera S, Passalacqua M, Alama A, Nhiri M, Cordera R, Maggi D. Caveolin-1 is essential for metformin inhibitory effect on IGF1 action in non-small-cell lung cancer cells. *FASEB J.* 26(2):788-98, 2012.
254. Muscogiuri G, Policola C, Priolella A, Sorice G, Mezza T, Lassandro A, Della Casa S, Pontecorvi A, Giaccari A. Low levels of 25(OH)D and insulin-resistance: 2 unrelated features or a cause-effect in PCOS? *Clin Nutr.* 31(4):476-80, 2012.
255. Gilardini L, Vallone L, Cottafava R, Redaelli G, Croci M, Conti A, Pasqualinotto L, Invitti C Insulin Sensitivity Deteriorates after Short-Term Lifestyle Intervention in the Insulin Sensitive Phenotype of Obesity. *Obesity Facts* 5:68-76, 2012.
256. Mencarelli M, Zulian A, Cancellò R, Alberti L, Gilardini L, Di Blasio AM, Invitti C A novel missense mutation in the signal peptide of the human POMC gene is associated with obesity and early-onset diabetes. *Eur J Hum Genet.*, 20:1290-4. 2012.
257. Masciopinto F, Di Pietro N, Corona C, Bomba M, Pipino C, Curcio M, Di Castelnuovo A, Ciavardelli D, Silvestri E, Canzoniero LM, Sekler I, Pandolfi A,

- Sensi SL. Effects of long-term treatment with pioglitazone on cognition and glucose metabolism of PS1-KI, 3xTg-AD, and wild-type mice. *Cell Death Dis.* 3:e448, 2012.
258. Bonapace S, Perseghin G, Molon G, Canali G, Bertolini L, Zenari L, Zoppini G, Barbieri E, Targher G. Non-alcoholic fatty liver disease is associated with left ventricular diastolic dysfunction in subjects with type 2 diabetes. *Diabetes Care*, 35:389–395, 2012.
 259. De Cobelli F, Esposito A, Perseghin G, Sallemi C, Belloni E, Ravelli S, Lanzani C, Del Maschio A. Intraindividual Comparison of Gadobutrol and Gadopentetate dimeglumine for detection of myocardial late enhancement in Cardiac MRI *AJR Am J Roentgenol*; 198: 809-816, 2012.
 260. Perseghin G, Calori G, Lattuada G, Dugnani E, Garancini MP, Crosignani P, Villa M, Bosi E, Ruotolo G, Piemonti L. Insulin resistance/hyperinsulinemia and cancer mortality: the Cremona Study at the 15th year of follow-up *Acta Diabetol* 49: 421-428, 2012.
 261. Ciresi A, Amato MC, Pizzolanti G, Giordano Galluzzo C Visceral adiposity index is associated with insulin sensitivity and adipocytokine levels in newly diagnosed acromegalic patients.. *J Clin Endocrinol Metab.* 97(8):2907-15, 2012.
 262. Aghemo A, Prati GM, Rumi MG, Soffredini R, D'Ambrosio R, Orsi E, De Nicola S, Degasperis E, Grancini V, Colombo M. Sustained virological response prevents development of insulin resistance in patients with chronic hepatitis C. *Hepatology.*; 56:1681-1687, 2012.
 263. Sambataro M, Perseghin G, Lattuada G, Beltramello G, Luzi L, Pacini G. Lipid accumulation in overweight type 2 diabetic subjects: relationships with insulin sensitivity and adipokines. *Acta Diabetol*; 50: 301-307, 2013.
 264. Spoladore R, Fragasso G, Perseghin G, De Cobelli F, Esposito A, Maranta F, Calori G, Locatelli M, Lattuada G, Scifo P, Del Maschio A, Margonato A. Beneficial effects of beta-blockers on left ventricular function and cellular energy reserve in patients with heart failure. *Fundam Clin Pharmacol*; 27: 455-464, 2013.
 265. Fragasso G, Spoladore R, Maranta F, Corti A, Lattuada G, Colombo B, Locatelli M, Salerno A, Calori G, Briceño L, Alfieri AB, Perseghin G, Margonato A. Increased low grade inflammation is associated with lack of functional response to Carvedilol in patients with systolic heart failure. *J Cardiovasc Med (Hagerstown)*; 14: 49-56, 2013.

266. Irace C, Carallo C, Scavelli FB, De Franceschi MS, Esposito T, Tripolino C, Gnasso A. Markers of insulin resistance and carotid atherosclerosis. A comparison of the homeostasis model assessment and triglyceride glucose index. *Int J Clin Pract*;67:665-672, 2013.
267. Irace C, Marini H, Bitto A, Altavilla D, Polito F, Adamo EB, Arcoraci V, Minutoli L, Di Benedetto A, Di Vieste G, de Gregorio C, Gnasso A, Corrao S, Licata G, Squadrito F. Genistein and endothelial function in postmenopausal women with metabolic syndrome. *Eur J Clin Invest*;43:1025-1031, 2013.
268. Bacci S, Prudente S, Copetti M, Spoto B, Rizza S, Baratta R, Di Pietro N, Morini E, Di Paola R, Testa A, Mallamaci F, Tripepi G, Zhang YY, Mercuri L, Di Silvestre S, Lauro R, Malatino L, Consoli A, Pellegrini F, Pandolfi A, Frittitta L, Zoccali C, Federici M, Doria A, Trischitta V. Joint effect of insulin signaling genes on cardiovascular events and on whole body and endothelial insulin resistance. *Atherosclerosis*. Jan;226(1):140-5, 2013.
269. Sciacqua A, Marini MA, Hribal ML, Perticone F, Sesti G. Association of insulin resistance indexes to carotid intima-media thickness. *PLoS One*. 8:e53968, 2013.
270. Marini MA, Frontoni S, Succurro E, Arturi F, Fiorentino TV, Sciacqua A, Hribal ML, Perticone F, Sesti G. Decreased insulin clearance in individuals with elevated 1-h post-load plasma glucose levels. *PLoS One* 8: e77440, 2013.
271. Sesti G, Fiorentino TV, Hribal ML, Sciacqua A, Perticone F. Association of hepatic insulin resistance indexes to nonalcoholic fatty liver disease and related biomarkers. *Nutr Metab Cardiovasc Dis* 23: 1182-1187, 2013.
272. Salani B, Marini C, Rio AD, Ravera S, Massollo M, Orengo AM, Amaro A, Passalacqua M, Maffioli S, Pfeiffer U, Cordera R, Maggi D, Sambuceti G. Metformin impairs glucose consumption and survival in Calu-1 cells by direct inhibition of hexokinase-II. *Sci Rep*.;3:2070, 2013.
273. Mazzucchelli C, Bordone C, Maggi D, Cordera R. Comments on ORIGIN trial. *Nutr Metab Cardiovasc Dis*. 23(8):e33-4, 2013.
274. Hamoudane M, Maffioli S, Cordera R, Maggi D, Salani B. Caveolin-1 and polymerase I and transcript release factor: new players in insulin-like growth factor-I receptor signaling. *J Endocrinol Invest*. 36(3):204-8, 2013.
275. Trombetta M., Boselli L., Cretti A., Cali A., Vettore M., Caruso B., Dorizzi R., Avogaro A., Muggeo M., Bonora E., Bonadonna R.C. Type 2 diabetes mellitus: a disease of the governance of the glucose-insulin system. An experimental metabolic

- control analysis study. *Nutr. Metab. Cardiovasc. Dis.* 23: 23-30, 2013.
276. Kiechl S, Wittmann J, Giaccari A, Knoflach M, Willeit P, Bozec A, Moschen AR, Muscogiuri G, Sorice GP, Kireva T, Summerer M, Wirtz S, Luther J, Mielenz D, Billmeier U, Egger G, Mayr A, Oberhollenzer F, Kronenberg F, Orthofer M, Penninger JM, Meigs JB, Bonora E, Tilg H, Willeit J, Schett G. Blockade of receptor activator of nuclear factor- κ B (RANKL) signaling improves hepatic insulin resistance and prevents development of diabetes mellitus. *Nat. Med.* 19:358-63, 2013.
 277. Moghetti P, Tosi F, Bonin C, Di Sarra D, Fiers T, Kaufman JM, Giagulli VA, Signori C, Zambotti F, Dall'Alda M, Spiazzi G, Zanolin ME, Bonora E. Divergences in insulin resistance between the different phenotypes of the polycystic ovary syndrome. *J. Clin. Endocrinol. Metab.* 98:E628-37, 2013.
 278. Di Sarra D, Tosi F, Bonin C, Fiers T, Kaufman JM, Signori C, Zambotti F, Dall'Alda M, Caruso B, Zanolin ME, Bonora E, Moghetti P. Metabolic inflexibility is a feature of women with polycystic ovary syndrome and is associated with both insulin resistance and hyperandrogenism. *J. Clin. Endocrinol. Metab.* 98: 2581-2588, 2013.
 279. Grasso V, Colombo C, Favalli V, Galderisi A, Rabbone I, Gombos S, Bonora E, Massa O, Meschi F, Cerutti F, Iafusco D, Bonfanti R, Monciotti C, Barbetti F. Six cases with severe insulin resistance (SIR) associated with mutations of insulin receptor: Is a Bartter-like syndrome a feature of congenital SIR *Acta Diabetol.* 50: 951-957, 2013.
 280. Rizzo M, Tyndall EK, Frontoni S, Jacoangeli F, Sarlo F, Panebianco F, Mistorni A, Di Renzo L, Calafiore R, Luca G, De Lorenzo A. Rapid and easy assessment of insulin resistance contributes to early detection of polycystic ovary syndrome. *J Endocrinol Invest.*,36:527-30, 2013.
 281. Bianchi C, Miccoli R, Trombetta M, Giorgino F, Frontoni S, Faloia E, Marchesini G, Dolci MA, Cavalot F, Cavallo G, Leonetti F, Bonadonna RC, Del Prato S; GENFIEV Investigators. Elevated 1-hour postload plasma glucose levels identify subjects with normal glucose tolerance but impaired β -cell function, insulin resistance, and worse cardiovascular risk profile: the GENFIEV study. *J Clin Endocrinol Metab.*,98:2100-5, 2013.
 282. Gilardini L, Lombardi C, Redaelli G, Vallone L, Faini , Mattaliano P, Parati G, Invitti C Glucose tolerance and weight loss in obese women with obstructive sleep

apnea. PLoS One.;8:e61382, 2013.

- 283.** Di Castelnuovo A, Di Pietro N, Di Tomo P, Di Silvestre S, Pipino C, Nenna G, Bonomini M, Iacoviello L, Pandolfi A. Metabolic syndrome in survivors from the 2009 earthquake in Italy. *Nutr Metab Cardiovasc Dis* 23(1):e5-8, 2013.
- 284.** Chandler-Laney P.C., Higgins P.B., Granger W., Alvarez J., Casazza K., Fernandez J.R., Dalla Man C., Cobelli C., Gower B.A.: Use of a simple liquid meal test to evaluate insulin sensitivity and beta-cell function in children. *Pediatric Obesity* 9:102-10, 2013.
- 285.** Squadrito F, Marini H, Bitto A, Altavilla D, Polito F, Adamo EB, D'Anna R, Arcoraci V, Burnett BP, Minutoli L, Di Benedetto A, Di Vieste G, Cucinotta D, de Gregorio C, Russo S, Corrado F, Saitta A, Irace C, Corrao S, Licata G. Genistein in the metabolic syndrome: results of a randomized clinical trial. *J Clin Endocrinol Metab.*;98:3366-3374, 2013.
- 286.** Marini MA, Frontoni S, Succurro E, Arturi F, Sciacqua A, Hribal ML, Perticone F, Sesti G. Insulin sensitivity, and β -cell function in relation to hemoglobin A1C. *Nutr Metab Cardiovasc Dis* 24: 27-33, 2014.
- 287.** Degasperi E, Valenti L, Aghemo A, De Francesco R, Rumi M, Soffredini R, Donnici L, Cheroni C, Fargion S, Zanoni V, Orsi E, Colombo M. Interleukin 28B genotype and insulin resistance in chronic Hepatitis C patients. *Antivir Ther.* 19:747-753, 2014.
- 288.** Anderwald CH, Tura A, Gessl A, Smajis S, Bieglmayer C, Marculescu R, Luger A, Pacini G, Krebs M. Whole-body insulin sensitivity rather than body-mass-index determines fasting and post-glucose-load growth hormone concentrations. *PLoS One.*17;9(12):e115184, 2014.
- 289.** Salani B, Del Rio A, Marini C, Sambuceti G, Cordera R, Maggi D. Metformin, cancer and glucose metabolism. *Endocr Relat Cancer.* 21(6):R461-71, 2014.
- 290.** Perticone M, Maio R, Tassone EJ, Tripepi G, Di Cello S, Miceli S, Caroleo B, Sciacqua A, Licata A, Sesti G, Perticone F. Insulin-resistance HCV infection-related affects vascular stiffness in normotensives. *Atherosclerosis* 238:108-112, 2015.
- 291.** Ciresi A, Amato MC, Giordano C. Reduction in insulin sensitivity and inadequate β -cell capacity to counteract the increase in insulin resistance in children with idiopathic growth hormone deficiency during 12 months of growth hormone treatment. *J Endocrinol Invest.*38:351-9, 2015.

