

1. Franzetti I, De Nale A, Bossi AC, Greco M, Morricone L, Ruggerini M, Meazza D, Sciolla A, Caviezel F, Oriani G. Epidural Spinal Electrostimulatory System (ESES) in the management of diabetic foot and peripheral arteriopathies. *Pacing Clin Electrophysiol* 12, 705-708, 1989.
2. Faglia E; Mantero M; Caminiti M; Caravaggi C; De Giglio R; Pritelli C; Clerici G; Fratino P; De Cata P; Dalla Paola L; Mariani G; Poli M; Settembrini PG; Sciangula L; Morabito A; Graziani L. Extensive use of peripheral angioplasty, particularly infrapopliteal, in the treatment of ischaemic diabetic foot ulcers: clinical results of a multicentric study of 221 consecutive diabetic subjects. *J Intern Med*, 252(3):225-32, 2002.
3. Dalla Paola L; Cogo A; Deanesi W; Stocchiero C; Colletta VC. Using hyaluronic acid derivates and cultured autologous fibroblasts and keratinocytes in a lower limb wound in a patient with diabetes: a case report. *Ostomy Wound Manage*, vol. 48(9) pp.46-9, 2002.
4. Dalla Paola L; Faglia E; Caminiti M; Clerici G; Ninkovic S; Deanesi V. Ulcer recurrence following first ray amputation in diabetic patients: a cohort prospective study. *Diabetes Care*, 26(6) pp. 1874-8, 2003.
5. Caravaggi C; De Giglio R; Pritelli C; Sommaria M; Dalla Noce S; Faglia E; Mantero M; Clerici G; Fratino P; Dalla Paola L; Mariani G; Mingardi R; Morabito A. HYAFF 11-based autologous dermal and epidermal grafts in the treatment of noninfected diabetic plantar and dorsal foot ulcers: a prospective, multicenter, controlled, randomized clinical trial. *Diabetes Care* 26(10) pp. 2853-9, 2003.
6. Saldalamacchia G, Lapice E, Cuomo V, De Feo E, D'Agostino E, Rivelles AA, Vaccaro O. A controlled study of the use of autologous platelet gel for the treatment of diabetic foot ulcers. *Nutr Metab Cardiovasc Dis*. 14(6):395-6, 2004.
7. van Schie CH, Vermigli C, Carrington AL, Boulton A. Muscle weakness and foot deformities in diabetes: relationship to neuropathy and foot ulceration in caucasian diabetic men. *Diabetes Care*. 27(7):1668-73, 2004.
8. Faglia E; Dalla Paola L; Clerici G; Clerissi J; Graziani L; Fusaro M; Gabrielli L; Losa S; Stella A; Gargiulo M; mantero M; Caminiti S; Curci V; Morabito A. Peripheral angioplasty as the first-choice revascularization procedure in diabetic patients with critical limb ischemia: prospective study of 993 consecutive patients hospitalized and followed between 1999 nd 2003. *Eur J Vasc Endovasc Surg*, 29(6) pp.620-7, 2005.

9. Giacco F, Perruolo G, D'Agostino E, Fratellanza G, Perna E, Misso S, Saldalamacchia G, Oriente F, Fiory F, Miele C, Formisano S, Beguinot F, Formisano P. Thrombin-activated platelets induce proliferation of human skin fibroblasts by stimulating autocrine production of insulin-like growth factor-1. *FASEB J.* 20(13):2402-4, 2006.
10. Biondi-Zoccai GG; Agostoni P; Sangiorgi G; Dalla Paola L; Armano F; Nicolini S; Alek J; Fusaro M. Mastering the antegrade femoral artery access in patients with symptomatic lower limb ischemia: learning curve, complications, and technical tips and tricks. *Catheter Cardiovasc Interv.* 68(6) pp 835-42, 2006.
11. Fusaro M; Dalla Paola L; Biondi-Zoccai GG. Retrograde posterior tibial artery access for below-the-knee percutaneous revascularization by means of sheathless approach and double wire technique. *Minerva cardioangiol.* 54(6) pp 773-7, 2006.
12. Dalla Paola L; Faglia E. Treatment of diabetic foot ulcer: an overview. Strategies for clinical approach. *Curr Diabetes Rev.* 2(4) 431-47, 2006.
13. Krishnan, S.T. Quattrini, C. Jeziorska, M. Malik, R.A. Rayman, G. Neurovascular factors in wound healing in the foot skin of type 2 diabetic subjects. *Diabetes Care.* 30(12):3058-62, 2007.
14. Fusaro M; Dalla Paola L; Biondi-Zoccai G. Pedal-plantar loop technique for a challenging below-the-knee chronic total occlusion: a novel approach to percutaneous revascularization in critical lower limb ischemia. *J Invasive Cardiol.* 19(2):E34-7, 2007.
15. Dalla Paola L; Volpe A; Varotto D; Postorino A; Brocco E; Senesi A; Merico M; De Vido D; Da Ros; Assaloni R. Use of a retrograde nail for ankle arthrodesis in Charcot neuroarthropathy: a limb salvage procedure. *Foot Ankle Int.* 28(9) :967-70, 2007.
16. Fusaro M; Dalla Paola L; Brigato C; Marangotto M; Nicolini S; Rripay R; Biondi Zoccai GG. Plantar to dorsalis pedis artery subintimal angioplasty in a patient with critical foot ischemia: a novel technique in the armamentarium of the peripheral interventionist. *J Cardiovasc Med (Hagerstown)*, vol.8(11) pp.977-80, 2007.
17. Dalla Paola L. Vacuum assisted closure: recommendations for use. A consensus document. *Int Wound J* vol 5 Suppl 4 pp iii-19, 2008.
18. Manzi M; Fusaro M; Ceccacci T; Erente G; Dalla Paola L; Brocco E. Clinical results of below-the knee intervention using pedal-plantar loop technique for the revascularization of foot arteries. *J Cardiovasc Surg (Torino)*, vol. 50(3) pp.331-7, 2009.

19. Dalla Paola L; Brocco E; Ceccacci T; Ninkovic S; Sorgentone S; Marinescu MG; Volpe A. Limb salvage in Charcot foot and ankle osteomyelitis: combined use single stage/double stage of arthrodesis and external fixation. *Foot Ankle Int*, 30(11) pp. 1065-70, 2009.
20. Uccioli L, Giurato L, Ruotolo V, Ciavarella A, Grimaldi MS, Piaggese A, Teobaldi I, Ricci L, Scionti L, Vermigli C, Seguro R, Mancini L, Ghirlanda G. Two-step autologous grafting using HYAFF scaffolds in treating difficult diabetic foot ulcers: results of a multicenter, randomized controlled clinical trial with long-term follow-up. *Int J Low Extrem Wounds* 10(2): 80-5, 2011.
21. Tolva V, Casana R, Guy Bianchi P., Bertoni G., Cireni L.V., Invitti C. Percutaneous transluminal angioplasty improves quality of life and glucose control in patients with critical limb ischemia. *Eur Rev Med Pharmacol Sci.*,16:2082-7, 2012.
22. Ciccone MM, Marchese A, Generali A, Loiodice C, Cortese F, Carbonara R, Scicchitano P, Laviola L, Giorgino F. Interventional therapy in diabetic foot: risk factors, clinical events and prognosis at one year follow-up (a study of 103 cases). *Pak J Biol Sci.*; 15:789-794, 2012.
23. Mancini M, Di Donato O, Saldalamacchia G, Liuzzi R, Rivellese A, Salvatore M. Contrast-enhanced ultrasound evaluation of peripheral microcirculation in diabetic patients: effects of cigarette smoking. *Radiol Med*. 118(2):206-14, 2013.
24. Dalla Paola L. Diabetic foot wounds: the value of negative pressure wound therapy with instillation. *Int Wound J*, vol. 10 Suppl 1 pp 25-31, 2013.
25. Dalla Paola L. Confronting a dramatic situation: the Charcot foot complicated by osteomyelitis. *Int J Low Extrem Wounds*, vol.13(4) pp.247-62, 2014.
26. Aiello A, Anichini R, Brocco E, Caravaggi C, Chiavetta A, Cioni R, Da Ros R, De Feo ME, Ferraresi R, Florio F, Gargiulo M, Galzerano G, Gandini R, Giurato L, Graziani L, Mancini L, Manzi M, Modugno P, Setacci C, Uccioli L; Italian Society of Diabetes.; Italian Society of Radiology.; Italian Society of Vascular Endovascular Surgery. Treatment of peripheral arterial disease in diabetes: a consensus of the Italian Societies of Diabetes (SID, AMD), Radiology (SIRM) and Vascular Endovascular Surgery (SICVE). *Nutr Metab Cardiovasc Dis*. 24(4):355-69, 2014.
27. Dalla Paola L, Carone A, Morisi C, Cardillo S, Pattavina M. Conservative surgical treatment of infected ulceration of the first metatarsophalangeal joint with osteomyelitis in diabetic patients. *J Foot Ankle Surg* 54:536-40, 2015.

28. Mantovani A, Trombetta M, Imbriaco C, Rigolon R, Mingolla L, Zamboni F, Dal Molin F, Cioccoloni D, Sanga V, Bruti M, Brocco E, Conti M, Ravenna G, Perrone F, Stoico V, Bonora E. Diabetic foot complicated by vertebral osteomyelitis and epidural abscess. *Endocrinol Diabetes Metab Case Rep* 2016:150132, 2016.
29. Troisi N, Turini F, Chisci E, Ercolini L, Frosini P, Lombardi R, Falciani F, Baggione C, Anichini R, Michelagnoli S. Pedal arch patency and not direct-angiosome revascularization predicts outcomes of endovascular interventions in diabetic patients with critical limb ischemia. *Int Angiol.* 36(5):438-444, 2017.
30. Francia P, Anichini R, Seghieri G, De Bellis A, Gulisano M. History, prevalence and assessment of limited joint mobility: from stiff hand syndrome to diabetic foot ulcer prevention. *Curr Diabetes Rev.* 14(5):411-426, 2018.
31. Mantovani A, Teobaldi I, Stoico V, Perrone F, Zannoni M, Cima L, Bruti M, Mingolla L, Trombetta M, Bonora E. Cutaneous squamous carcinoma in a patient with diabetic foot: an unusual evolution of a frequent complication. *Endocrinol Diabetes Metab Case Rep.* 31;2018, 2018.
32. Scarton A, Guiotto A, Malaquias T, Spolaor F, Sinigaglia G, Cobelli C, Jonkers I, Sawacha Z. A methodological framework for detecting ulcers' risk in diabetic foot subjects by combining gait analysis, a new musculoskeletal foot model and a foot finite element model. *Gait Posture.* 60:279-285, 2018.
33. Teobaldi I, Stoico V, Perrone F, Bruti M, Bonora E, Mantovani A. Honey dressing on a leg ulcer with tendon exposure in a patient with type 2 diabetes. *Endocrinol Diabetes Metab Case Rep.* 2018:18-0177, 2018.
34. Martinelli N, Curci V, Quarantiello A, Saldalamacchia G. The benefits of antimicrobial photodynamic therapy with RLP068 in the management of diabetic foot ulcers. *Drugs Context,* 8: 212610, 2019.
35. Dalla Paola L, Cimaglia P, Carone A, Scavone G, Boscarino G, Bernucci D, Sbarzaglia P, Censi S, Ferrari R, Campo G. Limb salvage in diabetic patients with no-option critical limb ischemia: outcomes of a specialized center experience. *Diabet Foot Ankle,* 10(1):1696012, 2019.
36. Bonora BM, Cappellari R, Mazzucato M, Rigato M, Grasso M, Menegolo M, Bruttocao A, Avogaro A, Fadini GP. Stem cell mobilization with plerixafor and healing of diabetic ischemic wounds: A phase IIa, randomized, double-blind, placebo-controlled trial. *Stem Cells Transl Med.* 9(9):965-973, 2020.

37. Dalla Paola L, Cimaglia P, Carone A, Boscarino G, Scavone G. Use of Integra Dermal Regeneration Template for Limb Salvage in Diabetic Patients With No-Option Critical Limb Ischemia. *Int J Low Extrem Wounds*. 1534734620905741, 2020.
38. Frykberg RG, Wukich DK, Kavarthapu V, Zgonis T, Dalla Paola L; Board of the Association of Diabetic Foot Surgeons. Surgery for the diabetic foot: A key component of care. *Diabetes Metab Res Rev*. 36 Suppl 1:e3251, 2020.
39. Martelli E, Federici M. Venous distal bypass should be considered as the gold standard to finally attempt limb salvage in patients with severe critical limb ischemia and diabetic foot, or under dialysis. *Acta Diabetol*. 57(8):1019-1020, 2020.