

1. Vaccaro O, Rivellese AA, Riccardi G, Tutino L, Annuzzi G, Mancini M. Impaired Glucose Tolerance and Risk Factors for Atherosclerosis. *Arteriosclerosis*, 4(6):592-597, 1984.
2. Vaccaro O, Pauciullo P, Rubba P, Annuzzi G, Rivellese AA, Riccardi G, Mancini M. Peripheral arterial circulation in individuals with impaired glucose tolerance. *Diabetes Care*, 8(6):594-597, 1985.
3. Zavaroni I., Dall'Aglia E., Bonora E., Alpi O., Passeri M., Reaven G.M. Evidence that multiple risk factors for coronary artery disease exist in persons with abnormal glucose tolerance. *Am. J. Med.* 83: 609-612, 1987.
4. Cignarelli M, Cicinelli E, Corso M, Cospite MR, Garruti G, Tafaro E, Giorgino R, Schonauer S. Biophysical and endocrine-metabolic changes during menopausal hot flashes: increase in plasma free fatty acid and norepinephrine levels. *Gynec. Obst. Invest.* 27: 34-37, 1989.
5. Bellodi G, Valeria Manicardi V, Malavasi V, Veneri L, Bernini G, Bossini P, Distefano S, Magnanini G, Muratori L, Rossi G. Hyperglycemia and prognosis of acute myocardial infarction in patients without diabetes mellitus. *The American Journal of Cardiology* 64(14):885-8, 1989.
6. Zavaroni I., Bonora E., Pagliara M., Dall'Aglia E., Luchetti L., Buonanno G., Bonati P.A., Bergonzani M., Gnudi L., Passeri M., Reaven G.M. Risk factors for coronary artery disease in healthy persons with hyperinsulinemia and normal glucose tolerance. *N. Engl. J. Med.* 320: 702-706, 1989.
7. Reaven G.M., Zavaroni I., Bonora E., Pagliara M., Dall'Aglia E., Luchetti L., Buonanno G., Bonati P.A. Bergonzani M., Gnudi L., Passeri M. Hyperinsulinemia and coronary artery disease. *N. Engl. J. Med.* 321: 618, 1989 (letter).
8. Zavaroni I., Bonati P.A., Luchetti L., Bonora E., Buonanno G., Bergonzani M., Pagliara M., Gnudi L., Butturini L., Passeri M., Reaven G.M. Habitual leisure-time physical activity is associated with differences in various risk factors for coronary artery disease. *J. Intern. Med.* 226: 417-421, 1989.
9. Rivellese A, Mancini M. Population diets influencing cardiovascular risk. *CURRENT OPINION IN LIPIDOLOGY*, vol. 2, p. 16-19, 1991.
10. Bonora E., Zenere M., Branzi P., Bagnani M., Maggiulli L., Tosi F., Travia D., Cacciatori V., Querena M., Moghetti P., Micciolo R., Ferrari A., Muggeo M. Influence of body fat and its regional localization on risk factors for atherosclerosis in young men. *Am. J. Epidemiol.* 135: 1271-1278, 1992.

11. Pujia A, Gnasso A, Irace C, Colonna A, Mattioli PL. Common carotid arterial wall thickness in NIDDM subjects. *Diabetes Care*. 17:1330-1336, 1994.
12. Bertello P, Veglio F, Pinna G, Gurioli L, Molino P, Alban S, Chiandussi L. Plasma Endothelin in NIDDM Patients with and without complication. *Diabetes Care*. 17: 574-577, 1994.
13. Nannipieri M, Rizzo L, Rapuano A, Pilo A, Penno G, Navalesi R. Increased transcapillary escape rate of albumin in microalbuminuric type II diabetic patients. *Diabetes Care*. 18: 1-9, 1995.
14. Letizia C, d'Ambrosio C, De Ciocchis A, Scavo D, Pozzilli P Serum angiotensin-converting enzyme levels in patients with recent-onset insulin-dependent diabetes after one year of low-dose cyclosporin therapy. IMDIAB Study Group. *Int J Clin Pharmacol Res*. 15(5-6):209-13, 1995.
15. Bonora E., Targher G., Branzi P., Zenere M., Saggiani F., Zenti M.G., Travia D., Tonoli M., Muggeo M., Cigolini M. Cardiovascular risk profile in 38-yr and 18-yr-old men. Contribution of body fat content and regional fat distribution. *Int. J. Obesity* 20: 28-36, 1996.
16. Avogaro A, Crepaldi C, Piarulli F, Milan D, Valerio A, Pavan P, Sacerdoti D, Calabrò A, Macdonald I, Crepaldi G, Scognamiglio R, Tiengo A. The hemodynamic abnormalities in short-term insulin deficiency: the role of prostaglandin inhibition. *Diabetes*. 45(5):602-9, 1996.
17. Pandolfi A, Iacoviello L, Capani F, Vitacolonna E, Donati MB, Consoli A. Glucose and insulin independently reduce the fibrinolytic potential of human vascular smooth muscle cells in culture. *Diabetologia*. 39(12):1425-3, 1996.
18. Bonora E., Targher G., Zenere M.B., Saggiani F., Cacciatori V., Tosi F., Travia D., Zenti M.G., Branzi P., Santi L., Muggeo M. Relationship of uric acid concentration to cardiovascular risk factors in young men. Role of obesity and central fat distribution. The Verona Young Men Atherosclerosis Risk Factors Study. *Int. J. Obes*. 20: 975-980, 1996.
19. Bonora E., Targher G., Zenere M.B., Saggiani F., Cacciatori V., Tosi F., Travia D., Zenti M.G., Branzi P., Santi L., Muggeo M. Relationship of fasting insulin to cardiovascular risk factor is already present in young men. The Verona Young Men Atherosclerosis Risk Factor Study. *Europ. J. Clin. Invest*. 27: 248-254, 1997.
20. Bonora E., Tessari R., Micciolo R., Zenere M., Targher G., Padovani R., Falezza G., Muggeo M. Intimal-medial thickness of the carotid artery in nondiabetic and non-insulin-

- dependent diabetic subjects. Relationship with insulin resistance. *Diabetes Care* 20: 627-631, 1997.
21. Willeit J., Kiechl S., Egger G., Oberhollenzer M., Oberhollenzer F., Muggeo M., Poewe W., Bonora E. The role of insulin in age-related sex differences of cardiovascular risk profile and morbidity. *Atherosclerosis* 130: 183-189, 1997.
 22. Bonora E., Willeit J., Kiechl S., Oberhollenzer F., Egger G., Bonadonna R., Muggeo M. Relationship between insulin and carotid atherosclerosis in the general population. The Bruneck Study. *Stroke* 28: 1147-1152, 1997
 23. Nannipieri M, Penno G, Rizzo L, Pucci L, Bandinelli S, Mattei P, Taddei S, Salvetti A, Navalesi R. Transcapillary escape rate of albumin in type II diabetic patients. The relationship with microalbuminuria and hypertension. *Diabetes Care*. 20: 1019-26, 1997.
 24. Avogaro A, Piarulli F, Valerio A, Miola M, Calverì M, Pavan P, Vicini P, Cobelli C, Tiengo A, Calò L, Del Prato S. Forearm nitric oxide balance, vascular relaxation, and glucose metabolism in NIDDM patients. *Diabetes*. 46(6):1040-6, 1997.
 25. Pedrinelli R, Penno G, Dell'Omo G, Bandinelli S, Giorgi D, Di Bello V, Nannipieri M, Navalesi R, Mariani M. Transvascular and urinary leakage of albumin in atherosclerotic and hypertensive men. *Hypertension*. 32: 318-23, 1998.
 26. Bonora E., Willeit J., Kiechl S., Oberhollenzer F., Egger G., Bonadonna R., Muggeo M. U-shaped and J-shaped relationships between serum insulin and coronary heart disease in the general population. *Diabetes Care* 21: 221-230, 1998.
 27. Kiechl S., Willeit J., Rungger G., Egger G., Oberhollenzer F., Bonora E. for the Bruneck Study Group. Alcohol consumption and atherosclerosis: What is the relation? Prospective results from the Bruneck Study. *Stroke* 29: 900-907, 1998.
 28. Bonora E., Kiechl S., Willeit J., Oberhollenzer F., Egger G., Bonadonna R., Muggeo M. Plasma glucose within the normal range is not associated with carotid atherosclerosis. Prospective results in subjects with normal glucose tolerance from the Bruneck Study. *Diabetes Care* 22: 1339-1346, 1999.
 29. Avogaro A, Calò L, Piarulli F, Miola M, deKreutzenberg S, Maran A, Burlina A, Mingardi R, Tiengo A, Del Prato S. Effect of acute ketosis on the endothelial function of type 1 diabetic patients: the role of nitric oxide. *Diabetes*. 48(2):391-7, 1999.
 30. Bertolotto A, Bandinelli S, Ruocco L, Lo Faro A, Penno G, Navalesi R. More on the effect of atorvastatin on plasma fibrinogen levels in primary hypercholesterolemia. *Atherosclerosis*.; 143: 455-7, 1999.

31. Irace C, Carallo C, Crescenzo A, Motti C, De Franceschi MS, Mattioli PL, Gnasso A. NIDDM is associated with lower wall shear stress of the common carotid artery. *Diabetes*. 48:193-197, 1999.
32. Marfella R, Rossi F, Giugliano D. QTc dispersion, hyperglycemia, and hyperinsulinemia. *Circulation*. 100(25):e149, 1999.
33. Mazza A, Motti C, Nulli A, Marra G, Gnasso A, Pastore A, Federici G, Cortese C. Lack of association between carotid intima-media thickness and methylenetetrahydrofolate reductase gene polymorphism or serum homocysteine in non-insulin-dependent diabetes mellitus. *Metabolism*. 49:718-723, 2000.
34. Marfella R, Giugliano D. Stress hyperglycaemia and death after myocardial infarction. *Lancet*. 355(9215):1647, 2000.
35. Marfella R, Giugliano D. Hyperglycemia leads to ECG reevaluation. *Aging*;12(4):246-8, 2000.
36. Nicolucci A, Ciccarone E, Consoli A, Di Martino G, La Penna G, Latorre A, Pandolfi A, Vitacolonna E, Capani F. Relationship between patient practice-oriented knowledge and metabolic control in intensively treated Type 1 diabetic patients: results of the validation of the Knowledge and Practices Diabetes Questionnaire. *Diabetes Nutr Metab*. 13(5):276-83, 2000.
37. Bonora E., Kiechl S., Oberhollenzer F., Egger G., Bonadonna R.C., Muggeo M. Impaired glucose tolerance, type 2 diabetes mellitus and carotid atherosclerosis. Prospective results from the Bruneck Study. *Diabetologia* 43: 156-164, 2000.
38. Lugari R, Dell'Anna C, Ugolotti D, Dei Cas A, Barilli AL, Zandomenighi R, Marani B, Iotti M, Orlandini A, Gnudi A. Effect of nutrient ingestion on glucagon-like peptide 1 (7-36 amide) secretion in human type 1 and type 2 diabetes. *Horm Metab Res*. 32(10):424-8, 2000.
39. Willeit J., Kiechl S., Oberhollenzer F., Rungger G, Egger G., Bonora E., Mitterer M., Muggeo M. Distinct risk profiles of early and advanced atherosclerosis. Prospective results from the Bruneck Study. *Arterioscl. Thromb. Vasc. Biol*. 20: 529-537, 2000.
40. Kiechl S., Willeit J., Bonora E., Schwartz S., Quingbo X. No association between dehydroepiandrosterone-sulphate and development of atherosclerosis in a prospective population study (Bruneck Study). *Arterioscl. Thromb. Vasc. Biol*. 20: 1094-1100, 2000.
41. Pandolfi A, Cetrullo D, Polishuck R, Alberta MM, Calafiore A, Pellegrini G, Vitacolonna E, Capani F, Consoli A. Plasminogen activator inhibitor type 1 is increased in the arterial wall of type II diabetic subjects. *Arterioscler Thromb Vasc Biol*. 21(8):1378-82, 2001.

42. Kiechl S., Egger G., Mayr M., Wiedermann C.J., Bonora E. Oberhollenzer F., Muggeo M., Xu Q., Wick G., Poewe W., Willeit J. Chronic infections and the risk of carotid atherosclerosis. prospective results from a large population study. *Circulation* 103: 1064-1070, 2001.
43. Pedrinelli R, Dell'Omo G, Penno G, Mariani M. Non-diabetic microalbuminuria, endothelial dysfunction and cardiovascular disease. *Vasc Med.* 6: 257-64, 2001.
44. Brun E., Zoppini G., Zamboni C., Bonora E., Muggeo M. Glucose instability is associated with a high level of circulating P-selectin. *Diabetes Care* 24: 1685, 2001 (letter).
45. Lugari R, Ugolotti D, Dei Cas A, Barilli AL, Iotti M, Marani B, Orlandini A, Gnudi A, Zandomenighi R. Urinary excretion of glucagon-like peptide 1 (GLP-1) 7-36 amide in human type 2 (non-insulin-dependent) diabetes mellitus. *Horm Metab Res.* 33(9):568-71, 2001.
46. Bonora E., Muggeo M. Post-prandial blood glucose as a risk factor for cardiovascular disease in type 2 diabetes. The epidemiological evidence. *Diabetologia* 44: 2107-2114, 2001.
47. Bacci S., Russo A., Rauseo A, Fanelli R., Trischitta V. The role of Dobutamine stress echocardiography in detecting severe coronary artery disease in asymptomatic at high risk type 2 diabetic patients. *Diabetes Care* 25: 1659, 2002.
48. Bacci S. Villella M, Villella A, Lancialonga T, Grilli M, Rauseo A, Mastroianno S, De Cosmo S, Fanelli R, Trischitta V. Screening for silent myocardial ischaemia in type 2 diabetic patients with additional atherogenic risk factors: applicability and accuracy of the exercise stress test. *Eur J Endocrinol* 147: 649-54, 2002.
49. Arcaro G., Cretti A., Balzano S., Lecgi A., Muggeo M., Bonora E., Bonadonna R.C. Insulin causes endothelial dysfunction in humans. Sites and mechanisms. *Circulation* 105: 576-582, 2002.
50. Bonora E. Postprandial peaks as a risk factor for cardiovascular disease: epidemiological perspectives. *Int. J. Clin. Pract. Suppl* 129: 5-11, 2002.
51. Bonora E., Formentini G., Calcaterra F., Lombardi S., Marini F., Zenari L., Saggiani F., Poli M., Perbellini S., Raffaelli A., Cacciatori V., Santi L., Targher G., Bonadonna R.C., Muggeo M. HOMA-estimated insulin resistance is an independent predictor of cardiovascular disease in type 2 diabetic subjects. Prospective data from the Verona Diabetes Complications Study. *Diabetes Care* 25: 1135-1141, 2002.
52. Lugari R, Dei Cas A, Ugolotti D, Finardi L, Barilli AL, Ognibene C, Luciani A, Zandomenighi R, Gnudi A. Evidence for early impairment of glucagon-like peptide 1-

- induced insulin secretion in human type 2 (non insulin-dependent) diabetes. *Horm Metab Res.* 34(3):150-4, 2002.
- 53.** Esposito K, Marfella R, Giugliano D. Hyperglycemia and heart dysfunction: an oxidant mechanism contributing to heart failure in diabetes. *J Endocrinol Invest.* 25(5):485-8, 2002.
- 54.** Veglio M, Bruno G, Borra M, Macchia G, Bargero G, D' Errico N, Pagano G, Cavallo Perin P. Prevalence of increased QT interval duration and dispersion in Type 2 diabetic patients and its relationship with coronary heart disease: a population-based cohort. *J Int Med* 251: 317-24, 2002.
- 55.** Federici M, Menghini R, Mauriello A, Hribal ML, Ferrelli F, Lauro D, Sbraccia P, Spagnoli LG, Sesti G, Lauro R. Insulin-dependent activation of endothelial nitric oxide synthase is impaired by O-linked glycosylation modification of signaling proteins in human coronary endothelial cells. *Circulation* 106:466-472, 2002.
- 56.** Sciacqua A, Candigliota M, Ceravolo R, Scozzafava A, Sinopoli F, Corsonello A, Sesti G, Perticone F. Weight loss in combination with physical activity improves endothelial dysfunction in human obesity. *Diabetes Care* 26:1673–1678, 2003.
- 57.** Bacci S., Langialonga T., Villella M., Rauseo A., Fanelli R, Trischitta V. Heart rate response at positive exercise stress test predicts coronary artery disease and its severity in silent at "high risk" type 2 diabetic patients. *Diabetes Care*, 26:2698-9, 2003.
- 58.** Marfella R, Verza M, Giugliano D. Ventricular remodeling does not accompany the development of heart failure in diabetic patients after myocardial infarction. *Circulation.* 107(13):e85, 2003.
- 59.** Bacci S., Villella M., Villella A., Langialonga T., Grilli M., Rauseo A, Mastroianno S., De Cosmo S., Fanelli R, Trischitta V. Response to Letter to the Editor - Screening of silent myocardial ischaemia in type 2 diabetic patients with additional atherogenic risk factors: applicability and accuracy. *European Journal of Endocrinology*, 149: 465-466, 2003.
- 60.** Bonora E., Kiechl S., Willeit J., Oberhollenzer F., Egger G., Bonadonna R., Muggeo M. Carotid atherosclerosis and coronary heart disease in the Metabolic Syndrome. Prospective data from the Bruneck Study. *Diabetes Care* 26: 1251-1257, 2003.
- 61.** Lapolla A, Piarulli F, Sartore G, Rossetti C, Martano L, Carraro P, De Paoli M, Fedele D. Peripheral artery disease in type 2 diabetes: the role of fibrinolysis. *Thromb Haemost.* 89(1):91-6, 2003.
- 62.** Iacobellis G, Ribaudo MC, Assael F, Vecchi E, Tiberti C, Zappaterreno A, Di Mario U, Leonetti F. Echocardiographic epicardial adipose tissue is related to anthropometric and

- clinical parameters of metabolic syndrome: a new indicator of cardiovascular risk. *J Clin Endocrinol Metab.* 88(11):5163-8, 2003.
- 63.** Dei Cas L, Metra M, Nodari S, Dei Cas A, Gheorghiade M. Prevention and management of chronic heart failure in patients at risk. *Am J Cardiol.*;91(9A):10F-17F, 2003.
- 64.** Andreozzi F, Laratta E, Sciacqua A, Perticone F, Sesti G. Angiotensin II impairs the insulin signalling pathway promoting production of nitric oxide by inducing phosphorylation of IRS-1 on Ser³¹² and Ser⁶¹⁶ in human umbilical vein endothelial cells. *Circ Res* 94:1211-1218, 2004.
- 65.** Sasso FC, Carbonara O, Nasti R, Campana B, Marfella R, Torella M, Nappi G, Torella R, Cozzolino D. Glucose metabolism and coronary heart disease in patients with normal glucose tolerance. *JAMA.* 291(15):1857-63, 2004.
- 66.** Montalcini T, Emanuele V, Ceravolo R, Gorgone G, Sesti G, Perticone F, Pujia A. Relation of low bone mineral density and carotid atherosclerosis in postmenopausal women. *Am J Cardiol* 94: 266-269, 2004.
- 67.** Marfella R, Esposito K, Nappo F, Siniscalchi M, Sasso FC, Portoghese M, Di Marino MP, Baldi A, Cuzzocrea S, Di Filippo C, Barboso G, Baldi F, Rossi F, D'Amico M, Giugliano D. Expression of angiogenic factors during acute coronary syndromes in human type 2 diabetes. *Diabetes.* 53(9):2383-91, 2004.
- 68.** Bonora E., Targher G., Formentini G., Calcaterra F., Lombardi S., Marini F., Zenari L., Saggiani F., Poli M., Perbellini S., Raffaelli A., Gemma M.L., Santi L., Bonadonna R.C., Muggeo M. The Metabolic Syndrome is an independent predictor of cardiovascular disease in type 2 diabetic subjects. Prospective data from the Verona Diabetes Complications Study. *Diabet. Med.* 21: 52-58, 2004.
- 69.** Bruno G, Giunti S, Bargero G, Ferrero S, Pagano G, Cavallo Perin P. Sex-differences in prevalence of electrocardiographic left ventricular hypertrophy in type 2 diabetes. The Casale Monferrato Study. *Diabetic Med.* 21:823-28, 2004.
- 70.** Bacci S., Villella M., Villella A., Rauseo A., Langialonga T., Trischitta V. Usefulness of pulse pressure for detection of extent and severity of coronary artery disease in type 2 diabetic patients with silent myocardial ischemia at exercise stress test. *Diabetologia* 48:1238-9, 2005.
- 71.** Marfella R, Verza M, Coppola L. Myocardial infarction and hyperglycemia. *Am J Cardiol.* 96(10):1468, 2005.

72. Miccoli R, Bianchi C, Odoguardi L, Penno G, Caricato F, Giovannitti MG, Pucci L, Del Prato S. Prevalence of the metabolic syndrome among Italian adults according to ATP III definition. *Nutr Metab Cardiovasc Dis.*; 15: 250-254, 2005.
73. Rivellese AA. Diet and cardiovascular disease: beyond cholesterol. *NUTRITION METABOLISM AND CARDIOVASCULAR DISEASES*, vol. 15, p. 395-398, 2005
74. Bonora E. The Metabolic Syndrome and cardiovascular disease. *Ann. Med.* 38: 64-80, 2006.
75. Giunti S, Bruno G, Veglio M, Gruden G, Webb D, Livingstone S, Chaturvedi N, Fuller JH, Cavallo-Perin P, and the EURODIAB IDDM complications Study Group. Electrocardiographic left ventricular hypertrophy in type 1 diabetes. Prevalence and relation to coronary heart disease and cardiovascular risk factors: The Eurodiab IDDM Complications Study. *Diabetes Care* 28:2255-57, 2006.
76. Secchiero P, Corallini F, Pandolfi A, Consoli A, Candido R, Fabris B, Celeghini C, Capitani S, Zauli G. An increased osteoprotegerin serum release characterizes the early onset of diabetes mellitus and may contribute to endothelial cell dysfunction. *Am J Pathol.* 169(6):2236-44, 2006.
77. Marfella R, Cacciapuoti F, Siniscalchi M, Sasso FC, Marchese F, Cinone F, Musacchio E, Marfella MA, Ruggiero L, Chiorazzo G, Liberti D, Chiorazzo G, Nicoletti GF, Saron C, D'Andrea F, Ammendola C, Verza M, Coppola L. Effect of moderate red wine intake on cardiac prognosis after recent acute myocardial infarction of subjects with Type 2 diabetes mellitus. *Diabet Med.* 23(9):974-81, 2006.
78. Di Filippo C, Cuzzocrea S, Rossi F, Marfella R, D'Amico M. Oxidative stress as the leading cause of acute myocardial infarction in diabetics. *Cardiovasc Drug Rev.* 24(2):77-87, 2006.
79. Zoccali C, Maio R, Mallamaci F, Sesti G, Perticone F. Uric acid and endothelial dysfunction in essential hypertension. *J Am Soc Nephrol* 17: 1466–1471, 2006.
80. Montalcini T, Gorgone G, Gazzaruso C, Sesti G, Perticone F, Pujia A. Endogenous testosterone and endothelial function in postmenopausal women. *Coron Artery Dis.* 18:9-13, 2007.
81. Cardellini M, Marini MA, Frontoni S, Hribal ML, Andreozzi F, Perticone F, Federici M, Lauro D, Sesti G. Carotid artery intima-media thickness is associated with insulin-mediated glucose disposal in nondiabetic normotensive offspring of type 2 diabetic patients. *Am J Physiol Endocrinol Metab* 292: E347-E352, 2007.
82. Sesti G, Sciacqua A, Scozzafava A, Vatrano M, Angotti E, Borrello F, Ruberto C, Santillo E, Parlato G, Perticone F. Effects of growth hormone and insulin-like growth factor-1 on cardiac hypertrophy of hypertensive patients. *J Hypertens* 25: 471-477, 2007.

83. Montalcini T, Gorgone G, Gazzaruso C, Sesti G, Perticone F, Pujia A. Carotid atherosclerosis associated to metabolic syndrome but not BMI in healthy menopausal women. *Diabetes Res Clin Pract.* 76:378-382, 2007.
84. Montalcini T, Gorgone G, Gazzaruso C, Sesti G, Perticone F, Pujia A. Relation between serum uric acid and carotid intima-media thickness in healthy postmenopausal women. *Intern Emerg Med.* 2:19-23, 2007.
85. Montalcini T, Gorgone G, Gazzaruso C, Sesti G, Perticone F, Pujia A. Role of endogenous androgens on carotid atherosclerosis in non-obese postmenopausal women. *Nutr Metab Cardiovasc Dis* 17:705-711, 2007.
86. Fourth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (Constituted by representatives of nine societies and by invited experts), Graham I, Atar D, Borch-Johnsen K, Boysen G, Burell G, Cifkova R, Dallongeville J, De Backer G, Ebrahim S, Gjelsvik B, Herrmann-Lingen C, Hoes A, Humphries S, Knapton M, Perk J, Priori SG, Pyorala K, Reiner Z, Ruilope L, Sans-Menendez S, Scholte OP, Reimer W, Weissberg P, Wood D, Yarnell J, Zamorano JL; Other experts who contributed to parts of the guidelines:, Walma E, Fitzgerald T, Cooney MT, Dudina A; European Society of Cardiology (ESC) Committee for Practice Guidelines (CPG), Vahanian A, Camm J, De Caterina R, Dean V, Dickstein K, Funck-Brentano C, Filippatos G, Hellemans I, Kristensen SD, McGregor K, Sechtem U, Silber S, Tendera M, Widimsky P, Zamorano JL, Hellemans I, Altiner A, Bonora E, Durrington PN, Fagard R, Giampaoli S, Hemingway H, Hakansson J, Kjeldsen SE, Larsen ML, Mancia G, Manolis AJ, Orth-Gomer K, Pedersen T, Rayner M, Ryden L, Sammut M, Schneiderman N, Stalenhoef AF, Tokgözoğlu L, Wiklund O, Zampelas A. European guidelines on cardiovascular disease prevention in clinical practice: executive summary. *Eur. Heart J.* 19:2375-2414, 2007.
87. Graham I, Atar D, Borch-Johnsen K, Boysen G, Burell G, Cifkova R, Dallongeville J, De Backer G, Ebrahim S, Gjelsvik B, Herrmann-Lingen C, Hoes A, Humphries S, Knapton M, Perk J, Priori SG, Pyorala K, Reiner Z, Ruilope L, Sans-Menendez S, Scholte OP, Reimer WS, Weissberg P, Wood D, Yarnell J, Zamorano JL, Walma E, Fitzgerald T, Cooney MT, Dudina A, Vahanian A, Camm J, De Caterina R, Dean V, Dickstein K, Funck-Brentano C, Filippatos G, Hellemans I, Kristensen SD, McGregor K, Sechtem U, Silber S, Tendera M, Widimsky P, Zamorano JL, Altiner A, Bonora E, Durrington PN, Fagard R, Giampaoli S, Hemingway H, Hakansson J, Kjeldsen SE, Larsen ML, Mancia G, Manolis AJ, Orth-Gomer K, Pedersen T, Rayner M, Ryden L, Sammut M, Schneiderman N, Stalenhoef AF,

Tokgözüoğlu L, Wiklund O, Zampelas A; European Society of Cardiology (ESC); European Association for Cardiovascular Prevention and Rehabilitation (EACPR); Council on Cardiovascular Nursing; European Association for Study of Diabetes (EASD); International Diabetes Federation Europe (IDF-Europe); European Stroke Initiative (EUSI); International Society of Behavioural Medicine (ISBM); European Society of Hypertension (ESH); European Society of General Practice/Family Medicine (ESGP/FM/WONCA); European Heart Network (EHN). European guidelines on cardiovascular disease prevention in clinical practice: executive summary. Fourth Joint Task Force of the European Society of Cardiology and other societies on cardiovascular disease prevention in clinical practice (constituted by representatives of nine societies and by invited experts). *Eur. J. Cardiovasc. Prev. Rehabil.* 14 Suppl 2:E1-40, 2007.

88. Graham I, Atar D, Borch-Johnsen K, Boysen G, Burell G, Cifkova R, Dallongeville J, De Backer G, Ebrahim S, Gjelsvik B, Herrmann-Lingen C, Hoes A, Humphries S, Knapton M, Perk J, Priori SG, Pyörälä K, Reiner Z, Ruilope L, Sans-Menéndez S, Scholte OP, Reimer WS, Weissberg P, Wood D, Yarnell J, Zamorano JL, Walma E, Fitzgerald T, Cooney MT, Dudina A, Vahanian A, Camm J, De Caterina R, Dean V, Dickstein K, Funck-Brentano C, Filippatos G, Hellemans I, Kristensen SD, McGregor K, Sechtem U, Silber S, Tendera M, Widimsky P, Zamorano JL, Altiner A, Bonora E, Durrington PN, Fagard R, Giampaoli S, Hemingway H, Hakansson J, Kjeldsen SE, Larsen L, Mancina G, Manolis AJ, Orth-Gomer K, Pedersen T, Rayner M, Ryden L, Sammut M, Schneiderman N, Stalenhoef AF, Tokgözüoğlu L, Wiklund O, Zampelas A; European Society of Cardiology (ESC); European Association for Cardiovascular Prevention and Rehabilitation (EACPR); Council on Cardiovascular Nursing; European Association for Study of Diabetes (EASD); International Diabetes Federation Europe (IDF-Europe); European Stroke Initiative (EUSI); Society of Behavioural Medicine (ISBM); European Society of Hypertension (ESH); WONCA Europe (European Society of General Practice/Family Medicine); European Heart Network (EHN); European Atherosclerosis Society (EAS). European guidelines on cardiovascular disease prevention in clinical practice: full text. Fourth Joint Task Force of the European Society of Cardiology and other societies on cardiovascular disease prevention in clinical practice (constituted by representatives of nine societies and by invited experts). *Eur. J. Cardiovasc. Prev. Rehabil.* 14: Suppl 2:S1-113, 2007.
89. Giunti S, Bruno G, Lillaz E, Gruden G, Lolli V, Chaturvedi N, Fuller JH, Veglio M, Cavallo-Perin P, and The EURODIAB IDDM Complications Study Group. Incidence and

- risk factors of prolonged QTc interval in type 1 diabetes: the Eurodiab Prospective Complications Study. *Diabetes Care*.30:2057-63, 2007.
90. Pandolfi A, De Filippis EA. Chronic hyperglycemia and nitric oxide bioavailability play a pivotal role in pro-atherogenic vascular modifications. *Genes Nutr*. 2(2):195-208, 2007.
 91. Di Cori A, Di Bello V, Miccoli R, Talini E, Palagi C, Delle Donne MG, Penno G, Nardi C, Bianchi C, Mariani M, Del Prato S, Balbarini A. Left ventricular function in normotensive young adults with well-controlled type 1 diabetes mellitus. *Am J Cardiol*. 99: 84-90, 2007.
 92. Bianchi C, Penno G, Pancani F, Civitelli A, Piaggese A, Caricato F, Pellegrini G, Del Prato S, Miccoli R. Non-traditional cardiovascular risk factors contribute to peripheral arterial disease in patients with type 2 diabetes. *Diabetes Res Clin Pract*.; 78: 246-253, 2007.
 93. Formoso G, De Filippis EA, Michetti N, Di Fulvio P, Pandolfi A, Bucciarelli T, Ciabattini G, Nicolucci A, Davì G, Consoli A. Decreased in vivo oxidative stress and decreased platelet activation following metformin treatment in newly diagnosed type 2 diabetic subjects. *Diabetes Metab Res Rev*. 24(3):231-7, 2008.
 94. Tesauro M, Schinzari F, Rovella V, Melina D, Mores N, Barini A, Mettimano M, Lauro D, Iantorno M, Quon MJ, Cardillo C. Tumor necrosis factor-alpha antagonism improves vasodilation during hyperinsulinemia in metabolic syndrome. *Diabetes Care*. 31(7):1439-41, 2008.
 95. Irace C, Tschakovsky ME, Carallo C, Cortese C, Gnasso A. Endothelial dysfunction or dysfunctions? Identification of three different FMD responses in males with type 2 diabetes. *Atherosclerosis*;200:439-445, 2008.
 96. Bianchi C, Penno G, Malloggi L, Barontini R, Corfini M, Giovannitti MG, Di Cianni G, Del Prato S, Miccoli R. Non-traditional markers of atherosclerosis potentiate the risk of coronary heart disease in patients with type 2 diabetes and metabolic syndrome. *Nutr Metab Cardiovasc Dis*.; 18: 31-38, 2008.
 97. Pattoneri P, Sozzi FB, Catellani E, Piazza A, Iotti R, Michelini M, Goldoni M, Borghetti A, Cappellini MD, Manicardi V. Myocardial involvement during the early course of type 2 diabetes mellitus: usefulness of myocardial performance index. *Cardiovascular Ultrasound*. 6:27, 2008.
 98. Bianchi C, Miccoli R, Penno G, Del Prato S. Primary prevention of cardiovascular disease in people with dysglycemia. *Diabetes Care*. 31 Suppl 2: S208-14, 2008.
 99. Montalcini T, Gorgone G, Gazzaruso C, Sesti G, Perticone F, Pujia A. Large brachial and common carotid artery diameter in postmenopausal women with carotid atherosclerosis. *Atherosclerosis* 196: 443–448, 2008.

100. Perticone F, Sciacqua A, Maio R, Perticone M, Laino I, Bruni R, Cello SD, Leone GG, Greco L, Andreozzi F, Sesti G. Renal function predicts cardiovascular outcomes in southern Italian postmenopausal women. *Eur J Cardiovasc Prev Rehabil.* 16: 481-486, 2009.
101. Bianchi C, Penno G, Del Prato S, Miccoli R. Serum gamma-glutamyltransferase is not associated to peripheral arterial disease in type 2 diabetic patients: Cross-sectional findings. *Atherosclerosis.* 203: 49-50, 2009.
102. Dell'Omo G, Penno G, Del Prato S, Mariani M, Pedrinelli R. Dysglycaemia in non-diabetic hypertensive patients: comparison of the impact of two different classifications of impaired fasting glucose on the cardiovascular risk profile. *J Hum Hypertens.* 23: 332-338, 2009.
103. Panero F, Novelli G, Zucco C, Fornengo P, Perotto M, Segre O, Grassi G, Cavallo-Perin P, Bruno G. Fasting plasma C-peptide and micro- and macrovascular complications in a large clinic-based cohort of type 1 diabetic people. *Diabetes Care.* 32:301-05, 2009.
104. Burt D, Bruno G, Chaturvedi N, Schalkwijk C, Stehouwer CD, Witte DR, Fuller JH, Pinach S, Cavallo Perin P, Gruden G. Anti-HSP27 Antibody Levels and Diabetic Complications in the EURODIAB Prospective Complications Study. *Diabetes Care* 32:1269-71, 2009.
105. Cacciapuoti F, Marfella R, Paolisso G, Cacciapuoti F. Is the aging heart similar to the diabetic heart? Evaluation of LV function of the aging heart with Tissue Doppler Imaging. *Aging Clin Exp Res.* 21(1):22-6, 2009.
106. Lamacchia O, Pinnelli S, Camarchio D, Fariello S, Gesualdo L, Stallone G, Cignarelli M. Waist-to-height ratio is the best anthropometric index in association with adverse cardiorenal outcomes in type 2 diabetes mellitus patients. *Am J Nephrol.*;29(6):615-9, 2009.
107. Marfella R, Di Filippo C, Portoghese M, Ferraraccio F, Rizzo MR, Siniscalchi M, Musacchio E, D'Amico M, Rossi F, Paolisso G. Tight glycemic control reduces heart inflammation and remodeling during acute myocardial infarction in hyperglycemic patients. *J Am Coll Cardiol.* 53(16):1425-36, 2009.
108. Marfella R, Di Filippo C, Portoghese M, Barbieri M, Ferraraccio F, Siniscalchi M, Cacciapuoti F, Rossi F, D'Amico M, Paolisso G. Myocardial lipid accumulation in patients with pressure-overloaded heart and metabolic syndrome. *J Lipid Res.* 50(11):2314-23, 2009.
109. Gruden G, Bruno G, Chaturvedi N, Burt D, Pinach S, Schalkwijk C, Stehouwer CD, Witte DR, Fuller JH, Cavallo Perin P, EURODIAB Prospective Complications Study Group. Anti-Hsp60 and anti-Hsp70 antibody levels and micro/macrovascular complications

- in Type 1 Diabetes: The Eurodiab Prospective Complications Study. *J Intern Med.* 266:527-36, 2009.
- 110.** Clodi M, Resl M, Stelzeneder D, Pacini G, Tura A, Mörtl D, Struck J, Morgenthaler NG, Bergmann A, Riedl M, Anderwald-Stadler M, Luger A, Pacher R, Hülsmann M. Interactions of glucose metabolism and chronic heart failure. *Exp Clin Endocrinol Diabetes.* 117(3):99-106, 2009.
- 111.** Piemonti L, Calori G, Lattuada G, Mercalli A, Ragogna F, Garancini MP, Ruotolo G, Luzi L, Perseghin G. Association between plasma monocyte chemoattractant protein-1 concentration and CVD mortality in middle-aged diabetic and non-diabetic individuals. *Diabetes Care.* 32(11):2105-10, 2009.
- 112.** Home PD, Pocock SJ, Beck-Nielsen H, Curtis PS, Gomis R, Hanefeld M, Jones NP, Komajda M, McMurray JJ; RECORD Study Team. Rosiglitazone evaluated for cardiovascular outcomes in oral agent combination therapy for type 2 diabetes (RECORD): a multicentre, randomised, open-label trial. *Lancet.* 373(9681):2125-35, 2009.
- 113.** Mazzucco S, Bifari F, Trombetta M, Guidi GC, Mazzi M, Anzola GP, Rizzuto N, Bonadonna R Evaluating endothelial function of the common carotid artery: an in vivo human model. *Nutr Metab Cardiovasc Dis.* 19(3):205-10, 2009.
- 114.** Targher G., Pichiri I., Zoppini G., Bonora E., Conchol M. Hemostatic and fibrinolytic abnormalities in endocrine diseases:a narrative review. *Sem. Thromb. Hemost.* 35: 605-612, 2009.
- 115.** Pontiroli AE, Morabito A. More than one target to prevent cardiovascular (CV) mortality in type 2 diabetes? *Nutr Metab Cardiovasc Dis.* 20(4): e13-4, 2010.
- 116.** . Giacco R., Clemente G, Cipriano D., Luongo D, Viscovo D., Patti L., Di Marino L., Giacco A, Naviglio D., Bianchi MA., Ciati R., Brighenti F, Rivellese AA., Riccardi G. Effects of the regular consumption of wholemeal wheat foods on cardiovascular risk factors in healthy people. *NUTRITION METABOLISM AND CARDIOVASCULAR DISEASES*, vol. 20, p. 186-194, 2010.
- 117.** Succurro E, Arturi F, Lugarà M, Grembiale A, Fiorentino TV, Caruso V, Andreozzi F, Sciacqua A, Hribal ML, Perticone F, Sesti G. One-hour postload plasma glucose levels are associated with kidney dysfunction. *Clin J Am Soc Nephrol* 5: 1922-1927, 2010.
- 118.** Amato, M.C., Giordano, C., Galia, M., Criscimanna, A., Vitabile, S., Midiri, M., Galluzzo, A. Visceral adiposity index: A reliable indicator of visceral fat function associated with cardiometabolic risk. *Diabetes Care* 33 (4) : 9, 2010.
- 119.** Giacco R., Clemente G, Cipriano D, Luongo D, Viscovo D., Patti L., Di Marino L,

- Giacco A., Naviglio D., Bianchi MA., Ciati R, Brighenti F., Rivellese AA., Riccardi G. Effects of the regular consumption of wholemeal wheat foods on cardiovascular risk factors in healthy people. *NUTRITION METABOLISM AND CARDIOVASCULAR DISEASES*, 20:186-194, 2010.
120. Rivellese A.A, Riccardi G., Vaccaro O. Cardiovascular risk in women with diabetes. *NUTRITION METABOLISM AND CARDIOVASCULAR DISEASES*, vol. 20, p. 474-480, 2010.
 121. Amato, M.C., Verghi, M., Nucera, M., Galluzzo, A., Giordano, C Low estradiol-to-testosterone ratio is associated with oligo-anovulatory cycles and atherogenic lipidic pattern in women with polycystic ovary syndrome. *Gynecological Endocrinology* 27 (8) :579-586, 2011.
 122. Amato, M.C., Giordano, C., Pitrone, M., Galluzzo, A Cut-off points of the visceral adiposity index (VAI) identifying a visceral adipose dysfunction associated with cardiometabolic risk in a Caucasian Sicilian population.. *Lipids in Health and Disease* 10:183, 2011.
 123. Tesouro M, Canale MP, Rodia G, Di Daniele N, Lauro D, Scuteri A, Cardillo C. Metabolic syndrome, chronic kidney, and cardiovascular diseases: role of adipokines. *Cardiol Res Pract.* 2011:653182, 2011.
 124. Presta I, Tassone EJ, Andreozzi F, Perticone M, Sciacqua A, Laino I, Musca D, Martino F, Sesti G, Perticone F. Angiotensin II type 1 receptor, but no type 2 receptor, interferes with the insulin-induced nitric oxide production in HUVECs. *Atherosclerosis* 219:463-467, 2011.
 125. Bonora E., Kiechl S., Zoppini G., Targher G., Meigs J.B., Willeit J., Oberhollenzer F., Bonadonna R.C., Muggeo M. Impact of reference category and number of traits in the cluster on risk of coronary heart disease in metabolic syndrome: prospective data from the Bruneck Study. *Metab. Syndr. Relat. Disord.* 9: 313-318, 2011.
 126. Dei Cas A, Spigoni V, Ardigò D, Pedrazzi G, Franzini L, Derlindati E, Urbani S, Monti L, Gnudi L, Zavaroni I. Reduced circulating endothelial progenitor cell number in healthy young adult hyperinsulinemic men. *Nutr Metab Cardiovasc Dis.* 21(7):512-7, 2011.
 127. Rivellese AA, Piatti P M. Consensus on: Screening and therapy of coronary heart disease in diabetic patients. *NUTRITION METABOLISM AND CARDIOVASCULAR DISEASES*, vol. 10, p. 757-764, 2011.
 128. Cignarelli M, Lamacchia O, Gesualdo L, Pilotti A, Pacilli A, Santini SA, De Cosmo S. Increased cardiovascular risk among type 2 diabetic patients with high-normal

- albuminuria and no evidence of kidney impairment. *Nutr Metab Cardiovasc Dis.* 21(2):e5-6, 2011.
- 129.** Di Flaviani A, Picconi F, Di Stefano P, Giordani I, Malandrucco I, Maggio P, Palazzo P, Sgreccia F, Peraldo C, Farina F, Frajese G, Frontoni S. Impact of glycemic and blood pressure variability on surrogate measures of cardiovascular outcomes in type 2 diabetic patients. *Diabetes Care.*,34:1605-9, 2011.
- 130.** Bianchi C, Miccoli R, Bonadonna RC, Giorgino F, Frontoni S, Faloia E, Marchesini G, Dolci MA, Cavalot F, Cavallo GM, Leonetti F, Del Prato S; GENFIEV Investigators. Pathogenetic mechanisms and cardiovascular risk: differences between HbA(1c) and oral glucose tolerance test for the diagnosis of glucose tolerance. *Diabetes Care.* 35:2607-12, 2012.
- 131.** Rizza S, Copetti M, Cardellini M, Porzio O, Luzi A, Pecchioli C, Martelli E, Valentini A, Ippoliti A, Romeo F, Pellegrini F, Lauro D, Lauro R, Federici M. Atherosclerosis severity but not undiagnosed diabetes predicts new cardiovascular events of subjects in secondary cardiovascular prevention. *Atherosclerosis.* 223(2):448-53, 2012.
- 132.** Perseghin G., Lattuada G, De Cobelli F., Esposito A., Canu T., Ragogna F., Maffi P., Scifo P., Secchi A, Del Maschio A., Luzi L. Left ventricular function and energy homeostasis in patients with type 1 diabetes with and without microvascular complications. *International Journal of Cardiology*, vol. 154, p. 111-115, 2012.
- 133.** Marfella R, Sasso FC, Cacciapuoti F, Portoghese M, Rizzo MR, Siniscalchi M, Carbonara O, Ferraraccio F, Torella M, Petrella A, Balestrieri ML, Stiuso P, Nappi G, Paolisso G. Tight glycemic control may increase regenerative potential of myocardium during acute infarction. *J Clin Endocrinol Metab.* 97(3):933-42, 2012.
- 134.** Pontiroli AE. Cardiovascular events after bariatric surgery. *JAMA.* 307(15):1577; author reply 1577-8, 2012.
- 135.** Marfella R, Sasso FC, Siniscalchi M, Paolisso P, Rizzo MR, Ferraro F, Stabile E, Sorropago G, Calabrò P, Carbonara O, Cinquegrana G, Piscione F, Ruocco A, D'Andrea D, Rapacciuolo A, Petronella P, Bresciani A, Rubino P, Mauro C, Paolisso G. Peri-Procedural Tight Glycemic Control during Early Percutaneous Coronary Intervention Is Associated with a Lower Rate of In-Stent Restenosis in Patients with Acute ST-Elevation Myocardial Infarction. *J Clin Endocrinol Metab.* 97(8):2862-71, 2012.
- 136.** Picconi F, Di Flaviani A, Malandrucco I, Giordani I, Frontoni S. Impact of glycemic variability on cardiovascular outcomes beyond glycated hemoglobin. Evidence and clinical perspectives. *Nutr Metab Cardiovasc Dis.*,22:691-6, 2012.

137. Giordani I, Picconi F, Malandrucchio I, Frontoni S. Comment on: Kromhout et al. n-3 fatty acids, ventricular arrhythmia-related events, and fatal myocardial infarction in postmyocardial infarction patients with diabetes. *Diabetes Care* 2011;34:2515-2520. *Diabetes Care*,35:e45, 2012.
138. Bruno RM, Penno G, Daniele G, Pucci L, Lucchesi D, Stea F, Landini L, Cartoni G, Taddei S, Ghiadoni L, Del Prato S. Type 2 diabetes mellitus worsens arterial stiffness in hypertensive patients through endothelial dysfunction. *Diabetologia*. 55: 1847-1855, 2012.
139. Franzini L, Ardigo D, Cavalot F, Miccoli R, Rivellese A, Trovati M, Zavaroni I, Vaccaro O. Women show worse control of type 2 diabetes and cardiovascular disease risk factors than men: Results from the MIND.IT Study Group of the Italian Society of Diabetology. *NMCD. NUTRITION METABOLISM AND CARDIOVASCULAR DISEASES*, p. 00-000, 2012.
140. Bruno RM, Ghiadoni L, Penno G, Taddei S, Del Prato S. Might hypovitaminosis D aggravate endothelial dysfunction-related increases in arterial stiffness seen in patients with hypertension and type 2 diabetes? Reply to Boucher BJ [letter]. *Diabetologia*.; 55: 3142-3143, 2012.
141. Marini MA, Succurro E, Castaldo E, Cufone S, Arturi F, Sciacqua A, Lauro R, Hribal ML, Perticone F, Sesti G. Cardiometabolic risk profiles and carotid atherosclerosis in individuals with prediabetes identified by fasting glucose, postchallenge glucose, and hemoglobin A1c criteria. *Diabetes Care* 35:1144–1149, 2012
142. Sciacqua A, Grillo N, Quero M, Sesti G, Perticone F. Asymmetric dimethylarginine plasma levels and endothelial function in newly diagnosed type 2 diabetic patients. *Int J Mol Sci*. 13:13804-15, 2012.
143. . Panero F, Gruden G, Perotto M, Fornengo P, Barutta F, Greco E, Runzo C, Ghezzo G, Cavallo-Perin P, Bruno G. Uric acid is not an independent predictor of cardiovascular mortality in type 2 diabetes: a population-based study. *Atherosclerosis* 221:183-188, 2012.
144. Gruden G, Giunti S, Barutta F, Chaturvedi N, Witte DR, Tricarico M, Fuller JH, Cavallo PerinP, Bruno G. QTc interval prolongation is independently associated with severe hypoglycemic attacks in type 1 diabetes from the EURODIAB IDDM Complications Study. *Diabetes Care*. 35:125-7, 2012.
145. Giunti S, Gruden G, Fornengo P, Barutta F, Amione C, Grezzo G, Cavallo-Perin P, Bruno G. Increased QT interval dispersion predicts 15-years cardiovascular mortality in type 2 diabetic subjects. The population-based Casale Monferrato Study. *Diabetes Care*;35:581-3, 2012.

146. Napoli N, Zardi E, Strollo R, Arigliani M, Daverio A, Olearo F, Tosi D, Dicuonzo G, Scarpa F, Pedone C, Tegue Simo HH, Mottini G, Pozzilli P. Increased carotid thickness in subjects with recently-diagnosed diabetes from rural Cameroon. *PLoS One*. 7(8):e41316, 2012.
147. Gruden G, Barutta F, Chaturvedi N, Schalkwijk C, Stehouwer CD, Pinach S, Manzo M, Loiacono M, Tricarico M, Mengozzi G, Witte DR, Fuller JH, Cavallo Perin P, Bruno G. N-terminal Pro-Brain Natriuretic Peptide is associated with diabetic complications in the EURODIAB Prospective Complications Study: the role of Tumor Necrosis Factor- α . *Diabetes Care* 35:1931-36, 2012.
148. Nodari S, Manerba A, Vaccari A, Milesi G, Carubelli V, Lazzarini V, Lombardi C, Etti F, Metra M, Dei Cas A. Six-year prognosis of diabetic patients with coronary artery disease. *Eur J Clin Invest*. 42(4):376-83, 2012.
149. Spigoni V, Picconi A, Cito M, Ridolfi V, Bonomini S, Casali C, Zavaroni I, Gnudi L, Metra M, Dei Cas A. Pioglitazone improves in vitro viability and function of endothelial progenitor cells from individuals with impaired glucose tolerance. *Plos One* 7(11):e48283, 2012.
150. Amato MC, Guarnotta V, Giordano C. Body composition assessment for the definition of cardiometabolic risk. *J Endocrinol Invest*. 36(7):537-43, 2013.
151. Marfella R, Rizzo MR, Siniscalchi M, Paolisso P, Barbieri M, Sardù C, Savinelli A, Angelico N, Del Gaudio S, Esposito N, Rambaldi PF, D'Onofrio N, Mansi L, Mauro C, Paolisso G, Balestrieri ML. Peri-procedural tight glycemic control during early percutaneous coronary intervention up-regulates endothelial progenitor cell level and differentiation during acute ST-elevation myocardial infarction: effects on myocardial salvage. *Int J Cardiol*. 168(4):3954-62, 2013.
152. Di Pietro N, Di Tomo P, Di Silvestre S, Giardinelli A, Pipino C, Morabito C, Formoso G, Marigliò MA, Pandolfi A. Increased iNOS activity in vascular smooth muscle cells from diabetic rats: potential role of Ca(2+)/calmodulin-dependent protein kinase II δ 2 (CaMKII δ 2). *Atherosclerosis*. 226(1):88-94, 2013.
153. Perticone F, Maio R, Tassone JE, Perticone M, Pascale A, Sciacqua A, Sesti G. Interaction between uric acid and endothelial dysfunction predicts new onset of diabetes in hypertensive patients. *Int J Cardiol* 167: 232-236, 2013.
154. Marini MA, Frontoni S, Succurro E, Arturi F, Fiorentino TV, Sciacqua A, Hribal ML, Perticone F, Sesti G. Insulin clearance is associated with carotid artery intima-media thickness. *Atherosclerosis* 229: 453-458, 2013.

155. Montesano A, Luzi L, Senesi P, Mazzocchi N, Terruzzi I. Resveratrol promotes myogenesis and hypertrophy in murine myoblasts. *J Transl Med.* 11:310, 2013.
156. Menzaghi C., Bacci S., Salvemini L., Mendonca C., Palladino G., Fontana A, De Bonis C., Marucci A., Goheen E., Prudente S., Morini E., Rizza S., Kanagaki A, Fini G., Mangiacotti D., Federici M, De Cosmo S., Pellegrini F., Doria A., Trischitta V. Serum resistin, cardiovascular disease and all-cause mortality in patients with type 2 diabetes. *PLoS ONE* 8:e64729, 2013.
157. Soedamah-Muthu SS, Chaturvedi N, Fuller JH, Toeller M; EURODIAB Prospective Complications Study Group. Do European people with type 1 diabetes consume a high atherogenic diet? 7-year follow-up of the EURODIAB Prospective Complications Study. *Eur J Nutr.* 52:1701-1710, 2013.
158. Irace C, Carallo C, Scavelli FB, De Franceschi MS, Esposito T, Tripolino C, Gnasso A. Markers of insulin resistance and carotid atherosclerosis. A comparison of the homeostasis model assessment and triglyceride glucose index. *Int J Clin Pract.* 67:665-672, 2013.
159. Giorgino F, Leonardini A, Laviola L. Cardiovascular disease and glycemic control in type 2 diabetes: now that the dust is settling from large clinical trials. *Ann N Y Acad Sci.* 1281:36-50, 2013.
160. Cioffi G, Faggiano P, Lucci D, Maggioni AP, Manicardi V, Travaglini A, Girfoglio D, Masson S, Giorda CB, Velussi M, Di Lenarda A, Verdecchia P, Comaschi M. Left ventricular dysfunction and outcome at two-year follow-up in patients with type 2 diabetes: The DYDA study. *Diabetes research and clinical practice* 06, 2013
161. Beltramello G, Manicardi V, Mazzuoli F, Rivellese A. Trialogue Plus : Management of cardiovascular risk in hyperglycaemic/diabetic patients at hospital discharge. *Acta Diabetologica* 10, 2013.
162. Gruden G, Barutta F, Pinach S, Lorenzati B, Cavallo-Perin P, Giunti S, Bruno G. Circulating anti-HSP70 levels in nascent metabolic syndrome: the Casale Monferrato Study. *Cell Stress Chaperone* 18:353-7, 2013.
163. Penno G, Solini A, Bonora E, Fondelli C, Orsi E, Zerbini G, Trevisan R, Vedovato M, Gruden G, Laviola L, Nicolucci A, Pugliese G. Gender differences in cardiovascular disease risk factors, treatments and complications in patients with type 2 diabetes: the RIACE Italian multicentre study.; Renal Insufficiency And Cardiovascular Events (RIACE) study, group. *J Intern Med.* 274(2):176-91, 2013.
164. Palazzo P, Maggio P, Altavilla R, Di Flaviani A, Giordani I, Malandrucco I, Picconi

- F, Passarelli F, Pasqualetti P, Ercolani M, Vernieri F, Frontoni S. Cerebral hemodynamics and systemic endothelial function are already impaired in well-controlled type 2 diabetic patients, with short-term disease. *PLoS One*. 8:e83287, 2013.
165. Targher G, Mantovani A, Pichiri I, Rigolon R, Dauriz M, Zoppini G, Morani G, Vassanelli C, Bonora E. Non-alcoholic fatty liver disease is associated with an increased prevalence of atrial fibrillation in hospitalized patients with type 2 diabetes. *Clin. Sci. (Lond)* 125: 301-309, 2013.
 166. Mannucci E, Dicembrini I, Lauria A, Pozzilli P. Is glucose control important for prevention of cardiovascular disease in diabetes? *Diabetes Care*. 36Suppl 2:S259-63, 2013.
 167. Dei Cas A, Spigoni V, Franzini L, Preti M, Ardigò D, Derlindati E, Metra M, Monti LD, Dell'Era P, Gnudi L, Zavaroni I. Lower endothelial progenitor cell number, family history of cardiovascular disease and reduced HDL-cholesterol levels are associated with shorter leukocyte telomere length in healthy young adults. *Nutr Metab Cardiovasc Dis*. 23(3):272-85, 2013.
 168. Dei Cas A, Spigoni V, Ridolfi V, Metra M. Diabetes and chronic heart failure: from diabetic cardiomyopathy to therapeutic approach. *Endocr Metab Immune Disord Drug Targets*.13(1):38-5011, 2013.
 169. Bruno G, Landi A, Barutta F, Ghezzi G, Baldin C, Spadafora L, Schimmenti A, Prinzi T, Cavallo Perin P, Gruden G. NH2-terminal pro-brain natriuretic peptide is a stronger predictor of cardiovascular mortality than C-reactive protein and albumin excretion rate in elderly patients with type 2 diabetes: the Casale Monferrato population-based study. *Diabetes Care* 36:2677-82, 2013.
 170. Giordani I, Di Flaviani A, Picconi F, Malandrucchio I, Ylli D, Palazzo P, Altavilla R, Vernieri F, Passarelli F, Donno S, Lauro D, Pasqualetti P, Frontoni S. Acute hyperglycemia reduces cerebrovascular reactivity: the role of glycemic variability. *J Clin Endocrinol Metab*. 99:2854-60, 2014.
 171. Marfella R, Rizzo MR, Capoluongo MC, Ambrosino M, Savinelli A, Cinone F, Martinelli G, Fava I, Petrella A, Barbieri M, Paolisso G. Cryptogenic stroke and diabetes: a probable link between silent atrial fibrillation episodes and cerebrovascular disease. *Expert Rev Cardiovasc Ther*. 12(3):323-9, 2014.
 172. Menzaghi C, Xu M, Salvemini L, De Bonis C, Palladino G, Huang T, Copetti M, Zheng Y, Li Y, Fini G, Hu FB, Bacci S, Qi L, Trischitta V. Circulating adiponectin and cardiovascular mortality in patients with type 2 diabetes mellitus: evidence of sexual dimorphism. *Cardiovasc Diabetol*. 13: 130, 2014

173. Targher G., Zoppini G., Bonora E., Moghetti P. Hemostatic and fibrinolytic abnormalities in polycystic ovary syndrome. *Semin Thromb Hemost.* 40: 600-618, 2014.
174. Menini S, Iacobini C, Ricci C, Blasetti Fantauzzi C, Pugliese G. Protection from diabetes-induced atherosclerosis and renal disease by D-carnosine-octylester: effects of early versus late inhibition of advanced glycation end-products in Apoe-null mice. *Diabetologia*, 2014.
175. Zanetti M., Zenti M.G., Barazzoni R., Zardi F., Semolic A., Messa M.G., Mearelli F., Russi G., Fonda M., Scarano L., Bonora E., Cattin L. HELP LDL Apheresis Reduces Plasma Pentraxin 3 in Familial Hypercholesterolemia. *PLoS One.* 11;9(7):e101290, 2014.
176. Bruno G, Barutta F, Landi A, Civera S, Cavallo Perin P, Gruden G. The effect of age and NT-proBNP on the association of central obesity with 6-years cardiovascular mortality of diabetic people: the population-based Casale Monferrato study mortality in type 2 diabetes. *Plos One*, 9:e96076, 2014.
177. Gruden G, Landi A, Bruno G. Natriuretic peptides, heart, and adipose tissue: new findings and future developments for diabetes research. *Diabetes Care* 37:2899-908, 2014.
178. Spigoni V, Lombardi C, Cito M, Picconi A, Ridolfi V, Andreoli R, Anelli N, Gnudi L, Goldoni M, Zavaroni I, Raddino R, Dei Cas A. N-3 PUFA increase bioavailability and function of endothelial progenitor cells. *Food Funct.* 5(8):1881-90, 2014.
179. Bruno G, Barutta F, Landi A, Pinach S, Caropreso P, Mengozzi G, Baldassarre S, Fragapani S, Civera S, Cavallo Perin P, Gruden G. Levels of N-terminal pro brain natriuretic peptide are enhanced in people with the uncomplicated metabolic syndrome: a nested case-control analysis of the population-based Casale Monferrato Study. *Diabetes&Metabolism Research &Reviews*, 2014
180. De Cosmo S, Rossi MC, Pellegrini F, Lucisano G, Bacci S, Gentile S, Ceriello A, Russo G, Nicolucci A, Giorda C, Viazzi F, Pontremoli R; AMD-Annals Study Group. Kidney dysfunction and related cardiovascular risk factors among patients with type 2 diabetes. *Nephrol Dial Transplant* 29(3):657-62, 2014.
181. Di Fulvio P, Pandolfi A, Formoso G, Di Silvestre S, Di Tomo P, Giardinelli A, De Marco A, Di Pietro N, Taraborrelli M, Sancilio S, Di Pietro R, Piantelli M, Consoli A. Features of endothelial dysfunction in umbilical cord vessels of women with gestational diabetes. *Nutr Metab Cardiovasc Dis.* 24(12):1337-45, 2014.
182. Ambra R, Manca S, Palumbo MC, Leoni G, Natarelli L, De Marco A, Consoli A, Pandolfi A, Virgili F. Transcriptome analysis of human primary endothelial cells (HUVEC) from umbilical cords of gestational diabetic mothers reveals candidate sites for an epigenetic

- modulation of specific gene expression. *Genomics*. 103(5-6):337-48, 2014.
183. Isidori AM, Corona G, Aversa A, Gianfrilli D, Jannini EA, Foresta C, Maggi M, Lenzi A; SIAMS-ED Study Group The SIAMS-ED Trial: A National, Independent, Multicentre Study on Cardiometabolic and Hormonal Impairment of Men with Erectile Dysfunction Treated with Vardenafil. *Int J Endocrinol*. 858715, 2014.
 184. Pozzilli P, Strollo R, Bonora E. One size does not fit all glycemic targets for type 2 diabetes. *J Diabetes Investig*. 5(2):134-41, 2014.
 185. Soare A, Weiss EP, Pozzilli P. Benefits of caloric restriction for cardiometabolic health, including type 2 diabetes mellitus risk. *Diabetes Metab Res Rev*. 30Suppl 1:41-7, 2014.
 186. Perticone M, Miceli S, Maio R, Caroleo B, Sciacqua A, Tassone EJ, Greco L, Staltari O, Sesti G, Perticone F. Chronic HCV infection increases cardiac left ventricular mass index in normotensive patients. *J Hepatol*. 61:755-760, 2014
 187. Perticone F, Sciacqua A, Perticone M, Tassone EJ, Sesti G, Violi F, Lip GY. Clinical risk factors and subclinical target organ damage as predictors of new-onset of atrial fibrillation: The Catanzaro atrial fibrillation project. *Int J Cardiol* 177:666-668, 2014
 188. Sesti G, Fiorentino TV, Perticone M, Sciacqua A, Andreozzi F, Hribal ML, Perticone F Characterization of left ventricular mass in individuals at risk for type 2 diabetes identified by HbA1c levels according to the American Diabetes Association Criteria. *Int J Cardiol* 179: 211–213, 2015
 189. De Franceschi MS, Palange AL, Mancuso A, Grande L, Muccari D, Scavelli FB, Irace C, Gnasso A, Carallo C. Decreased platelet aggregation by shear stress-stimulated endothelial cells in vitro: Description of a method and first results in diabetes. *Diab Vasc Dis Res*.12:53-61, 2015.
 190. Galli D, Carubbi C, Masselli E, Corradi D, Dei Cas A, Nouvenne A, Bucci G, Arcari ML, Mirandola P, Vitale M, Gobbi G. PKC ϵ is a negative regulator of PVAT-derived vessel formation. *Exp Cell Res*. 330(2):277-86, 2015.
 191. Dei Cas A, Fonarow GC, Gheorghiade M, Butler J. Concomitant diabetes mellitus and heart failure. *Curr Probl Cardiol*.40(1):7-43, 2015.
 192. Di Somma C, Ciresi A, Amato MC, Savastano S, Savanelli MC, Scarano E, Colao A, Giordano C. Alteration of the growth hormone axis, visceral fat dysfunction, and early cardiometabolic risk in adults: the role of the visceral adiposity index. *Endocrine*. 49:492-502, 2015.
 193. Peeters SA, Engelen L, Buijs J, Chaturvedi N, Fuller JH, Schalkwijk CG, Stehouwer

- CD; EURODIAB Prospective Complications Study Group. Plasma levels of matrix metalloproteinase-2, -3, -10, and tissue inhibitor of metalloproteinase-1 are associated with vascular complications in patients with type 1 diabetes: the EURODIAB Prospective Complications Study. *Cardiovasc Diabetol.* 14:31, 2015.
194. Bozzetto L, Annuzzi G, Pacini G, Costabile G, Vetrani C, Vitale M, Griffo E, Giacco A, De Natale C, Cocozza S, Della Pepa G, Tura A, Riccardi G, Rivelles AA. Polyphenol-rich diets improve glucose metabolism in people at high cardiometabolic risk: a controlled randomised intervention trial. *Diabetologia.* 58(7):1551-60, 2015.
 195. Costabile G, Della Pepa G, Bozzetto L, Annuzzi G, Vetrani C, Giacco R, Della Corte G, Conte FS, Di Marino L, Rivelles AA. Urine 8-isoprostane in relation to adiposity and insulin resistance in individuals at high cardiometabolic risk. *Metab Syndr Relat Disord.* 13(4):187-91, 2015.
 196. Di Pino A, Urbano F, Piro S, Purrello F, Rabuazzo AM. Update on pre-diabetes: Focus on diagnostic criteria and cardiovascular risk. *World J Diabetes.* 15;7(18):423-432, 2016.
 197. Zoppini G, Bonapace S, Bergamini C, Rossi A, Trombetta M, Lanzoni L, Bertolini L, Zenari L, Bonora E, Targher G. Evidence of left atrial remodeling and left ventricular diastolic dysfunction in type 2 diabetes mellitus with preserved systolic function. *Nutr Metab Cardiovasc Dis.* 26(11):1026-1032, 2016.
 198. Mantovani A, Mingolla L, Rigolon R, Pichiri I, Cavalieri V, Zoppini G, Lippi G, Bonora E, Targher G. Nonalcoholic fatty liver disease is independently associated with an increased incidence of cardiovascular disease in adult patients with type 1 diabetes. *Int J Cardiol.* 15;225:387-391, 2016.
 199. Mantovani A, Rigamonti A, Bonapace S, Bolzan B, Pernigo M, Morani G, Franceschini L, Bergamini C, Bertolini L, Valbusa F, Rigolon R, Pichiri I, Zoppini G, Bonora E, Violi F, Targher G. Nonalcoholic Fatty Liver Disease Is Associated With Ventricular Arrhythmias in Patients With Type 2 Diabetes Referred for Clinically Indicated 24-Hour Holter Monitoring. *Diabetes Care.* 39(8):1416-23, 2016.
 200. Mantovani A, Rigolon R, Pichiri I, Pernigo M, Bergamini C, Zoppini G, Bonora E, Targher G. Hyperuricemia is associated with an increased prevalence of atrial fibrillation in hospitalized patients with type 2 diabetes. *J Endocrinol Invest.* 39(2):159-67, 2016.
 201. Gazzaruso C, Coppola A, Pujia A, Falcone C, Collaviti S, Fodaro M, Gallotti P, Solerte SB, Giustina A, Pelissero G, Luzi L, Montalcini T. Erectile dysfunction as a predictor of asymptomatic coronary artery disease in elderly men with type 2 diabetes. *J*

Geriatr Cardiol. 13(6):552-6, 2016.

202. Fiorentino TV, Sesti F, Andreozzi F, Pedace E, Sciacqua A, Hribal ML, Perticone F, Sesti G. One-hour post-load hyperglycemia combined with HbA1c identifies pre-diabetic individuals with a higher cardio-metabolic risk burden. *Atherosclerosis*. 253:61-69, 2016.
203. Biondi-Zoccai G, Pinto A, Versaci F, Procaccini E, Neri G, Sesti G, Uccioli L, Vetere M, Peruzzi M, Nudi F. Comparative Impact of Hypoglycemic Agents on Severity and Extent of Myocardial Ischemia in Patients With Type 2 Diabetes Mellitus Undergoing Myocardial Perfusion Scintigraphy. *J Cardiovasc Pharmacol*. 68(2):162-70, 2016.
204. Bonadonna RC, Borghi C, Consoli A, Volpe M. Novel antidiabetic drugs and cardiovascular risk: Primum non nocere. *Nutr Metab Cardiovasc Dis*. 26(9):759-66, 2016.
205. Arturi F, Succurro E, Miceli S, Cloro C, Ruffo M, Maio R, Perticone M, Sesti G, Perticone F. Liraglutide improves cardiac function in patients with type 2 diabetes and chronic heart failure. *Endocrine* 57(3):464-473, 2017.
206. Vitale M, Vaccaro O, Masulli M, Bonora E, Del Prato S, Giorda CB, Nicolucci A, Squatrito S, Auciello S, Babini AC, Bani L, Buzzetti R, Cannarsa E, Cignarelli M, Cigolini M, Clemente G, Coccozza S, Corsi L, D'Angelo F, Dall'Aglia E, Di Cianni G, Fontana L, Gregori G, Grioni S, Giordano C, Iannarelli R, Iovine C, Lapolla A, Lauro D, Laviola L, Mazzucchelli C, Signorini S, Tonutti L, Trevisan R, Zamboni C, Riccardi G, Rivellese AA; TOSCA.IT Study Group. Polyphenol intake and cardiovascular risk factors in a population with type 2 diabetes: The TOSCA.IT study. *Clin Nutr*. 36(6):1686-1692, 2017.
207. Kautzky-Willer A, Baggio G, Rossi MC, Lapolla A, Russo GT. Type 2 Diabetes and Cardiovascular Risk in Women 2016. *Int J Endocrinol*. 2017:6905697, 2017.
208. Mantovani A, Rigolon R, Pichiri I, Morani G, Bonapace S, Dugo C, Zoppini G, Bonora E, Targher G. Relation of elevated serum uric acid levels to first-degree heart block and other cardiac conduction defects in hospitalized patients with type 2 diabetes. *J Diabetes Complications*. 31(12):1691-1697, 2017.
209. Dauriz M, Mantovani A, Bonapace S, Verlato G, Zoppini G, Bonora E, Targher G. Prognostic Impact of Diabetes on Long-term Survival Outcomes in Patients With Heart Failure: A Meta-analysis. *Diabetes Care*. 40(11):1597-1605, 2017.
210. Mantovani A, Rigolon R, Pichiri I, Bonapace S, Morani G, Zoppini G, Bonora E, Targher G. Nonalcoholic fatty liver disease is associated with an increased risk of heart block in hospitalized patients with type 2 diabetes mellitus. *PLoS One*. 12(10):e0185459, 2017.
211. Zoppini G, Bergamini C, Bonapace S, Rossi A, Trombetta M, Mantovani A,

- Toffalini A, Lanzoni L, Bertolini L, Zenari L, Bonora E, Targher G. Association between subclinical left ventricular systolic dysfunction and glycemic control in asymptomatic type 2 diabetic patients with preserved left ventricular function. *J Diabetes Complications*. 31(6):1035-1040, 2017.
- 212.** De Gregorio C, Marini H, Alibrandi A, Di Benedetto A, Bitto A, Adamo EB, Altavilla D, Irace C, Di Vieste G, Pancaldo D, Granese R, Atteritano M, Corrao S, Licata G, Squadrito F, Arcoraci V. Genistein Supplementation and Cardiac Function in Postmenopausal Women with Metabolic Syndrome: Results from a Pilot Strain-Echo Study. *Nutrients*. 9(6), 2017.
- 213.** Marzona I, Avanzini F, Lucisano G, Tettamanti M, Baviera M, Nicolucci A, Roncaglioni MC; Risk & Prevention Collaborative Group. Are all people with diabetes and cardiovascular risk factors or microvascular complications at very high risk? Findings from the Risk and Prevention Study. *Acta Diabetol*. 54(2):123-131, 2017.
- 214.** Di Pino A, Mangiafico S, Urbano F, Scicali R, Scandura S, D'Agate V, Piro S, Tamburino C, Purrello F, Rabuazzo AM. HbA1c Identifies Subjects With Prediabetes and Subclinical Left Ventricular Diastolic Dysfunction. *J Clin Endocrinol Metab*. 1;102(10):3756-3764, 2017.
- 215.** Maddaloni E, Xia Y, Park K, D'Eon S, Tinsley LJ, St-Louis R, Khamaisi M, Li Q, King GL, Keenan HA. High density lipoprotein modulates osteocalcin expression in circulating monocytes: a potential protective mechanism for cardiovascular disease in type 1 diabetes. *Cardiovasc Diabetol*. 16(1):116, 2017.
- 216.** Ballotari P, Venturelli F, Greci M, Giorgi Rossi P, Manicardi V. Sex Differences in the Effect of Type 2 Diabetes on Major Cardiovascular Diseases: Results from a Population-Based Study in Italy. *Int J Endocrinol*. 2017:6039356, 2017.
- 217.** Sardu C, Barbieri M, Santamaria M, Giordano V, Sacra C, Paolisso P, Spirito A, Marfella R, Paolisso G, Rizzo MR. Multipolar pacing by cardiac resynchronization therapy with a defibrillators treatment in type 2 diabetes mellitus failing heart patients: impact on responders rate, and clinical outcomes. *Cardiovasc Diabetol*. 16(1):75, 2017.
- 218.** Policardo L, Seghieri G, Francesconi P, Anichini R, Franconi F, Del Prato S. Gender difference in diabetes related excess risk of cardiovascular events: When does the 'risk window' open? *J Diabetes Complications*. 31(1):74-79, 2017.
- 219.** Mattioli AV, Palmiero P, Manfrini O, Puddu PE, Nodari S, Dei Cas A, Mercurio G, Scrutinio D, Palermo P, Sciomer S, Di Francesco S, Novo G, Novo S, Pedretti RFE, Zito A, Parati G, Pedrinelli R, Farinetti A, Maiello M, Moscucci F, Tenaglia RL, Sucato V,

- Triggiani M, Cugusi L, Scicchitano P, Saba PS, Ciccone MM. Mediterranean diet impact on cardiovascular diseases: a narrative review. *J Cardiovasc Med (Hagerstown)*. 18(12):925-935, 2017.
220. Katsiki N, Purrello F, Tsioufis C, Mikhailidis DP. Cardiovascular disease prevention strategies for type 2 diabetes mellitus. *Expert Opin Pharmacother*. 18(12):1243-1260, 2017.
 221. Marfella R, Sardu C, Calabrò P, Siniscalchi M, Minicucci F, Signoriello G, Balestrieri ML, Mauro C, Rizzo MR, Paolisso G, Barbieri M. Non-ST-elevation myocardial infarction outcomes in patients with type 2 diabetes with non-obstructive coronary artery stenosis: Effects of incretin treatment. *Diabetes Obes Metab*. 20(3):723-729, 2018.
 222. Zoppini G, Bergamini C, Mantovani A, Dauriz M, Targher G, Rossi A, Bonora E. The E/e' ratio difference between subjects with type 2 diabetes and controls. A meta-analysis of clinical studies. *PLoS One*. 13(12):e0209794, 2018.
 223. Zenti MG, Altomari A, Lupo MG, Botta M, Bonora E, Corsini A, Ruscica M, Ferri N. From lipoprotein apheresis to proprotein convertase subtilisin/kexin type 9 inhibitors: Impact on low-density lipoprotein cholesterol and C-reactive protein levels in cardiovascular disease patients. *Eur J Prev Cardiol*. 25(17):1843-1851, 2018.
 224. Vitale M, Masulli M, Calabrese I, Rivellese AA, Bonora E, Signorini S, Perriello G, Squatrito S, Buzzetti R, Sartore G, Babini AC, Gregori G, Giordano C, Clemente G, Grioni S, Dolce P, Riccardi G, Vaccaro O; TOSCA.IT Study Group. Impact of a Mediterranean Dietary Pattern and Its Components on Cardiovascular Risk Factors, Glucose Control, and Body Weight in People with Type 2 Diabetes: A Real-Life Study. *Nutrients*. 10;10(8), 2018.
 225. Zoppini G, Bergamini C, Bonapace S, Trombetta M, Mantovani A, Toffalini A, Lanzoni L, Bertolini L, Zenari L, Bonora E, Targher G, Rossi A. Left ventricular chamber dilation and filling pressure may help to categorise patients with type 2 diabetes. *BMJ Open Diabetes Res Care*. 6(1):e000529, 2018.
 226. Saglimbene V, Palmer SC, Ruospo M, Natale P, Maione A, Nicolucci A, Vecchio M, Tognoni G, Craig JC, Pellegrini F, Lucisano G, Hegbrant J, Ariano R, Lamacchia O, Sasso A, Morano S, Filardi T, De Cosmo S, Pugliese G, Procaccini DA, Gesualdo L, Palasciano G, Johnson DW, Tonelli M, Strippoli GFM; Long-Term Impact of RAS Inhibition on Cardiorenal Outcomes (LIRICO) Investigators. The Long-Term Impact of Renin-Angiotensin System (RAS) Inhibition on Cardiorenal Outcomes (LIRICO): A Randomized, Controlled Trial. *J Am Soc Nephrol*. 29(12):2890-2899, 2018.
 227. Cavallari I, Delli Veneri A, Maddaloni E, Melfi R, Patti G, Napoli N, Pozzilli P, Di Sciascio G. Comparison of Lipid-Lowering Medications and Risk for Cardiovascular

- Disease in Diabetes. *Curr Diab Rep.* 18(12):138, 2018.
- 228.** Sasso FC, Rinaldi L, Lascar N, Marrone A, Pafundi PC, Adinolfi LE, Marfella R. Role of Tight Glycemic Control during Acute Coronary Syndrome on CV Outcome in Type 2 Diabetes. *J Diabetes Res.* 2018:3106056, 2018.
- 229.** Nusca A, Tuccinardi D, Albano M, Cavallaro C, Ricottini E, Manfrini S, Pozzilli P, Di Sciascio G. Glycemic variability in the development of cardiovascular complications in diabetes. *Diabetes Metab Res Rev.* 34(8):e3047, 2018.
- 230.** Cavallari I, Delli Veneri A, Maddaloni E, Melfi R, Patti G, Napoli N, Pozzilli P, Di Sciascio G. Comparison of Lipid-Lowering Medications and Risk for Cardiovascular Disease in Diabetes. *Curr Diab Rep.* 18(12):138, 2018.
- 231.** Grieco D, Palamà Z, Borrelli A, De Ruvo E, Sciarra L, Scarà A, Goanta E, Calabrese V, Pozzilli P, Di Sciascio G, Calò L. Diabetes mellitus and atrial remodelling in patients with paroxysmal atrial fibrillation: Role of electroanatomical mapping and catheter ablation. *Diab Vasc Dis Res.* 15(3):185-195, 2018.
- 232.** D'Alessandro A, De Pergola G. The Mediterranean Diet: its definition and evaluation of a priori dietary indexes in primary cardiovascular prevention. *Int J Food Sci Nutr.* 69(6):647-659, 2018.
- 233.** Rizzo M, Nikolic D, Patti AM, Mannina C, Montalto G, McAdams BS, Rizvi AA, Cosentino F. GLP-1 receptor agonists and reduction of cardiometabolic risk: Potential underlying mechanisms. *Biochim Biophys Acta Mol Basis Dis.* 864(9 Pt B):2814-2821, 2018.
- 234.** De Rosa S, Arcidiacono B, Chiefari E, Brunetti A, Indolfi C, Foti DP. Type 2 Diabetes Mellitus and Cardiovascular Disease: Genetic and Epigenetic Links. *Front Endocrinol (Lausanne).* 9:2, 2018.
- 235.** Vetrani C, Vitale M, Bozzetto L, Della Pepa G, Coccozza S, Costabile G, Mangione A, Cipriano P, Annuzzi G, Rivellese AA. Association between different dietary polyphenol subclasses and the improvement in cardiometabolic risk factors: evidence from a randomized controlled clinical trial. *Acta Diabetol.* 55(2):149-153, 2018.
- 236.** Baldassarre MPA, Andersen A, Consoli A, Knop FK, Vilsbøll T. Cardiovascular biomarkers in clinical studies of type 2 diabetes. *Diabetes Obes Metab.* 20(6):1350-1360, 2018.
- 237.** Kappel BA, Federici M. Gut microbiome and cardiometabolic risk. *Rev Endocr Metab Disord.* 20(4):399-406, 2019.

- 238.** Castoldi G, di Gioia CRT, Roma F, Carletti R, Manzoni G, Stella A, Zerbini G, Perseghin G. Activation of angiotensin type 2 (AT2) receptors prevents myocardial hypertrophy in Zucker diabetic fatty rats. *Acta. Diabetol*, 2019 56:97-104, 2019.
- 239.** Maddaloni E, Coleman RL, Pozzilli P, Holman RR. Long-term risk of cardiovascular disease in individuals with latent autoimmune diabetes in adults (UKPDS 85). *Diabetes Obes. Metab*, 21:2115-2122, 2019.
- 240.** Nibali L, Donos N, Terranova V, Di Pino A, Di Marca S, Ferrara V, Pisano M, Scicali R, Rabuazzo AM, Purrello F, Malatino L. Left ventricular geometry and periodontitis in patients with the metabolic syndrome. *Clin. Oral. Investig*, 23:2695-2703, 2019.
- 241.** Fiorentino TV, Succurro E, Andreozzi F, Sciacqua A, Perticone F, Sesti G. One-hour post-load hyperglycemia combined with HbA1c identifies individuals with higher risk of cardiovascular diseases: Cross-sectional data from the CATAMERI study. *Diabetes Metab. Res. Rev*, 35:e3096, 2019.
- 242.** Zoppini G, Trombetta M, Sabbagh L, Dauriz M, Mantovani A, Bergamini C. Impaired Aortic Valve Growth in Type 1 Diabetes Mellitus. *Can. J. Cardiol*, 35:941.e5-941.e6, 2019.
- 243.** Zoppini G, Bergamini C, Trombetta M, Sabbagh L, Dauriz M, Mantovani A, Targher G, Fossà I, Rinaldi E, Bonora E. Increased aortic stiffness index in patients with type 1 diabetes without cardiovascular disease compared to controls. *J. Endocrinol. Invest*, 42:1109-1115, 2019.
- 244.** Morieri ML, Longato E, Mazzucato M, Di Camillo B, Cocchiglia A, Gubian L, Sparacino G, Avogaro A, Fadini GP, Vigili de Kreutzenberg S. Improved long-term cardiovascular outcomes after intensive versus standard screening of diabetic complications: an observational study. *Cardiovasc Diabetol*, 18(1):117, 2019.
- 245.** Rivellese AA, Ciciola P, Costabile G, Vetrani C, Vitale M. The Possible Role of Nutraceuticals in the Prevention of Cardiovascular Disease. *High Blood Press Cardiovasc. Prev*, 26:101-111, 2019.
- 246.** Milanese G, Silva M, Bruno L, Goldoni M, Benedetti G, Rossi E, Ferrari C, Grutta L, Maffei E, Toia P, Forte E, Bonadonna RC, Sverzellati N, Cademartiri F. Quantification of epicardial fat with cardiac CT angiography and association with cardiovascular risk factors in symptomatic patients: from the ALTER-BIO (Alternative Cardiovascular Bio-Imaging markers) registry. *Diagn Interv Radiol*, 25(1):35-41, 2019.

- 247.** Cardellini M, Rizza S, Casagrande V, Cardolini I, Ballanti M, Davato F, Porzio O, Canale MP, Legramante JM, Mavilio M, Menghini R, Martelli E, Farcomeni A, Federici M. Soluble ST2 is a biomarker for cardiovascular mortality related to abnormal glucose metabolism in high-risk subjects. *Acta Diabetol*, 56(3):273-280, 2019.
- 248.** Morieri ML, Avogaro A, Fadini GP; the DARWIN-T2D Network of the Italian Diabetes Society. Cholesterol lowering therapies and achievement of targets for primary and secondary cardiovascular prevention in type 2 diabetes: unmet needs in a large population of outpatients at specialist clinics. *Cardiovasc Diabetol*. 10;19(1):190, 2020.
- 249.** Boscari F, Morieri ML, Amato AML, Vallone V, Uliana A, Baritussio A, Vitturi N, Cipponeri E, Cavallin F, Avogaro A, Fadini GP, Bruttomesso D. Performance of the Steno type 1 risk engine for cardiovascular disease prediction in Italian patients with type 1 diabetes. *Nutr Metab Cardiovasc Dis* 30(10):1813-1819, 2020.
- 250.** Cosentino F, Grant PJ, Aboyans V, Bailey CJ, Ceriello A, Delgado V, Federici M, Filippatos G, Grobbee DE, Hansen TB, Huikuri HV, Johansson I, Jüni P, Lettino M, Marx N, Mellbin LG, Östgren CJ, Rocca B, Roffi M, Sattar N, Seferović PM, Sousa-Uva M, Valensi P, Wheeler DC; ESC Scientific Document Group. 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur Heart J*. 41(2):255-323, 2020.
- 251.** Avogaro A, Azzolina D, Fadini GP, Baldi I. Incidence of heart failure in patients with type 1 diabetes: a systematic review of observational studies. *J Endocrinol Invest*. [Epub ahead of print], 2020.
- 252.** Martini D, Rosi A, Tassotti M, Antonini M, Dall'Asta M, Bresciani L, Fantuzzi F, Spigoni V, Domínguez-Perles R, Angelino D, Ricci C, Del Pozo-Luengo S, Tornel PL, Scazzina F, Gil-Izquierdo A, Dei Cas A, Brighenti F, Bonadonna R, Del Rio D, Mena P. Effect of coffee and cocoa-based confectionery containing coffee on markers of cardiometabolic health: results from the pocket-4-life project. *Eur J Nutr* [Epub ahead of print], 2020.
- 253.** Giacco R, Costabile G, Fatati G, Frittitta L, Maiorino MI, Marelli G, Parillo M, Pistis D, Tubili C, Vetrani C, Vitale M. Effects of polyphenols on cardio-metabolic risk factors and risk of type 2 diabetes. A joint position statement of the Diabetes and Nutrition Study Group of the Italian Society of Diabetology (SID), the Italian Association of Dietetics and Clinical Nutrition (ADI) and the Italian Association of Medical Diabetologists (AMD). *Nutr Metab Cardiovasc Dis*. 30(3):355-367, 2020.
- 254.** Spaggiari G, Cignarelli A, Sansone A, Baldi M, Santi D. To beer or not to beer: A

- meta-analysis of the effects of beer consumption on cardiovascular health. *PLoS One*. 15(6):e0233619, 2020.
- 255.** Napoli R, Formoso G, Piro S, Targher G, Consoli A, Purrello F. Management of type 2 diabetes for prevention of cardiovascular disease. An expert opinion of the Italian Diabetes Society. *Nutr Metab Cardiovasc Dis*. 30(11):1926-1936, 2020.
- 256.** Cosentino F, Grant PJ, Aboyans V, Bailey CJ, Ceriello A, Delgado V, Federici M, Filippatos G, Grobbee DE, Hansen TB, Huikuri HV, Johansson I, Jüni P, Lettino M, Marx N, Mellbin LG, Östgren CJ, Rocca B, Roffi M, Sattar N, Seferović PM, Sousa-Uva M, Valensi P, Wheeler DC; ESC Scientific Document Group. 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur Heart J*. 41(2):255-323, 2020.
- 257.** Pastore I, Bolla AM, Montefusco L, Lunati ME, Rossi A, Assi E, Zuccotti GV, Fiorina P. The Impact of Diabetes Mellitus on Cardiovascular Risk Onset in Children and Adolescents. *Int J Mol Sci*. 21(14):4928, 2020.
- 258.** Sardu C, Paolisso P, Ducceschi V, Santamaria M, Sacra C, Massetti M, Ruocco A, Marfella R. Cardiac resynchronization therapy and its effects in patients with type 2 DIAbetes mellitus OPTimized in automatic vs. echo guided approach. Data from the DIA-OPTA investigators. *Cardiovasc Diabetol*. 19(1):202, 2020.
- 259.** Cavallari I, Maddaloni E, Pieralice S, Mulè MT, Buzzetti R, Ussia GP, Pozzilli P, Grigioni F. The Vicious Circle of Left Ventricular Dysfunction and Diabetes: From Pathophysiology to Emerging Treatments. *J Clin Endocrinol Metab*. 105(9):dgaa427, 2020.
- 260.** D'Onofrio L, Maddaloni E, Buzzetti R. Osteocalcin and sclerostin: Background characters or main actors in cardiovascular disease? *Diabetes Metab Res Rev*. 36(1):e3217, 2020.
- 261.** Cannistraci R, Mazzetti S, Mortara A, Perseghin G, Ciardullo S. Risk stratification tools for heart failure in the diabetes clinic. *Nutr Metab Cardiovasc Dis*. 30(7):1070-1079, 2020.
- 262.** Creta A, Providência R, Adragão P, de Asmundis C, Chun J, Chierchia G, Defaye P, Schmidt B, Anselme F, Finlay M, Hunter RJ, Papageorgiou N, Lambiase PD, Schilling RJ, Combes S, Combes N, Albenque JP, Pozzilli P, Boveda S. Impact of Type-2 Diabetes Mellitus on the Outcomes of Catheter Ablation of Atrial Fibrillation (European Observational Multicentre Study). *Am J Cardiol*. 125(6):901-906, 2020.
- 263.** Vaccaro O, Vitale M, Costanzo S, Masulli M, Riccardi G, De Curtis A, Donati MB, Rivellese AA, de Gaetano G, Iacoviello L. Cardiovascular risk factors control according to

diabetes status and prior cardiovascular events in patients managed in different settings. *Diabetes Res Clin Pract.* 168:108370, 2020.

264. Vitale M, Calabrese I, Massimino E, Shivappa N, Hebert JR, Auciello S, Grioni S, Krogh V, Sartore G, Signorini S, Rivellese AA, Riccardi G, Vaccaro O, Masulli M. Dietary inflammatory index score, glucose control and cardiovascular risk factors profile in people with type 2 diabetes. *Int J Food Sci Nutr.* 13:1-8, 2020.
265. Tecce N, Masulli M, Lupoli R, Della Pepa G, Bozzetto L, Palmisano L, Rivellese AA, Riccardi G, Capaldo B. Evaluation of cardiovascular risk in adults with type 1 diabetes: poor concordance between the 2019 ESC risk classification and 10-year cardiovascular risk prediction according to the Steno Type 1 Risk Engine. *Cardiovasc Diabetol.* 19(1):166, 2020.
266. Maranta F, Cianfanelli L, Rizzo M, Cianflone D. Filling the gap between Guidelines and Real World in the cardiovascular approach to the diabetic patients: the need for a call to action. *Int J Cardiol.* S0167-5273(20)34322-9, 2020.
267. Succurro E, Fiorentino TV, Miceli S, Perticone M, Sciacqua A, Andreozzi F, Sesti G. Relative Risk of Cardiovascular Disease Is Higher in Women With Type 2 Diabetes, but Not in Those With Prediabetes, as Compared With Men. *Diabetes Care.* 43(12):3070-3078, 2020.
268. Mancuso E, Perticone M, Spiga R, Averta C, Rubino M, Fiorentino TV, Miceli S, Mannino GC, Sciacqua A, Succurro E, Perticone F, Sesti G, Andreozzi F. Association between Serum Mg²⁺ Concentrations and Cardiovascular Organ Damage in a Cohort of Adult Subjects. *Nutrients.* 12(5):1264, 2020.
269. Succurro E, Pedace E, Andreozzi F, Papa A, Vizza P, Fiorentino TV, Perticone F, Veltri P, Cascini GL, Sesti G. Reduction in Global Myocardial Glucose Metabolism in Subjects With 1-Hour Postload Hyperglycemia and Impaired Glucose Tolerance. *Diabetes Care.* 43(3):669-676, 2020.
270. Targher G, Corey KE, Byrne CD. NAFLD, and cardiovascular and cardiac diseases: Factors influencing risk, prediction and treatment. *Diabetes Metab.* 6:101215, 2020.
271. Lonardo A, Targher G. Cardiovascular Risk in NAFLD: An Intimate Relationship? *Dig Dis Sci.* 65(6):1593-1595, 2020.
272. Henson JB, Roden M, Targher G, Corey KE. Is Nonalcoholic Fatty Liver Disease Not a Risk Factor for Cardiovascular Disease: Not Yet Time for a Change of Heart. *Hepatology.* 71(5):1867-1869, 2020.
273. Targher G, Byrne CD, Tilg H. NAFLD and increased risk of cardiovascular disease:

clinical associations, pathophysiological mechanisms and pharmacological implications. *Gut*. 69(9):1691-1705, 2020.

- 274.** Lombardi R, Airaghi L, Targher G, Serviddio G, Maffi G, Mantovani A, Maffei C, Colecchia A, Villani R, Rinaldi L, Orsi E, Pisano G, Adinolfi LE, Fargion S, Fracanzani AL. Liver fibrosis by FibroScan® independently of established cardiovascular risk parameters associates with macrovascular and microvascular complications in patients with type 2 diabetes. *Liver Int*. 40(2):347-354, 2020.